



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 11:29 pm GMT

PDB ID : 4P4E
Title : X-ray structure of human glutamate carboxypeptidase II (GCPII) in complex with a phosphoramidate inhibitor MP1D
Authors : Barinka, C.
Deposited on : 2014-03-12
Resolution : 1.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5252 (1.70-1.66)
Clashscore	112137	5803 (1.70-1.66)
Ramachandran outliers	110173	5704 (1.70-1.66)
Sidechain outliers	110143	5703 (1.70-1.66)
RSRZ outliers	101464	5298 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	707	<div> <div>6%</div> <div>87%</div> <div>10% ..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	802	-	-	-	X

2 Entry composition [i](#)


There are 10 unique types of molecules in this entry. The entry contains 6739 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	694	Total	C	N	O	S	0	75	0
			5928	3801	992	1109	26			

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula: C₆H₁₂O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		
3	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Zn	0	0
			2	2		


- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Cl	0	0
			1	1		

- Molecule 9 is N-[(S)-hydroxy(4-phenylbutoxy)phosphoryl]-L-glutamic acid (three-letter code: 2G4) (formula: C₁₅H₂₂NO₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	A	1	Total	C	N	O	P	0	0
			24	15	1	7	1		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	562	Total	O	0	0
			562	562		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glutamate carboxypeptidase 2



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	101.75Å 130.68Å 159.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.29 – 1.67 28.29 – 1.67	Depositor EDS
% Data completeness (in resolution range)	96.6 (28.29-1.67) 96.6 (28.29-1.67)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 1.67Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.154 , 0.183 0.154 , 0.183	Depositor DCC
R_{free} test set	1194 reflections (1.02%)	DCC
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.012	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6739	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, 2G4, NAG, CL, CA, FUC, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	2/6245 (0.0%)	0.83	2/8450 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	737	PHE	CD2-CE2	5.19	1.49	1.39
1	A	733	TYR	CD1-CE1	5.13	1.47	1.39

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	370	ARG	NE-CZ-NH1	6.60	123.60	120.30
1	A	673	ARG	NE-CZ-NH1	6.16	123.38	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5928	0	5782	59	1
2	A	168	0	147	1	0
3	A	20	0	20	0	0
4	A	22	0	19	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	11	0	10	0	0
6	A	2	0	0	0	0
7	A	1	0	0	0	0
8	A	1	0	0	0	0
9	A	24	0	19	0	0
10	A	562	0	0	13	2
All	All	6739	0	5997	60	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:651[B]:GLN:O	1:A:651[B]:GLN:NE2	1.65	1.27
1:A:660[A]:VAL:O	1:A:664[A]:MET:HG2	1.43	1.17
1:A:651[B]:GLN:O	1:A:651[B]:GLN:CD	1.88	1.11
1:A:658[A]:PRO:HB3	10:A:1344:HOH:O	1.62	1.00
1:A:412:PRO:HA	1:A:589[B]:ASN:HD21	1.34	0.87
1:A:215[B]:LYS:NZ	10:A:902:HOH:O	2.13	0.80
1:A:597[B]:CYS:SG	1:A:671:LEU:HD22	2.21	0.80
1:A:597[B]:CYS:SG	1:A:671:LEU:CD2	2.76	0.73
1:A:437[A]:GLU:OE1	10:A:901:HOH:O	2.07	0.73
1:A:641:GLU:HG3	10:A:1265:HOH:O	1.90	0.71
1:A:400:ARG:O	1:A:404[B]:THR:HG23	1.91	0.71
1:A:362:LEU:CD1	1:A:406:LYS:HD2	2.23	0.69
1:A:131[A]:ILE:HG22	1:A:137:GLU:HG2	1.75	0.69
1:A:276[B]:GLU:HG2	1:A:277:TYR:N	2.10	0.67
1:A:370:ARG:HD2	1:A:669[B]:MET:HE1	1.77	0.66
1:A:447:VAL:HG22	1:A:669[B]:MET:CE	2.26	0.65
1:A:719[A]:VAL:O	1:A:719[A]:VAL:HG12	1.98	0.64
1:A:90:PHE:CE2	1:A:94:LYS:HE2	2.37	0.59
1:A:480[B]:GLU:O	1:A:480[B]:GLU:HG3	2.03	0.58
1:A:719[A]:VAL:CG1	1:A:719[A]:VAL:O	2.52	0.58
1:A:516:GLY:O	1:A:526[B]:GLN:NE2	2.34	0.58
1:A:719[B]:VAL:HG22	10:A:1347:HOH:O	2.05	0.57
1:A:412:PRO:HA	1:A:589[B]:ASN:ND2	2.15	0.56
1:A:447:VAL:HG22	1:A:669[B]:MET:HE2	1.88	0.55
1:A:177:VAL:HG12	1:A:188:LEU:HD11	1.88	0.55
1:A:362:LEU:HD11	1:A:406:LYS:HD2	1.87	0.55
1:A:480[B]:GLU:O	1:A:480[B]:GLU:CG	2.54	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:GLU:OE2	10:A:903:HOH:O	2.19	0.52
1:A:447:VAL:HG22	1:A:669[B]:MET:HE1	1.90	0.52
1:A:579:VAL:O	1:A:583[B]:MET:HG2	2.09	0.52
1:A:545:LYS:HE3	1:A:546:PHE:CZ	2.48	0.49
1:A:169:MET:HA	1:A:344:MET:O	2.14	0.48
1:A:58[B]:MET:SD	1:A:586:GLU:HG2	2.54	0.48
1:A:215[B]:LYS:HE2	10:A:958:HOH:O	2.14	0.48
1:A:489:GLU:CD	1:A:489:GLU:H	2.18	0.47
1:A:106:ASP:OD1	1:A:406:LYS:HE3	2.14	0.47
1:A:620:GLN:NE2	10:A:920:HOH:O	2.47	0.47
1:A:506:PHE:CB	1:A:509:MET:HG3	2.45	0.47
1:A:188:LEU:HD21	1:A:329[A]:VAL:HG11	1.97	0.46
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.50	0.46
1:A:651[B]:GLN:CG	1:A:651[B]:GLN:O	2.58	0.45
1:A:278:ALA:HB3	1:A:280[A]:ARG:CZ	2.47	0.44
1:A:517:SER:OG	1:A:522:GLU:OE2	2.30	0.44
1:A:517:SER:HB2	1:A:694:PRO:HG3	1.98	0.44
1:A:645[A]:LYS:HD2	10:A:953:HOH:O	2.17	0.44
1:A:454[B]:SER:OG	1:A:457[B]:GLU:OE2	2.35	0.44
1:A:719[B]:VAL:HG13	10:A:1343:HOH:O	2.16	0.44
1:A:613[B]:SER:OG	10:A:904:HOH:O	2.20	0.44
1:A:443[A]:GLN:HG3	1:A:444:GLU:CD	2.38	0.44
1:A:648:GLU:O	1:A:652:ASP:HB2	2.17	0.44
2:A:802:NAG:H81	10:A:1404:HOH:O	2.17	0.43
1:A:276[B]:GLU:HB3	10:A:908:HOH:O	2.18	0.43
1:A:456[B]:ILE:O	1:A:456[B]:ILE:HG13	2.19	0.43
1:A:473:LEU:HG	1:A:583[B]:MET:SD	2.59	0.42
1:A:188:LEU:CD2	1:A:329[A]:VAL:HG11	2.49	0.42
1:A:656[B]:SER:O	1:A:658[B]:PRO:HD3	2.19	0.42
1:A:457[A]:GLU:HG2	1:A:538:THR:HA	2.02	0.41
1:A:506:PHE:HB3	1:A:509:MET:HG3	2.02	0.41
1:A:669[B]:MET:HE2	1:A:669[B]:MET:HB2	1.55	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:A:910:HOH:O	10:A:1150:HOH:O[2_565]	1.83	0.37
1:A:656[B]:SER:O	10:A:1344:HOH:O[4_566]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	760/707 (108%)	738 (97%)	19 (2%)	3 (0%)	38 18

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	541	TRP
1	A	382	VAL
1	A	652	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	662/603 (110%)	652 (98%)	10 (2%)	70 52

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	LYS
1	A	507	SER
1	A	519	ASN
1	A	536	ARG
1	A	537	TYR
1	A	539[A]	LYS
1	A	540	ASN
1	A	545	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	600	TYR
1	A	673	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	540	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 4 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	801	1,2	14,14,15	0.53	0	15,19,21	1.46	5 (33%)
2	NAG	A	802	2	14,14,15	0.77	0	15,19,21	1.44	2 (13%)
2	NAG	A	803	1,3,2	14,14,15	0.73	1 (7%)	15,19,21	1.36	3 (20%)
2	NAG	A	804	2	14,14,15	0.57	0	15,19,21	1.19	1 (6%)
3	FUC	A	805	2	9,10,11	0.75	0	13,14,16	1.04	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	806	1,3,2	14,14,15	0.61	0	15,19,21	0.86	0
2	NAG	A	807	2	14,14,15	0.51	0	15,19,21	0.96	1 (6%)
3	FUC	A	808	2	9,10,11	0.86	0	13,14,16	1.26	2 (15%)
2	NAG	A	809	1	14,14,15	0.51	0	15,19,21	2.40	2 (13%)
2	NAG	A	810	1	14,14,15	0.81	0	15,19,21	1.63	4 (26%)
2	NAG	A	811	1,2	14,14,15	0.92	1 (7%)	15,19,21	1.00	1 (6%)
2	NAG	A	812	2,4	14,14,15	0.52	0	15,19,21	1.27	2 (13%)
4	BMA	A	813	2	11,11,12	0.88	1 (9%)	13,15,17	1.17	1 (7%)
2	NAG	A	814	1,2	14,14,15	0.64	0	15,19,21	1.49	3 (20%)
2	NAG	A	815	2,4	14,14,15	0.48	0	15,19,21	1.42	3 (20%)
4	BMA	A	816	2,5	11,11,12	0.58	0	13,15,17	1.00	0
5	MAN	A	817	4	11,11,12	0.69	0	13,15,17	1.23	1 (7%)
9	2G4	A	822	6	17,24,24	2.70	4 (23%)	20,31,31	1.43	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	801	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	802	2	-	0/6/23/26	0/1/1/1
2	NAG	A	803	1,3,2	-	0/6/23/26	0/1/1/1
2	NAG	A	804	2	-	0/6/23/26	0/1/1/1
3	FUC	A	805	2	-	0/0/17/20	0/1/1/1
2	NAG	A	806	1,3,2	-	0/6/23/26	0/1/1/1
2	NAG	A	807	2	-	0/6/23/26	0/1/1/1
3	FUC	A	808	2	-	0/0/17/20	0/1/1/1
2	NAG	A	809	1	-	0/6/23/26	0/1/1/1
2	NAG	A	810	1	-	0/6/23/26	0/1/1/1
2	NAG	A	811	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	812	2,4	-	0/6/23/26	0/1/1/1
4	BMA	A	813	2	-	0/2/19/22	0/1/1/1
2	NAG	A	814	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	815	2,4	-	0/6/23/26	0/1/1/1
4	BMA	A	816	2,5	-	0/2/19/22	0/1/1/1
5	MAN	A	817	4	-	0/2/19/22	0/1/1/1
9	2G4	A	822	6	-	0/16/23/23	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	822	2G4	PAX-OAF	-2.46	1.50	1.56
2	A	803	NAG	O5-C1	-2.01	1.40	1.43
4	A	813	BMA	C2-C3	2.01	1.55	1.52
2	A	811	NAG	O7-C7	2.84	1.29	1.23
9	A	822	2G4	CB-CA	3.65	1.58	1.53
9	A	822	2G4	PAX-N	4.38	1.66	1.61
9	A	822	2G4	PAX-OAC	8.77	1.56	1.46

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	822	2G4	CB-CA-C	-3.72	106.88	112.28
2	A	814	NAG	O5-C1-C2	-3.40	106.75	111.47
2	A	810	NAG	C3-C4-C5	-3.15	104.67	110.22
2	A	810	NAG	C1-O5-C5	-2.85	108.23	112.17
2	A	809	NAG	C6-C5-C4	-2.75	106.56	113.00
2	A	814	NAG	O4-C4-C5	-2.62	102.68	109.28
4	A	813	BMA	O5-C1-C2	-2.55	106.80	110.79
2	A	803	NAG	C1-C2-N2	-2.41	106.37	110.49
2	A	801	NAG	C2-N2-C7	-2.41	119.43	122.94
2	A	803	NAG	C1-O5-C5	-2.34	108.94	112.17
2	A	810	NAG	C1-C2-N2	-2.33	106.51	110.49
2	A	801	NAG	O5-C1-C2	-2.28	108.31	111.47
2	A	812	NAG	O7-C7-C8	-2.27	117.93	122.06
2	A	814	NAG	C2-N2-C7	-2.25	119.66	122.94
9	A	822	2G4	CG-CB-CA	-2.16	108.88	113.19
3	A	808	FUC	O5-C1-C2	-2.14	107.44	110.79
2	A	803	NAG	O3-C3-C2	-2.13	104.83	109.39
2	A	810	NAG	O5-C1-C2	-2.07	108.59	111.47
2	A	801	NAG	O4-C4-C3	-2.07	105.85	110.36
2	A	801	NAG	O3-C3-C4	-2.05	105.89	110.36
2	A	815	NAG	O4-C4-C5	-2.02	104.19	109.28
2	A	801	NAG	C4-C3-C2	2.14	114.16	111.02
2	A	804	NAG	C1-O5-C5	2.16	115.15	112.17
9	A	822	2G4	CAJ-CAV-CAK	2.18	121.61	118.16
2	A	815	NAG	C8-C7-N2	2.25	120.17	116.11
5	A	817	MAN	O3-C3-C2	2.31	114.22	110.02
3	A	808	FUC	O2-C2-C3	2.40	114.89	110.17
2	A	807	NAG	O4-C4-C5	2.41	115.35	109.28
2	A	802	NAG	C4-C3-C2	2.44	114.60	111.02
2	A	811	NAG	C2-N2-C7	2.63	126.78	122.94
2	A	802	NAG	C2-N2-C7	2.85	127.10	122.94
2	A	815	NAG	C1-O5-C5	3.07	116.40	112.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	812	NAG	C8-C7-N2	3.17	121.84	116.11
2	A	809	NAG	C1-O5-C5	7.99	123.18	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	802	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	694/707 (98%)	-0.02	42 (6%) 22 23	16, 28, 48, 70	1 (0%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	541	TRP	6.7
1	A	542	GLU	5.1
1	A	719[A]	VAL	4.5
1	A	543	THR	4.0
1	A	55	LYS	4.0
1	A	505	GLU	3.9
1	A	134	ASP	3.5
1	A	226	ILE	3.4
1	A	153	ASN	3.4
1	A	155[A]	SER	3.4
1	A	201[A]	VAL	3.4
1	A	506	PHE	3.3
1	A	656[A]	SER	3.3
1	A	133	GLU	3.3
1	A	335	GLY	3.2
1	A	544	ASN	3.2
1	A	152	GLU	3.2
1	A	227	LEU	3.1
1	A	225	VAL	3.1
1	A	154	VAL	3.0
1	A	123	THR	3.0
1	A	135	GLY	2.9
1	A	507	SER	2.9
1	A	174[A]	LEU	2.9
1	A	336	ASN	2.8
1	A	283[A]	ILE	2.7
1	A	652	ASP	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	132	ASN	2.7
1	A	122	LYS	2.6
1	A	487	GLY	2.6
1	A	297	ILE	2.5
1	A	456[A]	ILE	2.4
1	A	504	PRO	2.4
1	A	237	PRO	2.3
1	A	136	ASN	2.2
1	A	124	HIS	2.2
1	A	337	PHE	2.2
1	A	284	ALA	2.2
1	A	253	VAL	2.1
1	A	294	VAL	2.1
1	A	408	GLU	2.0
1	A	203	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	A	802	14/15	0.76	0.23	5.29	44,51,59,60	0
2	NAG	A	803	14/15	0.79	0.26	1.92	48,55,67,73	0
7	CA	A	820	1/1	1.00	0.07	1.89	18,18,18,18	0
2	NAG	A	810	14/15	0.92	0.16	1.12	33,44,51,54	0
2	NAG	A	811	14/15	0.92	0.09	1.04	29,34,41,41	0
5	MAN	A	817	11/12	0.95	0.15	0.62	47,50,54,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAG	A	814	14/15	0.94	0.08	0.05	23,29,38,44	0
9	2G4	A	822	24/24	0.98	0.08	-0.50	20,24,43,45	0
6	ZN	A	819	1/1	1.00	0.05	-1.79	21,21,21,21	0
6	ZN	A	818	1/1	1.00	0.05	-3.02	20,20,20,20	0
8	CL	A	821	1/1	0.99	0.03	-3.07	28,28,28,28	0
4	BMA	A	813	11/12	0.77	0.28	-	61,66,70,71	0
2	NAG	A	812	14/15	0.89	0.18	-	37,45,53,53	0
2	NAG	A	807	14/15	0.80	0.30	-	59,65,70,71	0
2	NAG	A	804	14/15	0.81	0.41	-	72,76,78,79	0
2	NAG	A	809	14/15	0.82	0.20	-	69,75,77,77	0
3	FUC	A	808	10/11	0.87	0.39	-	56,62,64,66	0
2	NAG	A	815	14/15	0.89	0.23	-	34,40,50,53	0
2	NAG	A	806	14/15	0.92	0.15	-	45,49,56,59	0
4	BMA	A	816	11/12	0.93	0.15	-	39,43,46,47	0
2	NAG	A	801	14/15	0.95	0.10	-	35,43,49,52	0
3	FUC	A	805	10/11	0.91	0.40	-	64,68,69,69	0

6.5 Other polymers [i](#)

There are no such residues in this entry.