



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2018 – 09:19 AM EST

PDB ID : 5P9H
Title : BTK1 COCRYSTALLIZED WITH RN983
Authors : Gardberg, A.S.
Deposited on : 2016-09-20
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	<div> <div>20%</div> <div>86%</div> <div>8%</div> <div>5%</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2043 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase BTK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	1	0
			1973	1265	325	363	20			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	381	GLY	-	expression tag	UNP Q06187

- Molecule 2 is 6- {tert}-butyl-8-fluoranyl-2-[3-(hydroxymethyl)-4-[1-methyl-5-[[5-(1-methylpiperidin-4-yl)pyridin-2-yl]amino]-6-oxidanylidene-pyridazin-3-yl]pyridin-2-yl]phthalazin-1-one (three-letter code: 7G7) (formula: C₃₄H₃₇FN₈O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			46	34	1	8	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	24	Total	O	0	0
			24	24		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase BTK



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	62.76 Å 101.40 Å 38.10 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.36 – 1.95 24.22 – 1.95	Depositor EDS
% Data completeness (in resolution range)	85.2 (24.36-1.95) 85.2 (24.22-1.95)	Depositor EDS
R_{merge}	0.23	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.96 (at 1.95 Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.267 , 0.313 0.290 , 0.332	Depositor DCC
R_{free} test set	773 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.283	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 54.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	2043	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 7G7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2020	0.68	0/2749

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1973	0	1764	10	0
2	A	46	0	0	0	0
3	A	24	0	0	0	0
All	All	2043	0	1764	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:592:SER:HB3	1:A:595:LYS:HB2	1.69	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:399:PRO:HB3	1:A:467:GLN:HE22	1.56	0.69
1:A:501:MET:HG2	1:A:535:VAL:HG21	1.82	0.62
1:A:588:TRP:O	1:A:592:SER:HB2	2.06	0.56
1:A:577:LYS:HA	1:A:580:ILE:HB	1.90	0.53
1:A:456:LYS:HA	1:A:536:LYS:HG2	1.92	0.50
1:A:398:ASP:HB3	1:A:401:ASP:HB2	1.97	0.47
1:A:565:PRO:HG3	1:A:578:SER:HA	1.98	0.45
1:A:498:LEU:HB3	1:A:590:ILE:HG22	1.99	0.44
1:A:432:ILE:HD11	1:A:472:ILE:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/279 (94%)	256 (97%)	7 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/248 (75%)	182 (97%)	5 (3%)	50	39

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	410	THR
1	A	572	SER
1	A	586	LEU
1	A	592	SER
1	A	649	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	459	GLN
1	A	467	GLN
1	A	484	ASN
1	A	650	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	7G7	A	701	-	46,51,51	1.17	3 (6%)	55,76,76	2.06	14 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7G7	A	701	-	-	0/20/34/34	0/6/6/6

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	7G7	C28-C29	2.45	1.44	1.41
2	A	701	7G7	C14-C13	2.93	1.47	1.41
2	A	701	7G7	C11-C12	4.49	1.48	1.41

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	7G7	C17-C16-C12	-5.26	119.75	124.57
2	A	701	7G7	C13-C14-N15	-4.44	119.19	124.70
2	A	701	7G7	C2-C3-N4	-3.42	119.97	123.92
2	A	701	7G7	C37-C38-N39	-3.17	120.32	124.68
2	A	701	7G7	C35-C36-C37	-2.30	118.86	121.20
2	A	701	7G7	C6-C5-N10	2.10	120.34	118.01
2	A	701	7G7	C36-C37-C38	2.20	119.06	116.83
2	A	701	7G7	C18-C17-C16	2.69	121.50	118.48
2	A	701	7G7	C14-C13-C12	2.98	120.11	117.35
2	A	701	7G7	F21-C16-C12	3.24	122.21	118.51
2	A	701	7G7	C42-N43-C44	3.96	114.75	109.47
2	A	701	7G7	C1-C9-N26	4.45	120.39	115.85
2	A	701	7G7	C38-N39-C34	4.78	122.88	117.91
2	A	701	7G7	C3-N4-C5	5.85	122.13	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/279 (94%)	1.35	55 (20%) 1 1	33, 58, 100, 121	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	421	TRP	5.7
1	A	656	ASP	5.5
1	A	602	THR	5.4
1	A	415	VAL	4.0
1	A	607	ALA	3.9
1	A	402	LEU	3.8
1	A	450	MET	3.7
1	A	611	ALA	3.7
1	A	583	PHE	3.7
1	A	446	ALA	3.6
1	A	397	ILE	3.5
1	A	442	PHE	3.5
1	A	646	ILE	3.4
1	A	452	LEU	3.4
1	A	461	TYR	3.3
1	A	613	GLY	3.2
1	A	557	SER	3.2
1	A	654	VAL	3.2
1	A	429	ILE	3.1
1	A	601	PHE	3.0
1	A	574	PHE	2.9
1	A	547	LEU	2.9
1	A	464	CYS	2.8
1	A	476	TYR	2.8
1	A	471	PHE	2.7
1	A	586	LEU	2.6
1	A	424	GLN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	448	VAL	2.6
1	A	502	CYS	2.5
1	A	652	LEU	2.5
1	A	423	GLY	2.5
1	A	603	ASN	2.5
1	A	413	PHE	2.5
1	A	432	ILE	2.4
1	A	425	TYR	2.4
1	A	594	GLY	2.4
1	A	580	ILE	2.3
1	A	511	TYR	2.3
1	A	631	TYR	2.3
1	A	534	VAL	2.3
1	A	658	GLU	2.3
1	A	403	THR	2.3
1	A	434	GLU	2.3
1	A	416	VAL	2.3
1	A	522	LEU	2.2
1	A	422	ARG	2.2
1	A	435	GLY	2.1
1	A	523	ALA	2.1
1	A	634	TRP	2.1
1	A	505	VAL	2.1
1	A	524	ALA	2.1
1	A	585	VAL	2.1
1	A	409	GLY	2.1
1	A	548	ASP	2.0
1	A	453	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	7G7	A	701	46/46	0.77	0.22	1.56	40,50,72,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.