



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 05:41 pm GMT

PDB ID : 1PEV
Title : Crystal Structure of the Actin Interacting Protein from *Caenorhabditis Elegans*
Authors : Vorobiev, S.; Mohri, K.; Fedorov, A.A.; Ono, S.; Almo, S.C.; Burley, S.K.;
New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2003-05-22
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	611	<div> <div>2%</div> <div> <div></div> <div>77%</div> <div>21%</div> <div>.</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Actin interacting protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	610	Total	C	N	O	S	0	0	0
			4600	2901	788	902	9			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	330	Total	O	0	0
			330	330		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Actin interacting protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.66Å 90.82Å 76.98Å 90.00° 94.06° 90.00°	Depositor
Resolution (Å)	21.72 – 2.00 21.72 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.2 (21.72-2.00) 97.3 (21.72-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.01Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.197 , 0.234 0.197 , 0.234	Depositor DCC
R_{free} test set	1471 reflections (4.01%)	DCC
Wilson B-factor (Å ²)	18.8	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4930	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	1/4709 (0.0%)	0.73	4/6418 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	557	ASP	CB-CG	6.04	1.64	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	398	SER	N-CA-C	-6.79	92.66	111.00
1	A	10	PHE	N-CA-C	-5.99	94.83	111.00
1	A	330	LEU	CA-CB-CG	5.44	127.82	115.30
1	A	395	ALA	N-CA-C	5.13	124.85	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4600	0	4487	105	0
2	A	330	0	0	14	0
All	All	4930	0	4487	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 12.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:ASN:HD21	1:A:220:VAL:HG12	1.26	0.96
1:A:275:LYS:HE3	1:A:312:GLU:O	1.75	0.86
1:A:396:GLY:HA2	1:A:399:GLY:HA2	1.58	0.85
1:A:514:LYS:HE3	2:A:1015:HOH:O	1.80	0.81
1:A:142:ASN:ND2	1:A:144:THR:H	1.83	0.77
1:A:379:GLY:O	1:A:395:ALA:HB2	1.85	0.76
1:A:218:ASN:ND2	1:A:220:VAL:HG12	2.00	0.75
1:A:218:ASN:HD21	1:A:220:VAL:CG1	1.99	0.74
1:A:357:ILE:HD11	2:A:1089:HOH:O	1.86	0.74
1:A:361:VAL:HG22	1:A:400:VAL:HG21	1.70	0.73
1:A:104:GLY:HA3	1:A:123:GLU:O	1.88	0.72
1:A:357:ILE:HD12	2:A:956:HOH:O	1.89	0.72
1:A:26:ASN:HD22	1:A:291:ILE:HD11	1.56	0.71
1:A:393:VAL:HG22	1:A:405:ALA:HB2	1.76	0.66
1:A:395:ALA:O	1:A:397:GLY:N	2.29	0.65
1:A:522:ASN:O	1:A:525:GLU:HG2	1.97	0.64
1:A:310:ASN:HD22	1:A:310:ASN:C	2.01	0.63
1:A:142:ASN:C	1:A:142:ASN:HD22	2.02	0.63
1:A:126:GLU:CD	1:A:126:GLU:H	2.01	0.63
1:A:398:SER:HB3	1:A:401:ASP:OD1	1.99	0.62
1:A:123:GLU:HB3	2:A:1124:HOH:O	1.99	0.60
1:A:142:ASN:HD22	1:A:143:LEU:N	1.98	0.60
1:A:194:HIS:HE1	2:A:845:HOH:O	1.85	0.60
1:A:283:ILE:HD11	2:A:1128:HOH:O	2.02	0.60
1:A:429:ALA:HB2	1:A:454:VAL:HB	1.85	0.58
1:A:160:PRO:HB2	1:A:162:ARG:HH12	1.67	0.58
1:A:541:CYS:HB2	2:A:1078:HOH:O	2.02	0.58
1:A:392:VAL:HB	1:A:406:VAL:HG12	1.85	0.58
1:A:283:ILE:HG12	2:A:1123:HOH:O	2.02	0.57
1:A:537:ALA:HB3	1:A:557:ASP:HB3	1.87	0.57
1:A:53:GLU:HG3	2:A:959:HOH:O	2.04	0.56
1:A:514:LYS:HG2	1:A:533:THR:CG2	2.36	0.56
1:A:224:LYS:HE3	1:A:227:VAL:HG13	1.86	0.56
1:A:107:LYS:HG3	1:A:122:GLY:HA2	1.87	0.55
1:A:399:GLY:O	1:A:400:VAL:HG13	2.07	0.55
1:A:512:SER:O	1:A:513:ARG:HB2	2.06	0.55
1:A:558:ASN:N	1:A:558:ASN:OD1	2.40	0.55
1:A:275:LYS:HD3	1:A:314:GLY:HA3	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:431:TYR:O	1:A:450:ASN:HA	2.08	0.53
1:A:102:PHE:HB2	1:A:124:GLY:HA3	1.92	0.52
1:A:12:SER:HB2	1:A:602:SER:HB2	1.90	0.52
1:A:183:LYS:C	1:A:183:LYS:HD3	2.30	0.52
1:A:462:PHE:CE1	1:A:476:LYS:HE3	2.45	0.51
1:A:77:GLY:HA2	1:A:82:ASN:O	2.11	0.51
1:A:347:HIS:HD2	1:A:364:ASP:OD1	1.93	0.51
1:A:218:ASN:HD22	1:A:221:ASP:H	1.59	0.50
1:A:181:LYS:HG2	2:A:1022:HOH:O	2.11	0.49
1:A:406:VAL:HG13	1:A:406:VAL:O	2.12	0.49
1:A:142:ASN:C	1:A:142:ASN:ND2	2.64	0.49
1:A:393:VAL:HG22	1:A:405:ALA:CB	2.43	0.49
1:A:392:VAL:HB	1:A:406:VAL:CG1	2.43	0.47
1:A:373:ILE:HD11	1:A:381:LEU:HD11	1.95	0.47
1:A:310:ASN:HD21	1:A:312:GLU:HB2	1.79	0.47
1:A:36:CYS:HB3	1:A:62:THR:OG1	2.15	0.47
1:A:328:THR:HG21	1:A:369:MET:SD	2.55	0.47
1:A:488:ILE:HG21	1:A:518:TYR:CZ	2.50	0.46
1:A:562:VAL:HB	1:A:573:ILE:HB	1.96	0.46
1:A:546:PRO:HG3	2:A:1013:HOH:O	2.13	0.46
1:A:310:ASN:C	1:A:310:ASN:ND2	2.69	0.46
1:A:78:ASP:OD2	1:A:82:ASN:HB2	2.15	0.46
1:A:330:LEU:HD11	1:A:598:ALA:HB2	1.98	0.45
1:A:558:ASN:HA	1:A:582:SER:O	2.16	0.45
1:A:10:PHE:HB2	1:A:604:ILE:HB	1.98	0.45
1:A:176:GLU:OE2	1:A:183:LYS:HG2	2.17	0.45
1:A:94:ILE:HG12	2:A:1057:HOH:O	2.16	0.45
1:A:81:GLY:O	1:A:101:VAL:HG22	2.17	0.45
1:A:288:LEU:CD1	1:A:302:ALA:HA	2.46	0.45
1:A:514:LYS:HG2	1:A:533:THR:HG22	1.99	0.45
1:A:175:PHE:CD2	1:A:182:PHE:HA	2.52	0.45
1:A:425:ILE:HD12	1:A:460:LYS:HD2	1.99	0.44
1:A:34:GLN:HA	1:A:42:TYR:O	2.16	0.44
1:A:478:SER:O	1:A:481:SER:HB3	2.16	0.44
1:A:178:PRO:HA	1:A:179:PRO:C	2.37	0.44
1:A:237:ALA:HB1	1:A:266:TRP:CZ2	2.52	0.44
1:A:174:ILE:HG23	1:A:183:LYS:HB3	1.98	0.44
1:A:277:ILE:HD13	1:A:309:VAL:HG13	2.00	0.44
1:A:11:PRO:HA	1:A:603:ASN:HD22	1.83	0.43
1:A:343:ASP:OD2	1:A:347:HIS:HB2	2.18	0.43
1:A:191:LYS:HD3	1:A:211:ASP:HA	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:351:TRP:CE3	1:A:358:SER:HB3	2.53	0.43
1:A:277:ILE:HD13	1:A:309:VAL:CG1	2.49	0.43
1:A:325:LYS:HG3	1:A:345:GLU:HG3	2.00	0.43
1:A:169:ASP:O	1:A:170:ASN:HB2	2.19	0.42
1:A:218:ASN:ND2	1:A:220:VAL:CG1	2.74	0.42
1:A:362:PHE:CD1	1:A:363:PRO:HA	2.55	0.42
1:A:107:LYS:HE2	1:A:107:LYS:HB3	1.91	0.42
1:A:32:LYS:HG2	1:A:45:PRO:HA	2.00	0.42
1:A:472:VAL:HB	1:A:488:ILE:HB	2.02	0.42
1:A:69:PRO:HD2	1:A:115:SER:OG	2.20	0.42
1:A:142:ASN:ND2	1:A:144:THR:N	2.61	0.42
1:A:454:VAL:HG22	1:A:455:ALA:N	2.35	0.42
1:A:556:LEU:HA	1:A:583:SER:CB	2.50	0.42
1:A:13:LEU:HB3	1:A:14:PRO:HD2	2.02	0.41
1:A:181:LYS:HE2	2:A:1022:HOH:O	2.19	0.41
1:A:333:SER:HA	1:A:375:THR:HG21	2.01	0.41
1:A:390:LEU:C	1:A:390:LEU:HD23	2.39	0.41
1:A:282:ARG:HH11	1:A:282:ARG:HG2	1.85	0.41
1:A:395:ALA:HB1	1:A:396:GLY:H	1.48	0.41
1:A:130:HIS:CD2	2:A:1034:HOH:O	2.73	0.41
1:A:183:LYS:CD	1:A:183:LYS:C	2.89	0.41
1:A:257:ALA:HB1	1:A:287:GLN:HB3	2.02	0.41
1:A:404:LYS:HE3	1:A:404:LYS:HB2	1.81	0.41
1:A:323:HIS:CE1	1:A:349:ASN:HD22	2.39	0.40
1:A:105:PRO:HD2	1:A:123:GLU:HB2	2.03	0.40
1:A:10:PHE:HE1	1:A:50:THR:HG22	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	608/611 (100%)	585 (96%)	21 (4%)	2 (0%)	44 40

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	396	GLY
1	A	400	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	509/511 (100%)	498 (98%)	11 (2%)	57 60

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	102	PHE
1	A	126	GLU
1	A	134	PHE
1	A	142	ASN
1	A	183	LYS
1	A	310	ASN
1	A	317	ASP
1	A	385	SER
1	A	398	SER
1	A	400	VAL
1	A	558	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	A	34	GLN
1	A	130	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	142	ASN
1	A	146	GLN
1	A	194	HIS
1	A	218	ASN
1	A	310	ASN
1	A	318	GLN
1	A	347	HIS
1	A	413	GLN
1	A	473	HIS
1	A	603	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	610/611 (99%)	-0.10	14 (2%) 61 60	10, 18, 36, 60	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	396	GLY	10.2
1	A	399	GLY	5.9
1	A	397	GLY	5.6
1	A	398	SER	5.0
1	A	2	SER	3.6
1	A	18	ARG	3.2
1	A	395	ALA	3.2
1	A	400	VAL	3.1
1	A	72	TYR	2.9
1	A	401	ASP	2.8
1	A	531	SER	2.7
1	A	403	SER	2.7
1	A	90	GLN	2.3
1	A	91	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.