



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:50 am GMT

PDB ID : 2PFH
Title : Complex of Aldose Reductase with NADP⁺ and simultaneously bound competitive inhibitors Fidarestat and IDD594. Concentration of Fidarestat in soaking solution is less than concentration of IDD594.
Authors : Petrova, T.; Hazemann, I.; Cousido, A.; Mitschler, A.; Ginell, S.; Joachimiak, A.; Podjarny, A.
Deposited on : 2007-04-05
Resolution : 0.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 0.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1119 (1.08-0.64)
Clashscore	112137	1078 (1.06-0.66)
Ramachandran outliers	110173	1128 (1.08-0.64)
Sidechain outliers	110143	1000 (1.06-0.66)
RSRZ outliers	101464	1000 (1.06-0.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> % 82% 16% .. </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	CIT	A	450[A]	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3919 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aldose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	314	3098	1992	518	571	17	7	105	1


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4	ILE	LEU	SEE REMARK 999	UNP P15121

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	1

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is IDD594 (three-letter code: LDT) (formula: $C_{16}H_{12}BrF_2NO_3S$).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	Br	C	F	N	O	S	
			24	1	16	2	1	3	1	1

- Molecule 5 is (2S,4S)-2-AMINOFORMYL-6-FLUORO-SPIRO[CHROMAN-4,4'-IMIDAZO LIDINE]-2',5'-DIONE (three-letter code: FID) (formula: $C_{12}H_{10}FN_3O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	F	N	O	0	1
			20	12	1	3	4		

- Molecule 6 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	1
			13	6	7		
6	A	1	Total	C	O	0	1
			13	6	7		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	656	Total	O	0	305
			702	702		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aldose reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.15Å 66.55Å 47.24Å 90.00° 92.31° 90.00°	Depositor
Resolution (Å)	50.00 – 0.85 34.74 – 0.85	Depositor EDS
% Data completeness (in resolution range)	95.4 (50.00-0.85) 95.4 (34.74-0.85)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.02 (at 0.85Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.082 , 0.095 0.094 , 0.095	Depositor DCC
R_{free} test set	12783 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	5.3	Xtriage
Anisotropy	0.096	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 72.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.002 for -l,k,h 0.018 for -h,-k,l 0.015 for l,-k,h	Xtriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	3919	wwPDB-VP
Average B, all atoms (Å ²)	8.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, CIT, LDT, FID, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	6.18	17/3368 (0.5%)	2.58	60/4566 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	312[A]	HIS	CG-CD2	178.57	4.39	1.35
1	A	312[B]	HIS	CG-CD2	178.57	4.39	1.35
1	A	312[A]	HIS	CG-ND1	124.21	4.12	1.38
1	A	312[B]	HIS	CG-ND1	124.21	4.12	1.38
1	A	26[A]	GLN	CD-NE2	90.05	3.58	1.32
1	A	26[B]	GLN	CD-NE2	90.05	3.58	1.32
1	A	26[A]	GLN	CD-OE1	77.60	2.94	1.24
1	A	26[B]	GLN	CD-OE1	77.60	2.94	1.24
1	A	293[A]	ARG	CB-CG	-7.41	1.32	1.52
1	A	293[B]	ARG	CB-CG	-7.41	1.32	1.52
1	A	29[A]	GLU	CD-OE1	-6.96	1.18	1.25
1	A	29[B]	GLU	CD-OE1	-6.96	1.18	1.25
1	A	64[A]	GLU	CD-OE1	-5.65	1.19	1.25
1	A	64[B]	GLU	CD-OE1	-5.65	1.19	1.25
1	A	150	GLU	CD-OE2	-5.45	1.19	1.25
1	A	40[A]	ARG	CZ-NH2	5.21	1.39	1.33
1	A	40[B]	ARG	CZ-NH2	5.21	1.39	1.33

All (60) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	312[A]	HIS	ND1-CG-CD2	-54.82	29.25	106.00
1	A	312[B]	HIS	ND1-CG-CD2	-54.82	29.25	106.00
1	A	26[A]	GLN	CG-CD-OE1	-40.72	40.15	121.60
1	A	26[B]	GLN	CG-CD-OE1	-40.72	40.15	121.60
1	A	26[A]	GLN	OE1-CD-NE2	-36.71	37.48	121.90
1	A	26[B]	GLN	OE1-CD-NE2	-36.71	37.48	121.90
1	A	26[A]	GLN	CG-CD-NE2	-34.63	33.59	116.70
1	A	26[B]	GLN	CG-CD-NE2	-34.63	33.59	116.70
1	A	312[A]	HIS	CG-ND1-CE1	32.17	153.24	108.20
1	A	312[B]	HIS	CG-ND1-CE1	32.17	153.24	108.20
1	A	312[A]	HIS	CB-CG-ND1	-20.63	71.63	123.20
1	A	312[B]	HIS	CB-CG-ND1	-20.63	71.63	123.20
1	A	3	ARG	NE-CZ-NH2	-15.92	112.34	120.30
1	A	312[A]	HIS	CG-CD2-NE2	14.98	137.65	109.20
1	A	312[B]	HIS	CG-CD2-NE2	14.98	137.65	109.20
1	A	232	ARG	NE-CZ-NH1	11.01	125.80	120.30
1	A	293[A]	ARG	NE-CZ-NH1	-10.57	115.01	120.30
1	A	293[B]	ARG	NE-CZ-NH1	-10.57	115.01	120.30
1	A	276[A]	PHE	CB-CG-CD1	10.03	127.82	120.80
1	A	276[B]	PHE	CB-CG-CD1	10.03	127.82	120.80
1	A	312[A]	HIS	CB-CG-CD2	-9.74	100.59	130.80
1	A	312[B]	HIS	CB-CG-CD2	-9.74	100.59	130.80
1	A	217[A]	ARG	NE-CZ-NH2	-9.21	115.69	120.30
1	A	217[B]	ARG	NE-CZ-NH2	-9.21	115.69	120.30
1	A	217[A]	ARG	NE-CZ-NH1	8.29	124.44	120.30
1	A	217[B]	ARG	NE-CZ-NH1	8.29	124.44	120.30
1	A	268	ARG	NE-CZ-NH1	7.89	124.25	120.30
1	A	224	ASP	CB-CG-OD1	7.62	125.16	118.30
1	A	63[A]	ARG	NE-CZ-NH2	-7.59	116.51	120.30
1	A	216	ASP	CB-CG-OD2	-7.04	111.96	118.30
1	A	276[A]	PHE	CB-CG-CD2	-6.66	116.14	120.80
1	A	276[B]	PHE	CB-CG-CD2	-6.66	116.14	120.80
1	A	3	ARG	NH1-CZ-NH2	6.59	126.66	119.40
1	A	134[A]	ASP	CB-CG-OD2	-6.59	112.36	118.30
1	A	134[B]	ASP	CB-CG-OD2	-6.59	112.36	118.30
1	A	82	TYR	CB-CG-CD1	6.43	124.86	121.00
1	A	296[A]	ARG	NE-CZ-NH2	-5.96	117.32	120.30
1	A	296[B]	ARG	NE-CZ-NH2	-5.96	117.32	120.30
1	A	277[A]	ASP	CB-CG-OD2	-5.87	113.01	118.30
1	A	277[B]	ASP	CB-CG-OD2	-5.87	113.01	118.30
1	A	278[A]	PHE	CB-CG-CD1	5.82	124.87	120.80
1	A	278[B]	PHE	CB-CG-CD1	5.82	124.87	120.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	103	TYR	CB-CG-CD1	5.73	124.44	121.00
1	A	193[A]	GLU	CB-CA-C	-5.69	99.01	110.40
1	A	193[B]	GLU	CB-CA-C	-5.69	99.01	110.40
1	A	135[A]	THR	O-C-N	-5.62	113.72	122.70
1	A	135[B]	THR	O-C-N	-5.62	113.72	122.70
1	A	135[C]	THR	O-C-N	-5.62	113.72	122.70
1	A	291	TYR	CD1-CE1-CZ	5.59	124.83	119.80
1	A	73	PHE	CB-CG-CD1	5.48	124.64	120.80
1	A	67[A]	VAL	CB-CA-C	-5.41	101.12	111.40
1	A	67[B]	VAL	CB-CA-C	-5.41	101.12	111.40
1	A	232	ARG	NH1-CZ-NH2	-5.32	113.54	119.40
1	A	291	TYR	CB-CG-CD2	5.19	124.12	121.00
1	A	209	TYR	CA-CB-CG	5.16	123.21	113.40
1	A	70[A]	GLU	CB-CG-CD	5.06	127.85	114.20
1	A	70[B]	GLU	CB-CG-CD	5.06	127.85	114.20
1	A	115	PHE	CB-CG-CD2	-5.05	117.27	120.80
1	A	127[A]	SER	O-C-N	-5.02	114.67	123.20
1	A	127[B]	SER	O-C-N	-5.02	114.67	123.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	209	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3098	0	3002	28	0
2	A	1	0	0	0	0
3	A	48	0	26	4	0
4	A	24	0	11	1	0
5	A	20	0	10	3	0
6	A	26	0	10	0	0
7	A	702	0	0	13	0
All	All	3919	0	3059	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:298[B]:CYS:SG	5:A:321[B]:FID:H8I2	2.13	0.89
1:A:295[A]:TRP:CZ3	7:A:3675[A]:HOH:O	2.29	0.84
1:A:274:LYS:HE3	7:A:3509[A]:HOH:O	1.83	0.78
3:A:318:NDP:H42N	5:A:321[B]:FID:C2I	2.28	0.64
1:A:209:TYR:CE1	3:A:318:NDP:H41N	2.34	0.62
1:A:209:TYR:CZ	3:A:318:NDP:H41N	2.36	0.60
1:A:244[B]:THR:HG22	7:A:3120:HOH:O	2.01	0.59
1:A:181:VAL:HA	1:A:205:VAL:O	2.03	0.57
1:A:136[C]:ASN:O	1:A:139:ASP:HB2	2.04	0.56
1:A:225:PRO:HG2	1:A:295[B]:TRP:CD2	2.44	0.52
1:A:225:PRO:HG2	1:A:295[B]:TRP:CE2	2.44	0.52
1:A:262:LYS:O	3:A:318:NDP:H8A	2.11	0.50
1:A:70[A]:GLU:HG2	7:A:3350:HOH:O	2.12	0.50
1:A:41:HIS:HA	1:A:73:PHE:O	2.15	0.47
1:A:20:TRP:CE3	1:A:21:LYS:HE3	2.49	0.47
1:A:138:LEU:HG	7:A:3701[B]:HOH:O	2.15	0.47
1:A:309[B]:TYR:CE2	1:A:311[B]:PHE:HB2	2.51	0.46
1:A:301[B]:LEU:HD11	7:A:3260:HOH:O	2.16	0.45
1:A:193[A]:GLU:HG3	7:A:3503[A]:HOH:O	2.17	0.45
1:A:226:SER:N	7:A:3634[B]:HOH:O	2.45	0.45
1:A:154[B]:LYS:HD3	7:A:3334:HOH:O	2.18	0.44
1:A:116[A]:LYS:HZ3	1:A:123:PRO:HB2	1.83	0.43
1:A:101:LEU:HD23	7:A:3533[A]:HOH:O	2.19	0.43
1:A:116[A]:LYS:NZ	7:A:3267:HOH:O	2.48	0.42
1:A:136[A]:ASN:HB2	7:A:3076:HOH:O	2.20	0.42
1:A:113:THR:OG1	4:A:320[A]:LDT:BR8	2.85	0.42
1:A:277[B]:ASP:HB3	7:A:3250:HOH:O	2.20	0.40
1:A:48:TYR:OH	5:A:321[B]:FID:O3I	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	408/316 (129%)	404 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/281 (131%)	368 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	129	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NDP	A	318	-	43,52,52	1.56	7 (16%)	49,80,80	1.16	5 (10%)
4	LDT	A	320[A]	-	22,25,25	1.39	1 (4%)	30,34,34	1.52	6 (20%)
5	FID	A	321[B]	-	21,22,22	0.83	1 (4%)	21,34,34	3.03	10 (47%)
6	CIT	A	400[A]	-	3,12,12	0.02	0	3,17,17	1.24	0
6	CIT	A	450[A]	-	3,12,12	1.45	1 (33%)	3,17,17	1.42	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NDP	A	318	-	-	0/30/77/77	0/5/5/5
4	LDT	A	320[A]	-	-	0/12/14/14	0/2/2/2
5	FID	A	321[B]	-	-	0/4/34/34	0/3/3/3
6	CIT	A	400[A]	-	-	0/6/16/16	0/0/0/0
6	CIT	A	450[A]	-	-	0/6/16/16	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	318	NDP	C4N-C5N	-4.55	1.39	1.49
3	A	318	NDP	C4A-N3A	-2.30	1.32	1.35
5	A	321[B]	FID	O3I-C2I	2.08	1.27	1.23
6	A	450[A]	CIT	O7-C3	2.22	1.46	1.43
3	A	318	NDP	C3B-C2B	2.22	1.58	1.53
3	A	318	NDP	C2N-C3N	2.51	1.42	1.34
3	A	318	NDP	C1D-N1N	2.75	1.54	1.46
3	A	318	NDP	O4B-C1B	3.02	1.45	1.41
4	A	320[A]	LDT	C6-C5	3.16	1.43	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	318	NDP	C6N-C5N	4.33	1.41	1.33

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	321[B]	FID	C7I-C5-N4	-7.19	103.38	107.89
5	A	321[B]	FID	O6I-C5-C7I	-3.78	122.59	125.81
5	A	321[B]	FID	N4-C2I-N1I	-3.48	103.55	107.58
4	A	320[A]	LDT	F9-C5-C2	-2.98	113.90	118.24
5	A	321[B]	FID	O10-C11-C16	-2.67	111.91	116.61
5	A	321[B]	FID	O10-C9-C8I	-2.59	106.09	111.49
3	A	318	NDP	O2D-C2D-C3D	-2.44	104.01	111.83
4	A	320[A]	LDT	C13-N17-C11	-2.35	121.59	124.86
6	A	450[A]	CIT	C3-C2-C1	-2.04	111.76	114.95
4	A	320[A]	LDT	O15-C4-C2	-2.02	118.68	123.61
3	A	318	NDP	C5A-C6A-N1A	-2.02	113.58	119.70
4	A	320[A]	LDT	C3-C7-C4	2.10	120.95	118.21
3	A	318	NDP	C1B-N9A-C4A	2.13	130.31	126.64
4	A	320[A]	LDT	C26-C29-C25	2.24	122.22	119.19
4	A	320[A]	LDT	C28-C27-C24	2.40	127.09	124.09
5	A	321[B]	FID	C13-C12-C11	2.70	120.70	116.76
3	A	318	NDP	C5A-C6A-N6A	2.80	126.19	120.47
5	A	321[B]	FID	C5-N4-C2I	3.66	115.40	111.70
3	A	318	NDP	N3A-C2A-N1A	3.67	132.06	128.86
5	A	321[B]	FID	C8I-C7I-N1I	4.53	119.98	112.08
5	A	321[B]	FID	O10-C11-C12	4.75	128.91	122.31
5	A	321[B]	FID	O6I-C5-N4	5.42	133.79	126.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	318	NDP	4	0
4	A	320[A]	LDT	1	0
5	A	321[B]	FID	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/316 (99%)	-0.74	2 (0%) 89 79	3, 6, 13, 27	26 (8%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	0[A]	MET	6.0
1	A	1	ALA	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	CIT	A	450[A]	13/13	0.88	0.23	45.74	1,6,7,8	13
6	CIT	A	400[A]	13/13	0.98	0.10	1.25	7,10,12,16	13
5	FID	A	321[B]	20/20	0.99	0.05	0.33	3,4,7,9	20

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NDP	A	318	48/48	1.00	0.03	-1.00	2,3,5,6	0
4	LDT	A	320[A]	24/24	1.00	0.03	-1.45	3,4,5,5	24
2	CL	A	2000[A]	1/1	1.00	0.03	-2.80	5,5,5,5	1

6.5 Other polymers [i](#)

There are no such residues in this entry.