



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 20, 2017 – 07:29 PM EST

PDB ID : 1PGN  
Title : CRYSTALLOGRAPHIC STUDY OF COENZYME, COENZYME ANALOGUE AND SUBSTRATE BINDING IN 6-PHOSPHOGLUCONATE DEHYDROGENASE: IMPLICATIONS FOR NADP SPECIFICITY AND THE ENZYME MECHANISM  
Authors : Adams, M.J.; Phillips, C.; Gover, S.  
Deposited on : unknown  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	POP	A	506	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 6-PHOSPHOGLUCONATE DEHYDROGENASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	473	3654	2332	634	666	22	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is NICOTINAMIDE 8-BROMO-ADENINE DINUCLEOTIDE PHOSPHATE (three-letter code: NBP) (formula: C<sub>21</sub>H<sub>27</sub>BrN<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	Br	C	N	O	P	0	0
			49	1	21	7	17	3		

- Molecule 4 is PYROPHOSPHATE 2- (three-letter code: POP) (formula:  $\text{H}_2\text{O}_7\text{P}_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			9	7	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	420	Total 420	O 420	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: 6-PHOSPHOGLUCONATE DEHYDROGENASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.74Å 148.40Å 102.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.30 19.73 – 2.30	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.30) 77.5 (19.73-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.71 (at 2.30Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.204 , (Not available) (Not available) , (Not available)	Depositor DCC
$R_{free}$ test set	NotAvailable	DCC
Wilson B-factor (Å <sup>2</sup> )	32.5	Xtriage
Anisotropy	0.486	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 84.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4142	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NBP, SO4, POP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.59	0/3731	1.28	19/5028 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	362	ARG	NE-CZ-NH1	9.06	124.83	120.30
1	A	406	ARG	NE-CZ-NH2	-8.32	116.14	120.30
1	A	433	ARG	CA-CB-CG	7.91	130.81	113.40
1	A	113	ARG	NE-CZ-NH2	-7.90	116.35	120.30
1	A	433	ARG	NE-CZ-NH1	7.67	124.14	120.30
1	A	406	ARG	NE-CZ-NH1	7.33	123.97	120.30
1	A	395	LYS	CA-CB-CG	6.84	128.45	113.40
1	A	15	GLN	CA-CB-CG	6.06	126.72	113.40
1	A	342	ARG	NE-CZ-NH1	5.74	123.17	120.30
1	A	207	VAL	CA-CB-CG2	-5.58	102.53	110.90
1	A	15	GLN	N-CA-CB	-5.51	100.69	110.60
1	A	162	LYS	CA-CB-CG	5.45	125.39	113.40
1	A	61	VAL	CG1-CB-CG2	-5.16	102.65	110.90
1	A	75	LYS	CA-C-N	-5.15	105.87	117.20
1	A	453	THR	CA-CB-CG2	-5.15	105.19	112.40
1	A	433	ARG	CB-CA-C	-5.09	100.22	110.40

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	113	ARG	NE-CZ-NH1	5.06	122.83	120.30
1	A	75	LYS	O-C-N	5.05	130.77	122.70
1	A	207	VAL	CG1-CB-CG2	-5.04	102.84	110.90

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	136	TYR	Sidechain
1	A	273	TYR	Sidechain
1	A	354	TYR	Sidechain
1	A	83	PHE	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3654	0	3661	56	0
2	A	10	0	0	0	0
3	A	49	0	24	3	0
4	A	9	0	0	1	0
5	A	420	0	0	10	0
All	All	4142	0	3685	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:SER:HB3	1:A:369:SER:HB3	1.60	0.83
1:A:362:ARG:HG2	1:A:362:ARG:HH11	1.52	0.75
1:A:83:PHE:CE2	3:A:499:NBP:H2A	2.32	0.64
1:A:430:ASP:HA	1:A:433:ARG:HD3	1.81	0.63
1:A:108:THR:OG1	1:A:180:HIS:HD2	1.83	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:SER:HA	4:A:506:POP:O5	2.02	0.59
1:A:83:PHE:HE2	3:A:499:NBP:H2A	1.67	0.59
1:A:405:TRP:O	1:A:409:ILE:HG12	2.01	0.59
1:A:446:ARG:HG2	1:A:446:ARG:HH11	1.70	0.57
1:A:229:SER:CB	1:A:369:SER:HB3	2.35	0.56
1:A:369:SER:HB2	5:A:606:HOH:O	2.07	0.54
1:A:49:THR:OG1	1:A:51:VAL:HG12	2.08	0.53
1:A:29:CYS:SG	1:A:54:ALA:HB2	2.49	0.53
1:A:109:MET:HG2	1:A:176:ASP:HB3	1.90	0.52
1:A:15:GLN:HG3	1:A:41:PHE:CE1	2.44	0.52
1:A:399:GLU:HA	1:A:402:GLN:HG2	1.91	0.52
1:A:57:LEU:HD21	1:A:83:PHE:HE1	1.74	0.52
1:A:459:LYS:N	1:A:460:PRO:HD3	2.25	0.51
1:A:57:LEU:O	1:A:61:VAL:HG23	2.10	0.51
1:A:279:LEU:HD13	1:A:333:SER:HB2	1.94	0.49
1:A:42:LEU:HD11	1:A:53:GLY:HA3	1.95	0.49
1:A:449:PHE:HA	5:A:841:HOH:O	2.10	0.49
1:A:280:ILE:O	1:A:284:VAL:HG23	2.12	0.49
1:A:86:LYS:O	1:A:89:PRO:HD2	2.13	0.48
1:A:231:LEU:HB2	5:A:605:HOH:O	2.12	0.48
1:A:16:ASN:HD22	1:A:135:ARG:HD3	1.78	0.47
1:A:213:LYS:HG3	5:A:636:HOH:O	2.14	0.47
1:A:392:ASP:HA	1:A:395:LYS:HG3	1.96	0.47
1:A:12:VAL:HG23	3:A:499:NBP:O7N	2.14	0.47
1:A:441:LEU:O	1:A:445:GLN:HG2	2.14	0.47
1:A:5:ILE:HD11	1:A:71:ILE:HD11	1.96	0.47
1:A:350:TRP:HA	5:A:598:HOH:O	2.15	0.47
1:A:419:MET:HB3	1:A:422:PHE:HB2	1.97	0.46
1:A:89:PRO:HB2	1:A:90:LEU:HD22	1.97	0.46
1:A:162:LYS:HA	1:A:167:GLU:O	2.17	0.45
1:A:186:HIS:HD2	1:A:187:ASN:OD1	2.00	0.44
1:A:374:LYS:HD2	1:A:374:LYS:N	2.32	0.44
1:A:84:ILE:O	1:A:88:VAL:HG23	2.16	0.44
1:A:47:LYS:HA	5:A:537:HOH:O	2.17	0.44
1:A:330:LYS:O	1:A:334:TYR:HD1	2.00	0.44
1:A:302:LYS:HD2	5:A:1146:HOH:O	2.18	0.43
1:A:106:ARG:HB2	5:A:1086:HOH:O	2.19	0.42
1:A:280:ILE:HD13	1:A:280:ILE:N	2.35	0.42
1:A:456:LEU:HA	1:A:456:LEU:HD12	1.88	0.42
1:A:90:LEU:HD12	5:A:767:HOH:O	2.20	0.42
1:A:321:ASP:HA	1:A:404:SER:HB3	2.00	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:293:LYS:HE2	1:A:297:ILE:HD11	2.02	0.41
1:A:417:ILE:HA	1:A:418:PRO:HD2	1.87	0.41
1:A:73:LEU:HD23	1:A:73:LEU:HA	1.80	0.41
1:A:235:THR:O	1:A:239:LEU:HD13	2.20	0.41
1:A:409:ILE:HD12	1:A:422:PHE:HB3	2.02	0.41
1:A:80:VAL:O	1:A:84:ILE:HG13	2.21	0.41
1:A:409:ILE:O	1:A:413:VAL:HG23	2.20	0.41
1:A:309:ILE:HB	5:A:1131:HOH:O	2.21	0.41
1:A:8:ILE:HG12	1:A:31:PHE:HB3	2.03	0.40
1:A:21:MET:HG2	1:A:159:ILE:HG21	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	471/482 (98%)	437 (93%)	30 (6%)	4 (1%)	22	26

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	164	GLY
1	A	469	THR
1	A	76	ALA
1	A	310	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	378/386 (98%)	339 (90%)	39 (10%)	8 9

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	12	VAL
1	A	19	LEU
1	A	33	ARG
1	A	62	SER
1	A	69	ARG
1	A	70	ILE
1	A	83	PHE
1	A	85	GLU
1	A	92	ASP
1	A	123	VAL
1	A	132	ASP
1	A	138	PRO
1	A	142	PRO
1	A	162	LYS
1	A	167	GLU
1	A	176	ASP
1	A	208	LEU
1	A	210	LEU
1	A	217	LYS
1	A	271	LEU
1	A	276	PRO
1	A	280	ILE
1	A	289	LEU
1	A	292	LEU
1	A	293	LYS
1	A	298	GLN
1	A	340	LEU
1	A	341	LEU
1	A	359	LEU
1	A	362	ARG
1	A	374	LYS
1	A	390	LEU
1	A	395	LYS
1	A	400	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	414	GLN
1	A	445	GLN
1	A	456	LEU
1	A	457	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	102	ASN
1	A	180	HIS
1	A	186	HIS
1	A	195	GLN
1	A	440	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NBP	A	499	-	45,53,53	1.29	4 (8%)	52,82,82	1.55	5 (9%)
2	SO4	A	505	-	4,4,4	0.67	0	6,6,6	0.25	0
4	POP	A	506	-	8,8,8	1.08	0	8,13,13	1.21	0
2	SO4	A	508	-	4,4,4	0.61	0	6,6,6	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NBP	A	499	-	-	0/27/67/67	0/5/5/5
2	SO4	A	505	-	-	0/0/0/0	0/0/0/0
4	POP	A	506	-	-	0/6/6/6	0/0/0/0
2	SO4	A	508	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	499	NBP	C5A-N7A	-4.98	1.32	1.38
3	A	499	NBP	C3N-C7N	2.27	1.54	1.50
3	A	499	NBP	C6N-N1N	2.54	1.42	1.35
3	A	499	NBP	O4D-C1D	3.62	1.46	1.41

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	499	NBP	N3A-C2A-N1A	-6.81	122.93	128.86
3	A	499	NBP	O4B-C1B-C2B	-3.20	100.99	106.59
3	A	499	NBP	C4B-O4B-C1B	-2.96	106.62	109.77
3	A	499	NBP	O7N-C7N-C3N	2.20	122.20	119.62
3	A	499	NBP	C4A-C5A-N7A	3.88	113.11	109.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	499	NBP	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	506	POP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	473/482 (98%)	-0.30	9 (1%) 67 73	12, 33, 47, 64	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	ALA	4.7
1	A	473	GLY	4.0
1	A	78	GLN	2.7
1	A	469	THR	2.6
1	A	471	HIS	2.4
1	A	310	PRO	2.3
1	A	472	GLY	2.1
1	A	309	ILE	2.0
1	A	93	ILE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors



of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	POP	A	506	9/9	0.95	0.15	2.11	55,59,60,60	0
3	NBP	A	499	49/49	0.93	0.14	0.31	29,39,47,49	48
2	SO4	A	505	5/5	0.98	0.10	-0.59	31,35,36,38	0
2	SO4	A	508	5/5	0.96	0.22	-	57,58,58,58	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.