



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 10, 2017 – 02:09 AM EDT

PDB ID : 2PG5
Title : Crystal Structure of Human Microsomal P450 2A6 N297Q
Authors : Sansen, S.; Hsu, M.H.; Stout, C.D.; Johnson, E.F.
Deposited on : unknown
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	<div> <div>5%</div> <div>78%</div> <div>17%</div> <div>• •</div> </div>
1	B	476	<div> <div>7%</div> <div>71%</div> <div>24%</div> <div>• •</div> </div>
1	C	476	<div> <div>4%</div> <div>78%</div> <div>18%</div> <div>• •</div> </div>
1	D	476	<div> <div>9%</div> <div>74%</div> <div>21%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	501	-	-	-	X
3	EDO	B	501	-	-	-	X
3	EDO	C	501	-	-	-	X
3	EDO	D	501	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 15846 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 2A6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	464	Total	C	N	O	S	0	0	0
			3752	2409	648	677	18			
1	B	464	Total	C	N	O	S	0	0	0
			3758	2413	650	677	18			
1	C	463	Total	C	N	O	S	0	0	0
			3748	2407	647	676	18			
1	D	464	Total	C	N	O	S	0	0	0
			3752	2409	648	677	18			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	23	MET	-	CLONING ARTIFACT	UNP P11509
A	24	ALA	-	CLONING ARTIFACT	UNP P11509
A	25	LYS	-	CLONING ARTIFACT	UNP P11509
A	26	LYS	-	CLONING ARTIFACT	UNP P11509
A	27	THR	-	CLONING ARTIFACT	UNP P11509
A	28	SER	-	CLONING ARTIFACT	UNP P11509
A	160	LEU	HIS	VARIANT	UNP P11509
A	297	GLN	ASN	ENGINEERED	UNP P11509
A	495	HIS	-	EXPRESSION TAG	UNP P11509
A	496	HIS	-	EXPRESSION TAG	UNP P11509
A	497	HIS	-	EXPRESSION TAG	UNP P11509
A	498	HIS	-	EXPRESSION TAG	UNP P11509
B	23	MET	-	CLONING ARTIFACT	UNP P11509
B	24	ALA	-	CLONING ARTIFACT	UNP P11509
B	25	LYS	-	CLONING ARTIFACT	UNP P11509
B	26	LYS	-	CLONING ARTIFACT	UNP P11509
B	27	THR	-	CLONING ARTIFACT	UNP P11509
B	28	SER	-	CLONING ARTIFACT	UNP P11509
B	160	LEU	HIS	VARIANT	UNP P11509
B	297	GLN	ASN	ENGINEERED	UNP P11509
B	495	HIS	-	EXPRESSION TAG	UNP P11509

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	496	HIS	-	EXPRESSION TAG	UNP P11509
B	497	HIS	-	EXPRESSION TAG	UNP P11509
B	498	HIS	-	EXPRESSION TAG	UNP P11509
C	23	MET	-	CLONING ARTIFACT	UNP P11509
C	24	ALA	-	CLONING ARTIFACT	UNP P11509
C	25	LYS	-	CLONING ARTIFACT	UNP P11509
C	26	LYS	-	CLONING ARTIFACT	UNP P11509
C	27	THR	-	CLONING ARTIFACT	UNP P11509
C	28	SER	-	CLONING ARTIFACT	UNP P11509
C	160	LEU	HIS	VARIANT	UNP P11509
C	297	GLN	ASN	ENGINEERED	UNP P11509
C	495	HIS	-	EXPRESSION TAG	UNP P11509
C	496	HIS	-	EXPRESSION TAG	UNP P11509
C	497	HIS	-	EXPRESSION TAG	UNP P11509
C	498	HIS	-	EXPRESSION TAG	UNP P11509
D	23	MET	-	CLONING ARTIFACT	UNP P11509
D	24	ALA	-	CLONING ARTIFACT	UNP P11509
D	25	LYS	-	CLONING ARTIFACT	UNP P11509
D	26	LYS	-	CLONING ARTIFACT	UNP P11509
D	27	THR	-	CLONING ARTIFACT	UNP P11509
D	28	SER	-	CLONING ARTIFACT	UNP P11509
D	160	LEU	HIS	VARIANT	UNP P11509
D	297	GLN	ASN	ENGINEERED	UNP P11509
D	495	HIS	-	EXPRESSION TAG	UNP P11509
D	496	HIS	-	EXPRESSION TAG	UNP P11509
D	497	HIS	-	EXPRESSION TAG	UNP P11509
D	498	HIS	-	EXPRESSION TAG	UNP P11509

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\text{C}_2\text{H}_6\text{O}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	158	Total O 158 158	0	0
4	B	134	Total O 134 134	0	0
4	C	201	Total O 201 201	0	0
4	D	155	Total O 155 155	0	0

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- [illegible]

- Chain B:
-
- 7% 71% 24%
- MET ALA LYS THR SER LYS GLY K32 L33 P34 Q48 E52 Q53 M54 Y55 N56 M59 K60 I61 S62 E63 R64 Y65 F69 L73 R77 E89 Q94 E97 R101 G102 E103 V117 R123 L135 R136 D137 F138 L143 G141 K142 R143 E147 P149
- S270 F271 L272 I273 I274 R275 Q276 M281 P282 T283 T284 E285 L291 V292 M293 Q297 I300 L310 R311 Y312 R313 F314 L315 L316 L317 R318 K319 E320 P321 E324 V327 I335 G336 R337 P341 R346 H347 L348 R352 E353 I356 S369 L370 R373 T378 K379 E380
- R381 D382 F383 F394 T389 E390 V391 Y392 L395 R400 N412 P413 Q414 L417 N418 E419 Q422 F423 K424 K425 F429 S433 R437 N438 Q439 E442 R446 F452 F453 N457 F460 R461 L462 S467 P468 R469 K476 H477 V478 Q479 F490 F491

- 
- WORLD WIDE
PDB
PROTEIN DATA BANK



• Molecule 1: Cytochrome P450 2A6



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.85Å 157.97Å 103.74Å 90.00° 92.23° 90.00°	Depositor
Resolution (Å)	35.00 – 1.95 29.38 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.7 (35.00-1.95) 98.8 (29.38-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 1.95Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.215 , 0.239 0.212 , 0.212	Depositor DCC
R_{free} test set	8204 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.425	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.022 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15846	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/3843	0.60	1/5176 (0.0%)
1	B	0.40	0/3850	0.63	0/5186
1	C	0.39	0/3839	0.61	0/5171
1	D	0.40	0/3843	0.61	1/5176 (0.0%)
All	All	0.39	0/15375	0.61	2/20709 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	104	GLN	N-CA-C	-5.54	96.04	111.00
1	A	417	LEU	CA-CB-CG	5.24	127.36	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3752	0	3736	73	0
1	B	3758	0	3740	101	0
1	C	3748	0	3733	83	0
1	D	3752	0	3736	99	0
2	A	43	0	30	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	43	0	30	4	0
2	C	43	0	30	4	0
2	D	43	0	30	3	0
3	A	4	0	6	0	0
3	B	4	0	6	0	0
3	C	4	0	6	0	0
3	D	4	0	6	0	0
4	A	158	0	0	5	0
4	B	134	0	0	4	1
4	C	201	0	0	4	1
4	D	155	0	0	2	0
All	All	15846	0	15089	356	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 356 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:418:ASN:HD22	1:C:422:GLN:HB2	1.14	1.11
1:B:234:GLN:HE21	1:B:234:GLN:H	1.02	0.97
1:C:45:ASN:HD22	1:C:72:HIS:H	1.02	0.95
1:A:54:MET:HG3	1:A:218:GLN:HE21	1.32	0.91
1:C:234:GLN:H	1:C:234:GLN:HE21	1.12	0.91

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:609:HOH:O	4:C:671:HOH:O[1_554]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	462/476 (97%)	448 (97%)	14 (3%)	0	100	100
1	B	462/476 (97%)	445 (96%)	16 (4%)	1 (0%)	51	41
1	C	461/476 (97%)	448 (97%)	13 (3%)	0	100	100
1	D	462/476 (97%)	442 (96%)	18 (4%)	2 (0%)	38	25
All	All	1847/1904 (97%)	1783 (96%)	61 (3%)	3 (0%)	51	41

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	140	VAL
1	D	261	PRO
1	B	261	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/422 (97%)	397 (97%)	14 (3%)	42	29
1	B	412/422 (98%)	393 (95%)	19 (5%)	31	16
1	C	411/422 (97%)	399 (97%)	12 (3%)	48	35
1	D	411/422 (97%)	396 (96%)	15 (4%)	40	26
All	All	1645/1688 (98%)	1585 (96%)	60 (4%)	40	26

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	316	LEU
1	C	50	ASN
1	D	312	TYR
1	B	417	LEU
1	C	143	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 52 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	45	ASN
1	C	167	ASN
1	D	283	ASN
1	C	50	ASN
1	C	56	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	500	1,4	28,50,50	2.16	11 (39%)	17,82,82	1.62	4 (23%)
3	EDO	A	501	-	3,3,3	0.55	0	2,2,2	0.32	0
2	HEM	B	500	1,4	28,50,50	2.12	10 (35%)	17,82,82	1.64	3 (17%)
3	EDO	B	501	-	3,3,3	0.62	0	2,2,2	0.32	0
2	HEM	C	500	1,4	28,50,50	2.13	13 (46%)	17,82,82	1.76	3 (17%)
3	EDO	C	501	-	3,3,3	0.57	0	2,2,2	0.32	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	D	500	1,4	28,50,50	2.16	11 (39%)	17,82,82	1.66	4 (23%)
3	EDO	D	501	-	3,3,3	0.65	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	1,4	-	0/6/54/54	0/0/8/8
3	EDO	A	501	-	-	0/1/1/1	0/0/0/0
2	HEM	B	500	1,4	-	0/6/54/54	0/0/8/8
3	EDO	B	501	-	-	0/1/1/1	0/0/0/0
2	HEM	C	500	1,4	-	0/6/54/54	0/0/8/8
3	EDO	C	501	-	-	0/1/1/1	0/0/0/0
2	HEM	D	500	1,4	-	0/6/54/54	0/0/8/8
3	EDO	D	501	-	-	0/1/1/1	0/0/0/0

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	HEM	C3C-CAC	-5.55	1.36	1.47
2	D	500	HEM	C3C-CAC	-5.11	1.37	1.47
2	B	500	HEM	C3C-CAC	-5.04	1.37	1.47
2	C	500	HEM	C3C-CAC	-4.94	1.37	1.47
2	B	500	HEM	C1C-NC	-4.15	1.31	1.36

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	500	HEM	C4C-C3C-C2C	-3.51	104.45	106.90
2	C	500	HEM	C4C-C3C-C2C	-3.29	104.60	106.90
2	B	500	HEM	C4C-C3C-C2C	-3.15	104.70	106.90
2	A	500	HEM	C4C-C3C-C2C	-2.85	104.91	106.90
2	D	500	HEM	CAA-CBA-CGA	2.03	116.13	112.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	3	0
2	B	500	HEM	4	0
2	C	500	HEM	4	0
2	D	500	HEM	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	464/476 (97%)	0.34	25 (5%) 26 37	23, 34, 44, 50	1 (0%)
1	B	464/476 (97%)	0.44	32 (6%) 18 27	26, 36, 45, 50	0
1	C	463/476 (97%)	0.28	18 (3%) 40 50	22, 33, 44, 50	0
1	D	464/476 (97%)	0.48	44 (9%) 9 14	23, 35, 47, 51	0
All	All	1855/1904 (97%)	0.38	119 (6%) 20 30	22, 34, 45, 51	1 (0%)

The worst 5 of 119 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	370	LEU	12.1
1	B	495	HIS	7.5
1	D	143	ARG	5.9
1	B	143	ARG	4.7
1	D	280	LYS	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	EDO	A	501	4/4	0.87	0.37	7.79	42,42,43,44	0
3	EDO	B	501	4/4	0.79	0.35	7.08	40,42,42,47	0
3	EDO	D	501	4/4	0.80	0.60	6.34	40,40,43,48	0
3	EDO	C	501	4/4	0.84	0.51	6.19	40,40,42,45	0
2	HEM	C	500	43/43	0.98	0.19	0.73	24,28,31,36	0
2	HEM	B	500	43/43	0.97	0.16	0.42	25,29,31,38	0
2	HEM	D	500	43/43	0.97	0.18	0.20	25,30,33,37	0
2	HEM	A	500	43/43	0.97	0.17	0.17	23,27,30,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.