



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 03:44 am GMT

PDB ID : 3PGH  
Title : CYCLOOXYGENASE-2 (PROSTAGLANDIN SYNTHASE-2) COM-  
PLEXED WITH A NON-SELECTIVE INHIBITOR, FLURBIPROFEN  
Authors : Kurumbail, R.; Stallings, W.  
Deposited on : 1996-12-18  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	587	
1	B	587	
1	C	587	
1	D	587	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 18304 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLOOXYGENASE-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	552	Total	C	N	O	S	0	0	0
			4473	2886	748	814	25			
1	B	552	Total	C	N	O	S	0	0	0
			4473	2886	748	814	25			
1	C	552	Total	C	N	O	S	0	0	0
			4473	2886	748	814	25			
1	D	552	Total	C	N	O	S	0	0	0
			4473	2886	748	814	25			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	310	GLN	ASN	CONFLICT	UNP Q05769
A	333	LYS	ARG	CONFLICT	UNP Q05769
B	310	GLN	ASN	CONFLICT	UNP Q05769
B	333	LYS	ARG	CONFLICT	UNP Q05769
C	310	GLN	ASN	CONFLICT	UNP Q05769
C	333	LYS	ARG	CONFLICT	UNP Q05769
D	310	GLN	ASN	CONFLICT	UNP Q05769
D	333	LYS	ARG	CONFLICT	UNP Q05769

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 4 is FLURBIPROFEN (three-letter code: FLP) (formula:  $C_{15}H_{13}FO_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	F	O	0	0
			18	15	1	2		
4	B	1	Total	C	F	O	0	0
			18	15	1	2		
4	C	1	Total	C	F	O	0	0
			18	15	1	2		
4	D	1	Total	C	F	O	0	0
			18	15	1	2		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

#### • Molecule 1: CYCLOOXYGENASE-2




#### • Molecule 1: CYCLOOXYGENASE-2






• Molecule 1: CYCLOOXYGENASE-2

Chain C: 51% 34% 8% 6%



• Molecule 1: CYCLOOXYGENASE-2

Chain D: 52% 34% 7% 6%





E398	E399	Q400	E401	Y402	S403	F404	K405	Y409	S412	T413	L414	L415	E416	H417	G418	L419	T420	Q421	F422	V423	E424	S425	R428	Q429	I430	R433	V434	R438	N439	V440	P441	I442	Q445	A446	V447	Q454	E457	N458	K459	Y460	E465	Y466	R469	Y475	F478	E479	E480
L481	T482	G483	E484	K485	E486	N487	A488	L491	K492	Y495	D499	V500	M501	E502	I503	Y504	P505	A506	V509	E510	K511	P512	R513	P514	I517	P518	L525	G526	A527	P528	L531	L534	Q543	K546	G551	G552	E553	V554	G555	F556	K557	Q565	C569	K573	G574		
C575	P576	F577	T578	S579	F580	N581	V582	Q583	ASP	PRO	GLN	PRO	THR	LYS	THR	ALA	THR	TLE	ASN	ALA	SER	ALA	SER	HIS	SER	ARG	LEU	ASP	ASP	TLE	ASN	PRO	THR	VAL	LEU	TLE	LYS	ARG	ARG	SER	THR	GLU	LEU				

## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	179.50Å 133.80Å 117.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.50	Depositor
% Data completeness (in resolution range)	63.1 (8.00-2.50)	Depositor
$R_{merge}$	0.11	Depositor
$R_{sym}$	0.11	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.236 , 0.316	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	18304	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLP, NAG, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.67	1/4600 (0.0%)	0.85	5/6237 (0.1%)
1	B	0.68	1/4600 (0.0%)	0.86	6/6237 (0.1%)
1	C	0.65	0/4600	0.86	5/6237 (0.1%)
1	D	0.67	1/4600 (0.0%)	0.85	6/6237 (0.1%)
All	All	0.67	3/18400 (0.0%)	0.86	22/24948 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	2
1	D	0	1
All	All	0	5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	313	CYS	CB-SG	-5.72	1.72	1.81
1	B	236	GLU	CG-CD	5.45	1.60	1.51
1	D	479	GLU	CG-CD	5.12	1.59	1.51

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	390	LEU	CA-CB-CG	6.90	131.17	115.30
1	B	287	VAL	N-CA-C	6.78	129.30	111.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	287	VAL	N-CA-C	6.43	128.35	111.00
1	A	390	LEU	CA-CB-CG	6.42	130.08	115.30
1	A	287	VAL	N-CA-C	6.42	128.32	111.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	147	TYR	Sidechain
1	B	147	TYR	Sidechain
1	C	147	TYR	Sidechain
1	C	91	TYR	Sidechain
1	D	147	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4473	0	4374	207	0
1	B	4473	0	4374	217	0
1	C	4473	0	4375	215	0
1	D	4473	0	4375	201	0
2	A	42	0	39	4	0
2	B	42	0	39	3	0
2	C	42	0	39	0	0
2	D	42	0	39	0	0
3	A	43	0	30	6	0
3	B	43	0	30	9	0
3	C	43	0	30	6	0
3	D	43	0	30	9	0
4	A	18	0	12	1	0
4	B	18	0	12	6	0
4	C	18	0	12	3	0
4	D	18	0	12	3	0
All	All	18304	0	17822	829	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 829 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:424:GLU:HA	1:C:428:ARG:HH21	1.27	0.97
1:A:458:MET:HE2	1:A:460:TYR:HE1	1.27	0.96
1:A:424:GLU:HA	1:A:428:ARG:HH21	1.31	0.96
1:D:424:GLU:HA	1:D:428:ARG:HH21	1.30	0.95
1:B:479:GLU:HG2	1:B:485:LYS:NZ	1.82	0.95

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	550/587 (94%)	473 (86%)	62 (11%)	15 (3%)	6	9
1	B	550/587 (94%)	484 (88%)	49 (9%)	17 (3%)	5	6
1	C	550/587 (94%)	479 (87%)	54 (10%)	17 (3%)	5	6
1	D	550/587 (94%)	477 (87%)	57 (10%)	16 (3%)	5	7
All	All	2200/2348 (94%)	1913 (87%)	222 (10%)	65 (3%)	5	7

5 of 65 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	52	PHE
1	A	82	LEU
1	A	130	TYR
1	A	138	SER
1	A	282	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/525 (94%)	436 (88%)	57 (12%)	6	12
1	B	493/525 (94%)	437 (89%)	56 (11%)	7	12
1	C	493/525 (94%)	437 (89%)	56 (11%)	7	12
1	D	493/525 (94%)	437 (89%)	56 (11%)	7	12
All	All	1972/2100 (94%)	1747 (89%)	225 (11%)	7	12

5 of 225 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	484	GLU
1	C	165	VAL
1	D	422	PHE
1	B	518	PHE
1	C	52	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 76 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	417	HIS
1	C	232	HIS
1	D	396	ASN
1	B	454	GLN
1	C	95	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NAG	A	661	1	14,14,15	0.58	0	15,19,21	0.92	1 (6%)
2	NAG	A	671	1	14,14,15	0.72	0	15,19,21	1.71	5 (33%)
2	NAG	A	681	1	14,14,15	0.77	0	15,19,21	0.75	0
3	HEM	A	682	1	28,50,50	1.93	6 (21%)	17,82,82	1.17	0
4	FLP	A	701	-	16,19,19	1.53	3 (18%)	22,26,26	1.35	4 (18%)
2	NAG	B	661	1	14,14,15	0.51	0	15,19,21	1.06	1 (6%)
2	NAG	B	671	1	14,14,15	0.77	0	15,19,21	1.50	3 (20%)
2	NAG	B	681	1	14,14,15	0.73	0	15,19,21	0.90	0
3	HEM	B	682	1	28,50,50	1.96	6 (21%)	17,82,82	1.23	2 (11%)
4	FLP	B	701	-	16,19,19	1.80	4 (25%)	22,26,26	1.42	4 (18%)
2	NAG	C	661	1	14,14,15	0.26	0	15,19,21	0.90	1 (6%)
2	NAG	C	671	1	14,14,15	0.82	1 (7%)	15,19,21	1.66	5 (33%)
2	NAG	C	681	1	14,14,15	0.60	0	15,19,21	0.80	0
3	HEM	C	682	1	28,50,50	1.83	6 (21%)	17,82,82	1.27	1 (5%)
4	FLP	C	701	-	16,19,19	1.10	1 (6%)	22,26,26	1.40	3 (13%)
2	NAG	D	661	1	14,14,15	0.66	0	15,19,21	0.95	1 (6%)
2	NAG	D	671	1	14,14,15	0.63	0	15,19,21	1.59	3 (20%)
2	NAG	D	681	1	14,14,15	0.78	0	15,19,21	0.83	0
3	HEM	D	682	1	28,50,50	1.92	7 (25%)	17,82,82	1.00	1 (5%)
4	FLP	D	701	-	16,19,19	1.58	3 (18%)	22,26,26	1.28	4 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	661	1	-	0/6/23/26	0/1/1/1
2	NAG	A	671	1	-	0/6/23/26	0/1/1/1
2	NAG	A	681	1	-	0/6/23/26	0/1/1/1
3	HEM	A	682	1	-	0/6/54/54	0/0/8/8
4	FLP	A	701	-	-	0/8/12/12	0/2/2/2
2	NAG	B	661	1	-	0/6/23/26	0/1/1/1
2	NAG	B	671	1	-	0/6/23/26	0/1/1/1
2	NAG	B	681	1	-	0/6/23/26	0/1/1/1
3	HEM	B	682	1	-	0/6/54/54	0/0/8/8
4	FLP	B	701	-	-	0/8/12/12	0/2/2/2
2	NAG	C	661	1	-	0/6/23/26	0/1/1/1
2	NAG	C	671	1	-	0/6/23/26	0/1/1/1
2	NAG	C	681	1	-	0/6/23/26	0/1/1/1
3	HEM	C	682	1	-	0/6/54/54	0/0/8/8
4	FLP	C	701	-	-	0/8/12/12	0/2/2/2
2	NAG	D	661	1	-	0/6/23/26	0/1/1/1
2	NAG	D	671	1	-	0/6/23/26	0/1/1/1
2	NAG	D	681	1	-	0/6/23/26	0/1/1/1
3	HEM	D	682	1	-	0/6/54/54	0/0/8/8
4	FLP	D	701	-	-	0/8/12/12	0/2/2/2

The worst 5 of 37 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	682	HEM	C3C-CAC	-5.28	1.37	1.47
3	B	682	HEM	C3B-CAB	-5.10	1.37	1.47
3	A	682	HEM	C3C-C2C	-4.98	1.33	1.40
3	A	682	HEM	C3C-CAC	-4.89	1.38	1.47
3	C	682	HEM	C3C-C2C	-4.83	1.34	1.40

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	701	FLP	C2-C6-C11	-3.34	118.70	123.02
2	A	671	NAG	C4-C3-C2	-3.31	106.17	111.02
2	B	671	NAG	C4-C3-C2	-3.29	106.19	111.02
2	D	671	NAG	C4-C3-C2	-3.05	106.55	111.02
2	A	671	NAG	O5-C1-C2	-2.94	107.39	111.47

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

12 monomers are involved in 50 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	671	NAG	4	0
3	A	682	HEM	6	0
4	A	701	FLP	1	0
2	B	661	NAG	1	0
2	B	671	NAG	1	0
2	B	681	NAG	1	0
3	B	682	HEM	9	0
4	B	701	FLP	6	0
3	C	682	HEM	6	0
4	C	701	FLP	3	0
3	D	682	HEM	9	0
4	D	701	FLP	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.