



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 15, 2017 – 10:59 AM EDT

PDB ID : 4PHT  
Title : ATPase GspE in complex with the cytoplasmic domain of GspL from the  
Vibrio vulnificus type II Secretion system  
Authors : Lu, C.; Korotkov, K.; Hol, W.  
Deposited on : unknown  
Resolution : 2.83 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20029824
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3466 (2.88-2.80)
Clashscore	112137	3975 (2.88-2.80)
Ramachandran outliers	110173	3902 (2.88-2.80)
Sidechain outliers	110143	3905 (2.88-2.80)
RSRZ outliers	101464	3501 (2.88-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	246	
1	Y	246	
1	Z	246	
2	A	500	
2	B	500	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	C	500	<div><div><div>%</div><div><div></div><div>83%</div><div>5%</div><div>12%</div></div></div></div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 15452 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type II secretion system protein L.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	X	224	Total	C	N	O	S	0	0	0
			1739	1106	286	339	8			
1	Y	226	Total	C	N	O	S	0	0	0
			1742	1111	288	335	8			
1	Z	226	Total	C	N	O	S	0	0	0
			1732	1105	284	335	8			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	0	MET	-	initiating methionine	UNP Q8DDT8
X	238	LEU	-	expression tag	UNP Q8DDT8
X	239	GLU	-	expression tag	UNP Q8DDT8
X	240	HIS	-	expression tag	UNP Q8DDT8
X	241	HIS	-	expression tag	UNP Q8DDT8
X	242	HIS	-	expression tag	UNP Q8DDT8
X	243	HIS	-	expression tag	UNP Q8DDT8
X	244	HIS	-	expression tag	UNP Q8DDT8
X	245	HIS	-	expression tag	UNP Q8DDT8
Y	0	MET	-	initiating methionine	UNP Q8DDT8
Y	238	LEU	-	expression tag	UNP Q8DDT8
Y	239	GLU	-	expression tag	UNP Q8DDT8
Y	240	HIS	-	expression tag	UNP Q8DDT8
Y	241	HIS	-	expression tag	UNP Q8DDT8
Y	242	HIS	-	expression tag	UNP Q8DDT8
Y	243	HIS	-	expression tag	UNP Q8DDT8
Y	244	HIS	-	expression tag	UNP Q8DDT8
Y	245	HIS	-	expression tag	UNP Q8DDT8
Z	0	MET	-	initiating methionine	UNP Q8DDT8
Z	238	LEU	-	expression tag	UNP Q8DDT8
Z	239	GLU	-	expression tag	UNP Q8DDT8
Z	240	HIS	-	expression tag	UNP Q8DDT8
Z	241	HIS	-	expression tag	UNP Q8DDT8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
Z	242	HIS	-	expression tag	UNP Q8DDT8
Z	243	HIS	-	expression tag	UNP Q8DDT8
Z	244	HIS	-	expression tag	UNP Q8DDT8
Z	245	HIS	-	expression tag	UNP Q8DDT8

- Molecule 2 is a protein called General secretory pathway protein E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	438	Total	C	N	O	S	0	0	0
			3359	2118	598	626	17			
2	B	442	Total	C	N	O	S	0	0	0
			3385	2136	600	632	17			
2	C	441	Total	C	N	O	S	0	0	0
			3380	2132	598	633	17			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q8DDT1
B	0	MET	-	initiating methionine	UNP Q8DDT1
C	0	MET	-	initiating methionine	UNP Q8DDT1

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>6</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
4	B	1	Total 31	C 10	N 6	O 12	P 3	0	0
4	C	1	Total 31	C 10	N 6	O 12	P 3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Mg 1 1	0	0
5	A	1	Total Mg 1 1	0	0
5	C	1	Total Mg 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	X	3	Total O 3 3	0	0
6	A	1	Total O 1 1	0	0
6	Y	1	Total O 1 1	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	5	Total	O	0	0
			5	5		
6	C	6	Total	O	0	0
			6	6		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Type II secretion system protein L

Chain X: 





- Molecule 1: Type II secretion system protein L

Chain Y: 





- Molecule 1: Type II secretion system protein L

Chain Z: 




- Molecule 2: General secretory pathway protein E

Chain A: 



- Molecule 2: General secretory pathway protein E





GLOBAL-STATISTICS INFOmissingINFO

## 4 Model quality [i](#)

### 4.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ANP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	X	0.43	0/1778	0.54	0/2421
1	Y	0.44	0/1782	0.55	1/2428 (0.0%)
1	Z	0.41	0/1772	0.54	1/2417 (0.0%)
2	A	0.41	0/3404	0.53	0/4601
2	B	0.39	0/3430	0.54	0/4637
2	C	0.40	0/3425	0.54	0/4631
All	All	0.41	0/15591	0.54	2/21135 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	Z	134	ARG	NE-CZ-NH1	5.33	122.97	120.30
1	Y	134	ARG	NE-CZ-NH1	5.05	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

### 4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	1739	0	1672	9	0
1	Y	1742	0	1679	23	0
1	Z	1732	0	1657	24	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	3359	0	3403	12	0
2	B	3385	0	3427	12	0
2	C	3380	0	3419	12	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	31	0	13	0	0
4	B	31	0	13	0	0
4	C	31	0	13	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
6	A	1	0	0	0	0
6	B	5	0	0	0	0
6	C	6	0	0	0	0
6	X	3	0	0	0	0
6	Y	1	0	0	0	0
All	All	15452	0	15296	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 85 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:475:ARG:NH1	2:B:493:MET:SD	1.97	1.38
2:A:34:PRO:N	2:A:35:PRO:HD3	1.84	0.93
1:Z:149:ILE:CD1	1:Z:188:LEU:CD2	2.50	0.89
2:A:287:VAL:CG2	2:A:318:ILE:HD12	2.01	0.89
1:Z:149:ILE:HD12	1:Z:188:LEU:HD23	1.58	0.83

There are no symmetry-related clashes.

## 4.3 Torsion angles [i](#)

### 4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	X	218/246 (89%)	213 (98%)	4 (2%)	1 (0%)	32	64
1	Y	220/246 (89%)	216 (98%)	4 (2%)	0	100	100
1	Z	220/246 (89%)	216 (98%)	4 (2%)	0	100	100
2	A	426/500 (85%)	414 (97%)	11 (3%)	1 (0%)	51	81
2	B	430/500 (86%)	420 (98%)	8 (2%)	2 (0%)	32	64
2	C	429/500 (86%)	418 (97%)	10 (2%)	1 (0%)	51	81
All	All	1943/2238 (87%)	1897 (98%)	41 (2%)	5 (0%)	44	75

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	278	ASN
2	B	278	ASN
2	C	278	ASN
1	X	49	GLU
2	B	296	ASP

#### 4.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	X	190/212 (90%)	188 (99%)	2 (1%)	78	93
1	Y	189/212 (89%)	188 (100%)	1 (0%)	91	97
1	Z	187/212 (88%)	186 (100%)	1 (0%)	91	97
2	A	362/434 (83%)	358 (99%)	4 (1%)	78	93
2	B	365/434 (84%)	357 (98%)	8 (2%)	57	85
2	C	365/434 (84%)	357 (98%)	8 (2%)	57	85
All	All	1658/1938 (86%)	1634 (99%)	24 (1%)	71	91

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	217	ARG
2	B	431	HIS
2	C	431	HIS
2	B	318	ILE
2	B	429	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	50	GLN
2	A	386	GLN
1	Y	200	GLN
2	B	386	GLN
2	C	386	GLN

#### 4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

#### 4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 4.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

#### 4.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ANP	A	602	5	29,33,33	1.46	3 (10%)	28,52,52	1.70	6 (21%)
4	ANP	B	602	5	29,33,33	2.75	4 (13%)	28,52,52	1.78	6 (21%)
4	ANP	C	602	5	29,33,33	1.91	4 (13%)	28,52,52	1.73	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ANP	A	602	5	-	0/13/38/38	0/3/3/3
4	ANP	B	602	5	-	0/13/38/38	0/3/3/3
4	ANP	C	602	5	-	0/13/38/38	0/3/3/3

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	602	ANP	PG-O3G	-3.84	1.46	1.56
4	C	602	ANP	PB-O2B	-3.35	1.47	1.56
4	A	602	ANP	PG-O2G	-3.32	1.47	1.56
4	B	602	ANP	PB-O2B	-3.30	1.47	1.56
4	C	602	ANP	PG-O2G	-2.47	1.50	1.56

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	602	ANP	O1G-PG-N3B	-5.27	103.91	111.79
4	C	602	ANP	O1G-PG-N3B	-4.44	105.15	111.79
4	A	602	ANP	O1G-PG-N3B	-4.19	105.52	111.79
4	C	602	ANP	O1B-PB-N3B	-3.27	106.90	111.79
4	B	602	ANP	O1B-PB-N3B	-3.19	107.02	111.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	602	ANP	1	0

## 4.7 Other polymers [i](#)

There are no such residues in this entry.

## 4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 5 Fit of model and data [i](#)

### 5.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	X	224/246 (91%)	-0.22	0 100 100	27, 46, 78, 90	0
1	Y	226/246 (91%)	-0.12	1 (0%) 92 90	29, 47, 78, 107	0
1	Z	226/246 (91%)	-0.05	1 (0%) 92 90	33, 54, 88, 106	0
2	A	438/500 (87%)	-0.04	4 (0%) 84 80	31, 59, 96, 112	0
2	B	442/500 (88%)	-0.04	3 (0%) 87 84	34, 58, 97, 123	0
2	C	441/500 (88%)	-0.00	4 (0%) 84 80	31, 59, 101, 132	0
All	All	1997/2238 (89%)	-0.06	13 (0%) 87 84	27, 56, 95, 132	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	418	LEU	4.1
2	A	417	ALA	4.0
2	C	26	LEU	2.6
2	A	409	LEU	2.6
2	C	103	GLU	2.5

### 5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 5.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ZN	A	601	1/1	0.99	0.15	-0.08	50,50,50,50	0
3	ZN	C	601	1/1	0.97	0.15	-0.28	64,64,64,64	0
4	ANP	B	602	31/31	0.97	0.16	-0.45	36,40,43,45	0
4	ANP	C	602	31/31	0.98	0.16	-0.68	26,36,42,44	0
4	ANP	A	602	31/31	0.97	0.16	-0.72	29,35,44,50	0
3	ZN	B	601	1/1	0.99	0.11	-1.73	54,54,54,54	0
5	MG	C	603	1/1	0.94	0.17	-	27,27,27,27	0
5	MG	B	603	1/1	0.93	0.20	-	41,41,41,41	0
5	MG	A	603	1/1	0.94	0.19	-	24,24,24,24	0

## 5.5 Other polymers

There are no such residues in this entry.