



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2017 – 01:25 am GMT

PDB ID : 4PJI  
Title : Structure of human MR1-Ac-6-FP in complex with human MAIT C-C10 TCR  
Authors : Birkinshaw, R.W.; Rossjohn, J.  
Deposited on : 2014-05-12  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	<div> <div style="width: 91%;"></div> <div style="width: 6%;"></div> <div style="width: 3%;"></div> </div>
1	C	271	<div> <div style="width: 89%;"></div> <div style="width: 7%;"></div> <div style="width: 3%;"></div> </div>
2	B	100	<div> <div style="width: 97%;"></div> <div style="width: 2%;"></div> <div style="width: 1%;"></div> </div>
2	D	100	<div> <div style="width: 96%;"></div> <div style="width: 6%;"></div> <div style="width: 1%;"></div> </div>
3	E	205	<div> <div style="width: 92%;"></div> <div style="width: 8%;"></div> <div style="width: 1%;"></div> </div>
3	G	205	<div> <div style="width: 89%;"></div> <div style="width: 8%;"></div> <div style="width: 5%;"></div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	245	<div><div>%</div><div><div></div><div>91%</div><div>6%</div><div></div></div><div></div></div>
4	H	245	<div><div>3%</div><div><div></div><div>91%</div><div>6%</div><div></div></div><div></div></div>

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 13007 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major histocompatibility complex class I-related gene protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	262	Total	C	N	O	S	0	0	0
			2155	1381	371	393	10			
1	C	251	Total	C	N	O	S	0	0	0
			2070	1326	355	379	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q95460
A	261	SER	CYS	engineered mutation	UNP Q95460
C	0	MET	-	initiating methionine	UNP Q95460
C	261	SER	CYS	engineered mutation	UNP Q95460

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			812	517	137	155	3			
2	D	99	Total	C	N	O	S	0	0	0
			790	506	131	150	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
D	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called TCR-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	198	Total	C	N	O	S	0	0	0
			1509	958	240	302	9			

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	189	Total	C	N	O	S	0	0	0
			1423	914	225	275	9			

- Molecule 4 is a protein called TCR-beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	240	Total	C	N	O	S	0	0	0
			1853	1166	321	357	9			
4	H	238	Total	C	N	O	S	0	6	0
			1872	1179	322	362	9			

- Molecule 5 is N-(6-formyl-4-oxo-3,4-dihydropteridin-2-yl)acetamide (three-letter code: 30W) (formula: C<sub>9</sub>H<sub>7</sub>N<sub>5</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			16	9	5	2		
5	C	1	Total	C	N	O	0	0
			16	9	5	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	H	1	Total	Na	0	0
			1	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	96	Total	O	0	0
			96	96		
8	B	33	Total	O	0	0
			33	33		
8	C	80	Total	O	0	0
			80	80		
8	D	25	Total	O	0	0
			25	25		
8	E	57	Total	O	0	0
			57	57		
8	F	75	Total	O	0	0
			75	75		
8	G	44	Total	O	0	0
			44	44		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	H	74	Total	O	0	0
			74	74		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major histocompatibility complex class I-related gene protein



- Molecule 1: Major histocompatibility complex class I-related gene protein



- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin




- Molecule 3: TCR-alpha




- Molecule 3: TCR-alpha






● Molecule 4: TCR-beta



● Molecule 4: TCR-beta



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	217.69Å 71.02Å 144.98Å 90.00° 104.76° 90.00°	Depositor
Resolution (Å)	36.60 – 2.50 36.28 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.2 (36.60-2.50) 99.2 (36.28-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.84 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, $R_{free}$	0.188 , 0.228 0.189 , 0.229	Depositor DCC
$R_{free}$ test set	3726 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.9	Xtriage
Anisotropy	0.255	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	13007	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NA, 30W

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/2219	0.66	0/3014
1	C	0.52	0/2131	0.65	0/2893
2	B	0.51	0/835	0.66	0/1135
2	D	0.44	0/813	0.65	0/1109
3	E	0.51	0/1543	0.67	0/2100
3	G	0.52	0/1455	0.67	0/1980
4	F	0.51	0/1904	0.70	0/2600
4	H	0.50	0/1925	0.72	2/2630 (0.1%)
All	All	0.51	0/12825	0.68	2/17461 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	98[A]	GLY	C-N-CA	5.79	136.17	121.70
4	H	98[B]	GLY	C-N-CA	5.79	136.17	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2155	0	2042	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2070	0	1952	3	0
2	B	812	0	757	0	0
2	D	790	0	718	0	0
3	E	1509	0	1391	4	0
3	G	1423	0	1306	2	0
4	F	1853	0	1722	6	0
4	H	1872	0	1733	14	0
5	A	16	0	6	0	0
5	C	16	0	6	0	0
6	B	6	0	8	0	0
7	H	1	0	0	0	0
8	A	96	0	0	2	0
8	B	33	0	0	0	0
8	C	80	0	0	0	0
8	D	25	0	0	0	0
8	E	57	0	0	1	0
8	F	75	0	0	0	0
8	G	44	0	0	0	0
8	H	74	0	0	0	0
All	All	13007	0	11641	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:98[A]:GLY:HA2	4:H:99[A]:THR:HG22	1.67	0.77
1:A:127:LEU:HG	8:A:404:HOH:O	1.88	0.71
3:E:91:VAL:HG11	4:F:99:THR:HG22	1.72	0.69
3:E:159:LEU:HB3	4:F:170:CYS:HB3	1.78	0.65
4:H:98[B]:GLY:HA2	4:H:99[B]:THR:HG22	1.84	0.60
4:H:98[A]:GLY:HA3	4:H:100[A]:ASP:H	1.68	0.58
3:E:92:ASP:HB2	8:E:354:HOH:O	2.04	0.56
4:H:98[A]:GLY:HA3	4:H:100[A]:ASP:N	2.22	0.55
3:G:159:LEU:HB3	4:H:170:CYS:HB2	1.89	0.55
1:A:77:LEU:HD13	1:A:92:TYR:HB2	1.90	0.53
3:E:28:GLY:HA3	3:E:93:SER:HB3	1.89	0.52
4:H:95[B]:PRO:HD2	4:H:100[B]:ASP:HB2	1.92	0.51
4:H:98[B]:GLY:HA3	4:H:100[B]:ASP:N	2.28	0.48
4:H:129:PRO:HG3	4:H:140:ALA:HB1	1.95	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:HIS:HB3	4:H:99[B]:THR:OG1	2.14	0.48
1:C:9:ARG:HB2	1:C:94:ARG:HB3	1.96	0.47
4:H:98[A]:GLY:CA	4:H:99[A]:THR:HG22	2.42	0.46
4:F:129:PRO:HD3	4:F:142:LEU:HG	1.98	0.46
4:H:83:PRO:HA	4:H:112:VAL:HB	1.98	0.46
3:G:109:PRO:HG3	3:G:158:VAL:HG11	1.97	0.46
1:A:215:MET:HG3	1:A:257:HIS:CD2	2.53	0.44
4:F:92:ALA:HA	4:F:103:TYR:O	2.18	0.43
4:H:95[A]:PRO:HD2	4:H:100[A]:ASP:HB3	2.00	0.43
1:C:169:LEU:HD23	1:C:176:LEU:HD13	2.01	0.43
4:F:83:PRO:HA	4:F:112:VAL:HB	2.00	0.43
1:C:215:MET:HG3	1:C:257:HIS:CD2	2.54	0.43
1:A:223:GLN:HG3	8:A:460:HOH:O	2.20	0.42
1:A:186:VAL:HG11	1:A:269:VAL:HG22	2.02	0.42
4:H:98[B]:GLY:CA	4:H:99[B]:THR:HG22	2.47	0.42
4:F:149:PHE:CE1	4:F:187:TYR:HB2	2.55	0.41
4:H:129:PRO:HD3	4:H:142:LEU:HG	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/271 (94%)	252 (98%)	4 (2%)	0	100	100
1	C	243/271 (90%)	238 (98%)	5 (2%)	0	100	100
2	B	97/100 (97%)	96 (99%)	1 (1%)	0	100	100
2	D	97/100 (97%)	96 (99%)	1 (1%)	0	100	100
3	E	196/205 (96%)	192 (98%)	4 (2%)	0	100	100
3	G	183/205 (89%)	178 (97%)	5 (3%)	0	100	100
4	F	238/245 (97%)	232 (98%)	6 (2%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	H	242/245 (99%)	235 (97%)	5 (2%)	2 (1%)	22	39
All	All	1552/1642 (94%)	1519 (98%)	31 (2%)	2 (0%)	55	76

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	H	96[A]	PRO
4	H	96[B]	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/241 (95%)	222 (97%)	6 (3%)	51	78
1	C	219/241 (91%)	214 (98%)	5 (2%)	56	81
2	B	90/95 (95%)	88 (98%)	2 (2%)	57	82
2	D	84/95 (88%)	81 (96%)	3 (4%)	40	67
3	E	162/182 (89%)	158 (98%)	4 (2%)	53	79
3	G	146/182 (80%)	142 (97%)	4 (3%)	50	77
4	F	197/212 (93%)	190 (96%)	7 (4%)	40	67
4	H	199/212 (94%)	194 (98%)	5 (2%)	53	79
All	All	1325/1460 (91%)	1289 (97%)	36 (3%)	50	77

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	ASP
1	A	66	LEU
1	A	73	PHE
1	A	221	ILE
1	A	224	GLU
1	A	246	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	4	THR
2	B	70	PHE
1	C	16	ILE
1	C	57	ASP
1	C	73	PHE
1	C	82	ARG
1	C	183	LEU
2	D	4	THR
2	D	34	ASP
2	D	70	PHE
3	E	27	SER
3	E	62	SER
3	E	130	SER
3	E	149	SER
4	F	51	SER
4	F	100	ASP
4	F	126	VAL
4	F	170	CYS
4	F	183	ASN
4	F	192	ARG
4	F	204	ARG
3	G	27	SER
3	G	77	GLU
3	G	164	MET
3	G	182	CYS
4	H	14	LYS
4	H	51	SER
4	H	101	THR
4	H	126	VAL
4	H	192	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
3	E	145	GLN
4	H	206	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	30W	A	301	1	16,17,18	1.53	3 (18%)	20,24,25	3.13	7 (35%)
6	GOL	B	101	-	5,5,5	0.23	0	5,5,5	0.52	0
5	30W	C	301	1	16,17,18	2.09	3 (18%)	20,24,25	3.07	8 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	30W	A	301	1	-	0/4/4/6	0/2/2/2
6	GOL	B	101	-	-	0/4/4/4	0/0/0/0
5	30W	C	301	1	-	0/4/4/6	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	301	30W	C2-N2	-4.87	1.32	1.38
5	A	301	30W	C2-N2	-3.61	1.34	1.38
5	A	301	30W	C2-N1	2.12	1.41	1.34
5	A	301	30W	C4-N3	2.79	1.38	1.33

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	301	30W	C4-N3	3.16	1.38	1.33
5	C	301	30W	C4-C4A	4.83	1.50	1.41

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	301	30W	C4A-C4-N3	-6.39	114.38	123.48
5	A	301	30W	N1-C2-N3	-6.31	116.69	126.23
5	A	301	30W	C4A-C4-N3	-5.82	115.19	123.48
5	C	301	30W	N1-C2-N3	-5.17	118.42	126.23
5	A	301	30W	C2-N2-C10	-4.42	123.45	130.28
5	C	301	30W	N2-C2-N1	-2.26	109.57	116.49
5	C	301	30W	C2-N2-C10	-2.22	126.86	130.28
5	C	301	30W	C4-C4A-N5	2.24	121.14	118.68
5	A	301	30W	N8-C8A-N1	2.66	119.47	116.04
5	A	301	30W	N2-C2-N3	4.40	129.65	116.44
5	A	301	30W	C2-N1-C8A	4.70	120.48	115.11
5	C	301	30W	N2-C2-N3	5.10	131.77	116.44
5	C	301	30W	C2-N1-C8A	5.40	121.27	115.11
5	C	301	30W	C4-N3-C2	6.42	126.67	115.18
5	A	301	30W	C4-N3-C2	6.94	127.62	115.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	262/271 (96%)	-0.22	4 (1%) 74 75	9, 23, 50, 66	0
1	C	251/271 (92%)	-0.07	7 (2%) 53 56	12, 25, 58, 85	0
2	B	99/100 (99%)	-0.18	2 (2%) 65 67	10, 26, 56, 75	0
2	D	99/100 (99%)	0.28	6 (6%) 22 22	15, 43, 82, 112	0
3	E	198/205 (96%)	-0.15	2 (1%) 82 83	12, 27, 54, 70	0
3	G	189/205 (92%)	0.12	11 (5%) 24 24	9, 34, 78, 96	0
4	F	240/245 (97%)	-0.22	3 (1%) 77 78	13, 27, 51, 74	0
4	H	238/245 (97%)	-0.05	7 (2%) 52 55	11, 29, 68, 98	0
All	All	1576/1642 (95%)	-0.09	42 (2%) 55 58	9, 27, 65, 112	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	98	ASP	5.2
1	C	246	LEU	4.7
4	H	219	ASN	4.3
1	A	222	VAL	3.7
4	H	203	PRO	3.5
2	D	76	ASP	3.4
3	G	179	ASP	3.4
4	H	97[A]	GLY	3.2
1	A	104	GLY	3.2
1	C	218	GLY	3.1
4	H	96[A]	PRO	3.0
3	E	182	CYS	2.9
1	C	224	GLU	2.9
3	G	196	PHE	2.9
3	G	180	PHE	2.9
1	A	223	GLN	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	244	ILE	2.7
2	B	0	MET	2.7
4	F	181	ALA	2.6
2	D	75	LYS	2.6
4	F	26	ASP	2.6
4	H	100[A]	ASP	2.6
3	G	149	SER	2.6
4	H	223	THR	2.6
4	H	227	ALA	2.6
3	G	191	ILE	2.5
2	D	96	ASP	2.4
2	D	97	ARG	2.4
3	G	198	PRO	2.4
1	A	18	GLY	2.3
3	E	165	ASP	2.3
3	G	190	ILE	2.3
1	C	103	ASP	2.2
2	D	77	GLU	2.2
4	F	182	LEU	2.2
3	G	132	CYS	2.2
1	C	197	ALA	2.1
3	G	163	SER	2.1
3	G	199	SER	2.1
3	G	119	TYR	2.1
2	B	47	GLU	2.1
1	C	245	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	30W	A	301	16/17	0.98	0.13	-0.15	10,14,21,26	0
5	30W	C	301	16/17	0.98	0.12	-0.20	20,22,26,28	0
6	GOL	B	101	6/6	0.98	0.12	-0.35	15,17,20,24	0
7	NA	H	301	1/1	0.93	0.13	-0.66	31,31,31,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.