



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 12, 2017 – 07:45 PM EDT

PDB ID : 2PO6
Title : Crystal structure of CD1d-lipid-antigen complexed with Beta-2-Microglobulin, NKT15 Alpha-Chain and NKT15 Beta-Chain
Authors : Borg, N.A.
Deposited on : unknown
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1015 (3.22-3.18)
Clashscore	112137	1009 (3.20-3.20)
Ramachandran outliers	110173	1118 (3.22-3.18)
Sidechain outliers	110143	1117 (3.22-3.18)
RSRZ outliers	101464	1020 (3.22-3.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	278	<div> <div>4%</div> <div> <div></div> <div>71%</div> <div>26%</div> <div>..</div> </div> </div>
1	E	278	<div> <div>3%</div> <div> <div></div> <div>68%</div> <div>24%</div> <div>5%</div> <div>..</div> </div> </div>
2	B	99	<div> <div>3%</div> <div> <div></div> <div>73%</div> <div>21%</div> <div>5%</div> <div>.</div> </div> </div>
2	F	99	<div> <div>3%</div> <div> <div></div> <div>74%</div> <div>21%</div> <div>..</div> </div> </div>
3	C	204	<div> <div>3%</div> <div> <div></div> <div>81%</div> <div>16%</div> <div>.</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	204	
4	D	244	
4	H	244	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	A	1000	X	-	-	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 13298 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-cell surface glycoprotein CD1d.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2192	1400	383	402	7			
1	E	272	Total	C	N	O	S	0	0	0
			2171	1388	377	399	7			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	278	HIS	-	EXPRESSION TAG	UNP P15813
A	279	HIS	-	EXPRESSION TAG	UNP P15813
A	280	HIS	-	EXPRESSION TAG	UNP P15813
A	281	HIS	-	EXPRESSION TAG	UNP P15813
A	282	HIS	-	EXPRESSION TAG	UNP P15813
A	283	HIS	-	EXPRESSION TAG	UNP P15813
E	278	HIS	-	EXPRESSION TAG	UNP P15813
E	279	HIS	-	EXPRESSION TAG	UNP P15813
E	280	HIS	-	EXPRESSION TAG	UNP P15813
E	281	HIS	-	EXPRESSION TAG	UNP P15813
E	282	HIS	-	EXPRESSION TAG	UNP P15813
E	283	HIS	-	EXPRESSION TAG	UNP P15813

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			824	525	139	157	3			
2	F	98	Total	C	N	O	S	0	0	0
			813	517	138	156	2			


- Molecule 3 is a protein called NKT15 alpha-chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	203	Total	C	N	O	S	0	0	0
			1581	979	267	326	9			
3	G	204	Total	C	N	O	S	0	0	0
			1588	984	268	327	9			

- Molecule 4 is a protein called NKT15 beta-chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	244	Total	C	N	O	S	0	0	0
			1957	1230	340	379	8			
4	H	244	Total	C	N	O	S	0	0	0
			1957	1230	340	379	8			

- Molecule 5 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	E	1	Total	C	N	O	0	0
			14	8	1	5		
5	E	1	Total	C	N	O	0	0
			14	8	1	5		
5	E	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 7 is N-{(1S,2R,3S)-1-[(ALPHA-D-GALACTOPYRANOSYLOXY)METHYL]-2,3-DIHYDROXYHEPTADECYL}HEXACOSANAMIDE (three-letter code: AGH) (formula: $C_{50}H_{99}NO_9$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	C	1	Total	C	N	O	0	0
			60	50	1	9		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	E	1	Total	C	N	O	0	0
			60	50	1	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: T-cell surface glycoprotein CD1d



- Molecule 1: T-cell surface glycoprotein CD1d



- Molecule 2: Beta-2-microglobulin





- Molecule 2: Beta-2-microglobulin









• Molecule 3: NKT15 alpha-chain





• Molecule 3: NKT15 alpha-chain



• Molecule 4: NKT15 beta-chain



• Molecule 4: NKT15 beta-chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	204.01Å 155.64Å 85.98Å 90.00° 94.78° 90.00°	Depositor
Resolution (Å)	29.18 – 3.20 29.19 – 3.20	Depositor EDS
% Data completeness (in resolution range)	95.6 (29.18-3.20) 95.6 (29.19-3.20)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 3.18Å)	Xtriage
Refinement program	REFMAC refmac_5.2.0019	Depositor
R, R_{free}	0.226 , 0.293 0.217 , 0.279	Depositor DCC
R_{free} test set	2135 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	57.6	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 10.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	13298	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, AGH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	2/2257 (0.1%)	0.57	0/3076
1	E	0.37	0/2236	0.60	1/3048 (0.0%)
2	B	0.37	0/847	0.60	1/1148 (0.1%)
2	F	0.35	0/836	0.55	1/1134 (0.1%)
3	C	0.39	0/1612	0.56	1/2189 (0.0%)
3	G	0.38	0/1620	0.55	2/2201 (0.1%)
4	D	0.38	0/2011	0.54	0/2736
4	H	0.38	0/2011	0.58	0/2736
All	All	0.39	2/13430 (0.0%)	0.57	6/18268 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	E	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	180	GLU	CD-OE1	9.77	1.36	1.25
1	A	180	GLU	CD-OE2	9.34	1.35	1.25

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	54	LEU	CA-CB-CG	6.29	129.77	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	54	LEU	CA-CB-CG	6.01	129.13	115.30
3	G	166	LEU	CA-CB-CG	5.51	127.98	115.30
1	E	154	THR	C-N-CA	5.32	134.99	121.70
3	G	32	LEU	CA-CB-CG	5.15	127.14	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	91	SER	Peptide
1	E	91	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2192	0	2102	42	0
1	E	2171	0	2086	57	0
2	B	824	0	783	21	0
2	F	813	0	762	11	0
3	C	1581	0	1508	22	0
3	G	1588	0	1515	23	0
4	D	1957	0	1855	27	0
4	H	1957	0	1855	36	0
5	A	42	0	38	0	0
5	E	42	0	38	0	0
6	A	11	0	10	0	0
7	C	60	0	99	6	0
7	E	60	0	99	2	0
All	All	13298	0	12750	217	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 217 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:27:VAL:CA	2:B:28:SER:HB2	1.83	1.06
1:E:7:LEU:HB3	1:E:8:PHE:HA	1.36	1.06
2:B:27:VAL:HA	2:B:28:SER:CB	1.94	0.98
2:B:27:VAL:HA	2:B:28:SER:HB2	0.99	0.98
1:E:154:THR:HG22	1:E:155:ARG:HB2	1.51	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/278 (98%)	244 (89%)	24 (9%)	5 (2%)	10	47
1	E	270/278 (97%)	241 (89%)	16 (6%)	13 (5%)	2	20
2	B	97/99 (98%)	85 (88%)	10 (10%)	2 (2%)	8	42
2	F	96/99 (97%)	87 (91%)	8 (8%)	1 (1%)	18	61
3	C	201/204 (98%)	187 (93%)	12 (6%)	2 (1%)	18	61
3	G	202/204 (99%)	184 (91%)	16 (8%)	2 (1%)	18	61
4	D	242/244 (99%)	218 (90%)	18 (7%)	6 (2%)	6	38
4	H	242/244 (99%)	217 (90%)	16 (7%)	9 (4%)	4	26
All	All	1623/1650 (98%)	1463 (90%)	120 (7%)	40 (2%)	6	38

5 of 40 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	7	ILE
2	B	28	SER
3	C	156	SER
3	C	194	ASN
4	D	52	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/244 (97%)	211 (89%)	25 (11%)	8	32
1	E	235/244 (96%)	217 (92%)	18 (8%)	15	50
2	B	93/94 (99%)	81 (87%)	12 (13%)	5	22
2	F	91/94 (97%)	78 (86%)	13 (14%)	4	18
3	C	185/186 (100%)	169 (91%)	16 (9%)	12	43
3	G	186/186 (100%)	170 (91%)	16 (9%)	12	43
4	D	215/215 (100%)	198 (92%)	17 (8%)	14	49
4	H	215/215 (100%)	199 (93%)	16 (7%)	16	52
All	All	1456/1478 (98%)	1323 (91%)	133 (9%)	11	39

5 of 133 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	78	THR
1	E	92	TYR
4	H	98	ARG
4	D	98	ARG
4	D	190	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 37 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	86	HIS
1	E	55	GLN
4	H	183	GLN
4	D	106	GLN
4	D	236	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	A	1000	1	14,14,15	0.53	0	15,19,21	1.31	2 (13%)
5	NAG	A	1100	6	14,14,15	0.71	0	15,19,21	1.26	1 (6%)
6	BMA	A	1200	5	11,11,12	0.63	0	13,15,17	0.81	1 (7%)
5	NAG	A	2000	1	14,14,15	0.45	0	15,19,21	1.37	1 (6%)
7	AGH	C	208	-	60,60,60	0.40	0	64,69,69	0.76	0
7	AGH	E	2	-	60,60,60	0.51	1 (1%)	64,69,69	0.88	2 (3%)
5	NAG	E	3000	1,5	14,14,15	0.50	0	15,19,21	0.82	0
5	NAG	E	3100	5	14,14,15	0.59	0	15,19,21	1.17	2 (13%)
5	NAG	E	4000	1	14,14,15	0.44	0	15,19,21	1.38	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	1000	1	1/1/5/7	0/6/23/26	0/1/1/1
5	NAG	A	1100	6	-	0/6/23/26	0/1/1/1
6	BMA	A	1200	5	-	0/2/19/22	0/1/1/1
5	NAG	A	2000	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	AGH	C	208	-	-	0/58/78/78	0/1/1/1
7	AGH	E	2	-	-	0/58/78/78	0/1/1/1
5	NAG	E	3000	1,5	-	0/6/23/26	0/1/1/1
5	NAG	E	3100	5	-	0/6/23/26	0/1/1/1
5	NAG	E	4000	1	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	E	2	AGH	O1A-C1A	2.63	1.44	1.40

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	E	2	AGH	C1-C2-N2	-3.02	105.58	109.76
5	A	1000	NAG	O5-C1-C2	-2.40	108.14	111.47
7	E	2	AGH	C1A-O6A-C5M	2.09	117.65	113.72
6	A	1200	BMA	C1-C2-C3	2.27	112.53	109.65
5	E	3100	NAG	C3-C4-C5	2.51	114.64	110.22

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	1000	NAG	C1

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	E	4000	NAG	O7-C7-N2-C2

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	208	AGH	6	0
7	E	2	AGH	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/278 (98%)	0.20	11 (4%) 39 25	52, 57, 61, 63	0
1	E	272/278 (97%)	0.15	8 (2%) 52 37	53, 58, 62, 65	0
2	B	99/99 (100%)	0.41	3 (3%) 51 35	53, 58, 60, 64	0
2	F	98/99 (98%)	0.20	3 (3%) 49 33	54, 58, 61, 62	0
3	C	203/204 (99%)	0.11	6 (2%) 51 35	52, 58, 61, 65	0
3	G	204/204 (100%)	0.11	4 (1%) 65 50	51, 58, 62, 64	0
4	D	244/244 (100%)	0.03	6 (2%) 58 43	54, 57, 62, 64	0
4	H	244/244 (100%)	-0.01	3 (1%) 79 67	49, 57, 63, 66	0
All	All	1639/1650 (99%)	0.13	44 (2%) 55 40	49, 58, 62, 66	0

The worst 5 of 44 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	135	ASP	5.1
2	B	1	ILE	4.1
1	A	108	ASN	3.7
1	E	271	GLN	3.6
3	C	135	ASP	3.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	AGH	E	2	60/60	0.95	0.20	-0.70	30,35,47,48	0
7	AGH	C	208	60/60	0.95	0.19	-0.92	29,37,55,55	0
5	NAG	A	1100	14/15	0.48	0.45	-	154,155,156,156	0
5	NAG	A	2000	14/15	0.87	0.30	-	66,68,69,69	0
5	NAG	E	3100	14/15	0.78	0.38	-	82,83,83,83	0
5	NAG	A	1000	14/15	0.92	0.22	-	62,63,64,64	0
5	NAG	E	3000	14/15	0.83	0.30	-	73,76,78,80	0
6	BMA	A	1200	11/12	0.42	0.41	-	156,156,156,156	0
5	NAG	E	4000	14/15	0.85	0.27	-	65,65,67,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.