



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2017 – 11:00 PM EDT

PDB ID : 3PO4
Title : Structure of a mutant of the large fragment of DNA polymerase I from *Thermus aquaticus* in complex with a blunt-ended DNA and ddATP
Authors : Marx, A.; Diederichs, K.; Obeid, S.
Deposited on : unknown
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	<div> <div>2%</div> <div>87%</div> <div>12%</div> </div>
2	B	12	<div> <div>42%</div> <div>58%</div> </div>
3	C	12	<div> <div>67%</div> <div>17%</div> <div>17%</div> </div>

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5385 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	538	Total	C	N	O	S	0	4	0
			4313	2736	781	784	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	614	LYS	ILE	ENGINEERED MUTATION	UNP P19821
A	747	LYS	MET	ENGINEERED MUTATION	UNP P19821

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*GP*CP*(2DA))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	P	0	0	0
			242	115	50	66	11			

- Molecule 3 is a DNA chain called DNA (5'-D(*TP*GP*CP*GP*CP*CP*GP*TP*GP*GP*TP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	12	Total	C	N	O	P	0	0	0
			243	116	43	73	11			


- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Mg	0	0
			2	2		
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is 2',3'-dideoxyadenosine triphosphate (three-letter code: DDS) (formula: $C_{10}H_{16}N_5O_{11}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	P	0	0
			29	10	5	11	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	438	Total	O	0	0
			438	438		
8	B	44	Total	O	0	0
			44	44		
8	C	66	Total	O	0	0
			66	66		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase I



- Molecule 2: DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*GP*CP*(2DA))-3')



- Molecule 3: DNA (5'-D(*TP*GP*CP*GP*CP*CP*GP*TP*GP*GP*TP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	113.53 Å 113.53 Å 91.24 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.20 – 1.80 48.20 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.20-1.80) 100.0 (48.20-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 1.79 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_572)	Depositor
R, R_{free}	0.164 , 0.194 0.160 , 0.190	Depositor DCC
R_{free} test set	3136 reflections (4.97%)	DCC
Wilson B-factor (Å ²)	33.1	Xtriage
Anisotropy	0.184	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 38.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.032 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5385	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DDS, GOL, MG, 2DA, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/4406	0.48	0/5964
2	B	0.73	0/249	1.40	6/382 (1.6%)
3	C	1.25	3/271 (1.1%)	1.80	10/417 (2.4%)
All	All	0.45	3/4926 (0.1%)	0.72	16/6763 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	205	DT	C5-C6	11.66	1.42	1.34
3	C	205	DT	N1-C2	6.43	1.43	1.38
3	C	205	DT	O4'-C1'	5.69	1.49	1.42

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	205	DT	C2-N3-C4	-10.54	120.87	127.20
3	C	205	DT	N3-C4-C5	8.62	120.37	115.20
2	B	106	DC	O4'-C1'-N1	-8.02	102.39	108.00
3	C	205	DT	C5-C6-N1	-8.02	118.89	123.70
3	C	205	DT	N1-C1'-C2'	7.63	127.09	112.60
2	B	107	DG	O4'-C1'-N9	-6.41	103.51	108.00
2	B	108	DG	C1'-O4'-C4'	-6.30	103.80	110.10
3	C	205	DT	N1-C2-N3	6.25	118.35	114.60
3	C	205	DT	N1-C2-O2	-6.20	118.14	123.10
2	B	109	DC	O4'-C1'-N1	5.96	112.17	108.00
2	B	108	DG	O4'-C1'-N9	5.90	112.13	108.00
3	C	208	DG	O4'-C1'-N9	-5.85	103.90	108.00
3	C	205	DT	C6-C5-C7	-5.83	119.41	122.90
3	C	210	DC	O4'-C1'-N1	-5.10	104.43	108.00
3	C	205	DT	C1'-O4'-C4'	-5.06	105.04	110.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	105	DA	O4'-C1'-N9	-5.04	104.47	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4313	0	4363	50	0
2	B	242	0	134	2	0
3	C	243	0	137	2	0
4	A	1	0	0	0	0
4	B	2	0	0	0	0
5	A	1	0	0	0	0
6	A	6	0	8	1	0
7	A	29	0	12	2	0
8	A	438	0	0	6	0
8	B	44	0	0	0	0
8	C	66	0	0	0	0
All	All	5385	0	4654	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:644:SER:O	1:A:649:VAL:HG12	1.73	0.86
1:A:557:THR:OG1	1:A:561:HIS:HE1	1.79	0.65
1:A:649:VAL:HB	1:A:650:PRO:HA	1.79	0.63
1:A:665:ILE:CD1	1:A:678:LEU:HD21	2.29	0.63
1:A:649:VAL:HG23	1:A:650:PRO:CA	2.29	0.62
1:A:649:VAL:HG23	1:A:650:PRO:N	2.12	0.62
1:A:771:ARG:O	1:A:774:GLU:HG2	2.00	0.61
1:A:557:THR:HG1	1:A:561:HIS:HE1	1.50	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:665:ILE:HD13	1:A:678:LEU:HD11	1.85	0.59
1:A:786:GLU:HG3	8:A:1014:HOH:O	2.03	0.58
1:A:822:GLY:HA3	1:A:830:ALA:O	2.02	0.58
1:A:649:VAL:CB	1:A:650:PRO:HA	2.34	0.58
1:A:592:GLN:HG2	8:A:1026:HOH:O	2.04	0.56
7:A:835:DDS:O5'	7:A:835:DDS:H8	2.07	0.54
1:A:547:ASP:HB2	1:A:548:PRO:HD3	1.89	0.54
1:A:642:THR:O	1:A:646:MET:HG2	2.08	0.53
3:C:210:DC:H2'	3:C:211:DG:C8	2.43	0.53
1:A:655:ASP:HB2	1:A:656:PRO:HD2	1.90	0.53
2:B:103:DC:H2''	2:B:104:DC:H5'	1.90	0.53
1:A:731:PRO:O	1:A:732:ASP:HB2	2.08	0.52
1:A:648:GLY:O	1:A:649:VAL:HG22	2.10	0.52
1:A:445:GLU:O	1:A:561:HIS:HD2	1.93	0.51
1:A:671:TYR:HE2	7:A:835:DDS:H2	1.74	0.51
1:A:786:GLU:CG	8:A:1014:HOH:O	2.57	0.50
1:A:556:ARG:NE	1:A:556:ARG:HA	2.26	0.50
1:A:299:TRP:CG	1:A:300:PRO:HA	2.46	0.50
1:A:818:GLU:HG3	8:A:1030:HOH:O	2.11	0.50
1:A:769:PHE:HB3	1:A:770:PRO:HD3	1.93	0.49
2:B:103:DC:H2''	2:B:104:DC:C5'	2.43	0.49
1:A:526:HIS:ND1	1:A:527:PRO:HD2	2.28	0.48
1:A:608:ALA:HB1	8:A:1014:HOH:O	2.13	0.47
1:A:473:GLU:HG2	1:A:476[A]:ARG:HH22	1.80	0.47
1:A:388:GLU:HG3	1:A:398:TRP:CD1	2.52	0.45
1:A:469:ARG:O	1:A:473:GLU:HG3	2.16	0.45
1:A:469:ARG:NH2	8:A:860:HOH:O	2.50	0.45
1:A:687:GLU:H	1:A:687:GLU:CD	2.21	0.43
1:A:490:LEU:HD11	1:A:532:ILE:HD13	2.00	0.43
1:A:655:ASP:HB2	1:A:656:PRO:CD	2.49	0.43
1:A:576:SER:O	3:C:208:DG:H4'	2.18	0.43
1:A:648:GLY:C	1:A:649:VAL:HG13	2.39	0.43
1:A:428:TRP:CD2	6:A:834:GOL:H32	2.54	0.43
1:A:563:ARG:O	1:A:576:SER:HA	2.18	0.43
1:A:649:VAL:CG2	1:A:650:PRO:N	2.81	0.43
1:A:664:THR:HG21	1:A:682:LEU:HD13	2.02	0.41
1:A:554:HIS:HA	1:A:555:PRO:HD3	1.93	0.41
1:A:650:PRO:HD2	1:A:653:ALA:CB	2.49	0.41
1:A:673:MET:HE2	1:A:673:MET:HB2	1.90	0.41
1:A:631:VAL:HG11	1:A:638:ILE:HA	2.03	0.41
1:A:665:ILE:HD13	1:A:678:LEU:HD21	2.02	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:596:ARG:HG2	1:A:828:LEU:HD23	2.02	0.40
1:A:453:VAL:O	1:A:457:ARG:HG2	2.21	0.40
1:A:596:ARG:CG	1:A:828:LEU:HD23	2.50	0.40
1:A:427:LEU:HA	1:A:427:LEU:HD23	1.96	0.40
1:A:605:LEU:HG	1:A:793:LYS:HG2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	540/540 (100%)	523 (97%)	13 (2%)	4 (1%)	25 11

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	649	VAL
1	A	831	LYS
1	A	586	VAL
1	A	503	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	444/441 (101%)	436 (98%)	8 (2%)	64 53

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	487	ARG
1	A	492	ARG
1	A	538	LEU
1	A	649	VAL
1	A	673	MET
1	A	682	LEU
1	A	695	ARG
1	A	794	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	333	HIS
1	A	561	HIS
1	A	750	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	2DA	B	112	3,2,4	16,22,23	2.07	6 (37%)	13,31,34	3.81	3 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2DA	B	112	3,2,4	-	0/3/18/19	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	112	2DA	O5'-C5'	-3.93	1.39	1.44
2	B	112	2DA	C2'-C3'	-2.21	1.47	1.54
2	B	112	2DA	C6-N6	2.01	1.42	1.34
2	B	112	2DA	O4'-C1'	2.46	1.48	1.42
2	B	112	2DA	C2-N1	3.63	1.40	1.33
2	B	112	2DA	C4-N3	3.91	1.41	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	112	2DA	N3-C2-N1	-10.88	119.39	128.86
2	B	112	2DA	O4'-C1'-C2'	-2.84	103.59	106.67
2	B	112	2DA	C2'-C1'-N9	6.99	125.77	112.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
6	GOL	A	834	-	5,5,5	0.34	0	5,5,5	0.31	0
7	DDS	A	835	4	25,31,31	1.25	4 (16%)	21,48,48	2.31	4 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	834	-	-	0/4/4/4	0/0/0/0
7	DDS	A	835	4	-	0/18/31/31	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	835	DDS	C5'-C4'	-2.85	1.42	1.50
7	A	835	DDS	C3'-C2'	-2.48	1.47	1.54
7	A	835	DDS	PG-O3B	-2.15	1.56	1.60
7	A	835	DDS	C6-N6	2.87	1.45	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	835	DDS	N3-C2-N1	-9.18	120.86	128.86
7	A	835	DDS	C4-C5-N7	-2.23	107.25	109.41
7	A	835	DDS	C3'-C2'-C1'	2.05	104.97	102.69
7	A	835	DDS	O4'-C1'-N9	2.59	112.15	107.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	834	GOL	1	0
7	A	835	DDS	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	538/540 (99%)	-0.45	9 (1%) 70 67	20, 35, 75, 137	0
2	B	11/12 (91%)	-0.96	0 100 100	24, 26, 50, 51	0
3	C	12/12 (100%)	-0.97	0 100 100	22, 27, 46, 47	0
All	All	561/564 (99%)	-0.47	9 (1%) 72 69	20, 35, 73, 137	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	648	GLY	5.4
1	A	503	ILE	4.5
1	A	649	VAL	4.2
1	A	647	PHE	3.1
1	A	645	TRP	2.8
1	A	657	LEU	2.3
1	A	646	MET	2.2
1	A	644	SER	2.2
1	A	653	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	2DA	B	112	20/21	0.98	0.09	-	21,25,28,29	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	DDS	A	835	29/29	0.94	0.11	1.30	27,45,66,271	0
4	MG	B	113	1/1	0.95	0.12	0.17	34,34,34,34	0
4	MG	A	1	1/1	0.96	0.09	-	36,36,36,36	0
5	NA	A	833	1/1	0.69	0.12	-	50,50,50,50	0
4	MG	B	1	1/1	0.98	0.14	-	36,36,36,36	0
6	GOL	A	834	6/6	0.94	0.11	-	49,60,64,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.