



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 12, 2017 – 10:25 AM EST

PDB ID : 4PQK  
Title : C-Terminal domain of DNA binding protein  
Authors : Schumacher, M.A.; Chinnam, N.; Tonthat, N.K.  
Deposited on : unknown  
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1679 (3.50-3.30)
Clashscore	112137	1832 (3.50-3.30)
Ramachandran outliers	110173	1789 (3.50-3.30)
Sidechain outliers	110143	1789 (3.50-3.30)
RSRZ outliers	101464	1709 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	487	<div> <div>8%</div> <div>64%</div> <div>30%</div> <div>• •</div> </div>
1	B	487	<div> <div>7%</div> <div>70%</div> <div>26%</div> <div>• •</div> </div>
1	C	487	<div> <div>7%</div> <div>68%</div> <div>25%</div> <div>5% •</div> </div>
1	D	487	<div> <div>7%</div> <div>63%</div> <div>28%</div> <div>6% •</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 14876 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Maltose ABC transporter periplasmic protein, Truncated replication protein RepA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	478	Total	C	N	O	S	0	0	0
			3700	2371	603	713	13			
1	B	479	Total	C	N	O	S	0	0	0
			3722	2392	615	703	12			
1	C	476	Total	C	N	O	S	0	0	0
			3648	2346	601	690	11			
1	D	472	Total	C	N	O	S	0	0	0
			3670	2351	601	707	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP U6NJU2
B	1	MET	-	EXPRESSION TAG	UNP U6NJU2
C	1	MET	-	EXPRESSION TAG	UNP U6NJU2
D	1	MET	-	EXPRESSION TAG	UNP U6NJU2

- Molecule 2 is MALTOTRIOSE (three-letter code: MLR) (formula: C<sub>18</sub>H<sub>32</sub>O<sub>16</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total 34	C 18	O 16	0	0
2	B	1	Total 34	C 18	O 16	0	0
2	C	1	Total 34	C 18	O 16	0	0
2	D	1	Total 34	C 18	O 16	0	0


### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose ABC transporter periplasmic protein, Truncated replication protein RepA




- Molecule 1: Maltose ABC transporter periplasmic protein, Truncated replication protein RepA






- Molecule 1: Maltose ABC transporter periplasmic protein, Truncated replication protein RepA



- Molecule 1: Maltose ABC transporter periplasmic protein, Truncated replication protein RepA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.22Å 93.67Å 100.81Å 107.67° 108.30° 84.93°	Depositor
Resolution (Å)	46.62 – 3.40 46.62 – 3.40	Depositor EDS
% Data completeness (in resolution range)	(Not available) (46.62-3.40) 63.2 (46.62-3.40)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.43 (at 3.40Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, $R_{free}$	0.237 , 0.275 0.240 , 0.273	Depositor DCC
$R_{free}$ test set	1963 reflections (10.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	73.9	Xtriage
Anisotropy	0.378	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 25.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	14876	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/3784	0.57	0/5142
1	B	0.44	0/3805	0.54	0/5157
1	C	0.43	0/3730	0.63	1/5063 (0.0%)
1	D	0.45	0/3751	0.65	2/5090 (0.0%)
All	All	0.44	0/15070	0.60	3/20452 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	8	LEU	CA-CB-CG	5.36	127.62	115.30
1	D	458	LEU	CA-CB-CG	5.14	127.12	115.30
1	C	463	LEU	CA-CB-CG	5.02	126.84	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3700	0	3562	116	0
1	B	3722	0	3663	87	0
1	C	3648	0	3540	85	0
1	D	3670	0	3560	108	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	34	0	31	7	0
2	B	34	0	32	7	0
2	C	34	0	31	5	0
2	D	34	0	32	6	0
All	All	14876	0	14451	411	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 411 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:700:MLR:O5C	2:C:700:MLR:C5C	1.63	1.46
2:A:700:MLR:C5C	2:A:700:MLR:O5C	1.64	1.45
2:D:700:MLR:C5C	2:D:700:MLR:O5C	1.64	1.43
2:B:700:MLR:C5C	2:B:700:MLR:O5C	1.64	1.41
2:B:700:MLR:C1C	2:B:700:MLR:O5C	1.72	1.38

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	476/487 (98%)	463 (97%)	13 (3%)	0	100	100
1	B	477/487 (98%)	472 (99%)	5 (1%)	0	100	100
1	C	474/487 (97%)	469 (99%)	5 (1%)	0	100	100
1	D	468/487 (96%)	464 (99%)	4 (1%)	0	100	100
All	All	1895/1948 (97%)	1868 (99%)	27 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	375/407 (92%)	326 (87%)	49 (13%)	5	23
1	B	380/407 (93%)	337 (89%)	43 (11%)	7	29
1	C	363/407 (89%)	308 (85%)	55 (15%)	3	18
1	D	375/407 (92%)	315 (84%)	60 (16%)	3	15
All	All	1493/1628 (92%)	1286 (86%)	207 (14%)	4	21

5 of 207 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	4	GLU
1	C	201	LYS
1	D	434	ARG
1	C	42	ASP
1	C	128	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	324	ASN
1	B	336	GLN
1	B	378	GLN
1	C	369	ASN
1	D	470	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	MLR	A	700	-	36,36,36	3.41	11 (30%)	53,53,53	2.58	18 (33%)
2	MLR	B	700	-	36,36,36	3.31	10 (27%)	53,53,53	1.96	9 (16%)
2	MLR	C	700	-	36,36,36	3.25	13 (36%)	53,53,53	2.24	13 (24%)
2	MLR	D	700	-	36,36,36	3.31	11 (30%)	53,53,53	1.74	8 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLR	A	700	-	-	0/14/74/74	0/3/3/3
2	MLR	B	700	-	-	0/14/74/74	0/3/3/3
2	MLR	C	700	-	-	0/14/74/74	0/3/3/3
2	MLR	D	700	-	-	0/14/74/74	0/3/3/3

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	700	MLR	C4C-C3C	-5.06	1.39	1.52
2	C	700	MLR	C4C-C3C	-4.38	1.41	1.52
2	D	700	MLR	C4C-C3C	-4.09	1.41	1.52
2	B	700	MLR	C4C-C3C	-4.08	1.42	1.52
2	A	700	MLR	C3C-C2C	-4.07	1.42	1.52

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	700	MLR	C1C-O5C-C5C	-9.05	96.66	113.72
2	C	700	MLR	C1C-O5C-C5C	-7.85	98.92	113.72
2	B	700	MLR	C1C-O5C-C5C	-7.71	99.20	113.72
2	D	700	MLR	C1C-O5C-C5C	-6.27	101.90	113.72
2	C	700	MLR	O3C-C3C-C2C	-4.66	100.21	110.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	700	MLR	7	0
2	B	700	MLR	7	0
2	C	700	MLR	5	0
2	D	700	MLR	6	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	478/487 (98%)	0.31	38 (7%) 13 13	31, 61, 95, 121	0
1	B	479/487 (98%)	0.34	32 (6%) 19 19	39, 66, 99, 135	0
1	C	476/487 (97%)	0.26	34 (7%) 17 17	41, 68, 105, 140	0
1	D	472/487 (96%)	0.31	34 (7%) 16 17	45, 73, 100, 138	0
All	All	1905/1948 (97%)	0.31	138 (7%) 16 17	31, 67, 101, 140	0

The worst 5 of 138 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	372	ASP	5.9
1	C	380	PHE	5.9
1	C	173	GLU	5.9
1	C	375	MET	5.8
1	B	386	ASN	5.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MLR	D	700	34/34	0.83	0.28	0.25	45,60,73,74	0
2	MLR	B	700	34/34	0.89	0.24	0.14	33,53,65,76	0
2	MLR	A	700	34/34	0.87	0.24	-0.05	30,42,62,76	0
2	MLR	C	700	34/34	0.95	0.20	-0.43	53,65,78,81	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.