



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 16, 2018 – 03:07 PM EST

PDB ID : 3PSB  
Title : Furo[2,3-c]pyridine-based Indanone Oximes as Potent and Selective B-Raf Inhibitors  
Authors : Morales, T.; Vigers, G.P.A.; Brandhuber, B.J.  
Deposited on : 2010-12-01  
Resolution : 3.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030736  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1679 (3.50-3.30)
Clashscore	112137	1832 (3.50-3.30)
Ramachandran outliers	110173	1789 (3.50-3.30)
Sidechain outliers	110143	1789 (3.50-3.30)
RSRZ outliers	101464	1709 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	<div> <div>3%</div> <div> <div></div> <div>72%</div> <div>11%</div> <div>•</div> <div>15%</div> </div> </div>
1	B	307	<div> <div>3%</div> <div> <div></div> <div>70%</div> <div>13%</div> <div>•</div> <div>14%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called B-RAF PROTO-ONCOGENE SERINE/THREONINE-PROTEIN KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	262	Total	C	N	O	S	0	0	0
			2093	1343	364	373	13			
1	B	264	Total	C	N	O	S	0	0	0
			2107	1352	366	376	13			

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	420	MET	-	EXPRESSION TAG	UNP P15056
A	421	ASP	-	EXPRESSION TAG	UNP P15056
A	422	ARG	-	EXPRESSION TAG	UNP P15056
A	423	GLY	-	EXPRESSION TAG	UNP P15056
A	424	SER	-	EXPRESSION TAG	UNP P15056
A	425	HIS	-	EXPRESSION TAG	UNP P15056
A	426	HIS	-	EXPRESSION TAG	UNP P15056
A	427	HIS	-	EXPRESSION TAG	UNP P15056
A	428	HIS	-	EXPRESSION TAG	UNP P15056
A	429	HIS	-	EXPRESSION TAG	UNP P15056
A	430	HIS	-	EXPRESSION TAG	UNP P15056
A	431	GLY	-	EXPRESSION TAG	UNP P15056
A	432	SER	-	EXPRESSION TAG	UNP P15056
B	420	MET	-	EXPRESSION TAG	UNP P15056
B	421	ASP	-	EXPRESSION TAG	UNP P15056
B	422	ARG	-	EXPRESSION TAG	UNP P15056
B	423	GLY	-	EXPRESSION TAG	UNP P15056
B	424	SER	-	EXPRESSION TAG	UNP P15056
B	425	HIS	-	EXPRESSION TAG	UNP P15056
B	426	HIS	-	EXPRESSION TAG	UNP P15056
B	427	HIS	-	EXPRESSION TAG	UNP P15056
B	428	HIS	-	EXPRESSION TAG	UNP P15056
B	429	HIS	-	EXPRESSION TAG	UNP P15056
B	430	HIS	-	EXPRESSION TAG	UNP P15056

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	431	GLY	-	EXPRESSION TAG	UNP P15056
B	432	SER	-	EXPRESSION TAG	UNP P15056

- Molecule 2 is ethyl 3-{{1-(hydroxyamino)-2H-inden-5-yl}amino}thieno[2,3-c]pyridine-2-carboxylate (three-letter code: SM6) (formula: C<sub>19</sub>H<sub>17</sub>N<sub>3</sub>O<sub>3</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			26	19	3	3	1		
2	B	1	Total	C	N	O	S	0	0
			26	19	3	3	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: B-RAF PROTO-ONCOGENE SERINE/THREONINE-PROTEIN KINASE



- Molecule 1: B-RAF PROTO-ONCOGENE SERINE/THREONINE-PROTEIN KINASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.07Å 101.07Å 162.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.63 – 3.40 28.63 – 3.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.63-3.40) 99.7 (28.63-3.40)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	0.17	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.63 (at 3.39Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.224 , 0.268 0.210 , 0.253	Depositor DCC
$R_{free}$ test set	621 reflections (5.40%)	DCC
Wilson B-factor (Å <sup>2</sup> )	64.9	Xtriage
Anisotropy	0.196	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 18.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	4252	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SM6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/2138	0.55	0/2886
1	B	0.36	0/2152	0.52	0/2905
All	All	0.36	0/4290	0.54	0/5791

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2093	0	2135	17	0
1	B	2107	0	2151	25	0
2	A	26	0	17	1	0
2	B	26	0	17	3	0
All	All	4252	0	4320	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:658:ASN:H	1:B:658:ASN:HD22	1.22	0.88
2:B:2:SM6:H19	2:B:2:SM6:C3	2.11	0.80
2:B:2:SM6:H19	2:B:2:SM6:H3	1.65	0.77
1:A:551:ILE:HD12	1:A:551:ILE:H	1.55	0.71
1:B:573:ILE:HG22	1:B:575:ARG:HG3	1.72	0.70
2:B:2:SM6:H3	2:B:2:SM6:C19	2.23	0.69
1:B:551:ILE:HD12	1:B:551:ILE:H	1.61	0.66
1:A:511:VAL:O	1:A:591:LYS:HE2	1.97	0.65
1:A:573:ILE:HG22	1:A:575:ARG:HG3	1.79	0.62
1:B:684:ASN:H	1:B:684:ASN:HD22	1.47	0.62
1:B:511:VAL:O	1:B:591:LYS:HE2	1.99	0.62
1:B:721:LEU:HB3	1:B:722:PRO:HD3	1.83	0.60
1:B:575:ARG:NH1	1:B:597:LEU:O	2.35	0.58
1:B:660:ASN:C	1:B:660:ASN:HD22	2.08	0.56
1:B:578:LYS:HG3	1:B:580:ASN:HB2	1.88	0.55
1:A:484:MET:HG3	1:A:526:ALA:HB2	1.89	0.55
1:B:617:ILE:HD13	1:B:666:ILE:HG13	1.91	0.53
1:B:658:ASN:H	1:B:658:ASN:ND2	2.00	0.53
1:A:684:ASN:HD22	1:A:684:ASN:H	1.57	0.52
1:B:668:MET:HG2	1:B:673:TYR:HB2	1.92	0.52
1:B:658:ASN:N	1:B:658:ASN:HD22	2.01	0.52
1:B:684:ASN:N	1:B:684:ASN:HD22	2.07	0.51
1:A:537:LEU:HD21	1:A:649:LEU:HD21	1.94	0.50
1:B:707:PHE:HA	1:B:710:ILE:HB	1.94	0.50
1:A:642:PHE:O	1:A:645:VAL:HG22	2.13	0.49
1:B:491:THR:HB	1:B:494:GLN:H	1.76	0.48
1:B:555:ASP:OD1	1:B:558:ARG:NH1	2.46	0.48
1:A:454:ASP:H	1:A:522:LYS:HD3	1.78	0.48
1:A:658:ASN:HD22	1:A:658:ASN:H	1.61	0.48
1:A:486:ASN:HA	1:A:524:GLN:CD	2.35	0.47
1:A:482:VAL:HG22	1:A:528:VAL:HG22	1.95	0.47
1:A:692:LEU:HD11	1:A:710:ILE:HG23	1.97	0.47
1:A:623:GLU:HG3	1:A:623:GLU:H	1.39	0.46
1:A:500:ASN:O	1:A:504:VAL:HG23	2.16	0.46
1:B:651:THR:HG22	1:B:681:VAL:HA	1.98	0.45
1:B:510:HIS:HB3	1:B:513:ILE:HG12	1.98	0.45
1:A:664:GLN:O	1:A:668:MET:HB2	2.17	0.44
1:B:576:ASP:HB2	1:B:597:LEU:HD12	2.00	0.43
1:B:575:ARG:HG2	1:B:633:TYR:CD2	2.55	0.42
1:A:541:LEU:HD23	1:A:648:GLU:HB3	2.02	0.42
1:B:622:PRO:HA	1:B:625:ILE:HG22	2.01	0.42
1:A:684:ASN:N	1:A:684:ASN:HD22	2.16	0.41

Continued on next page...



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1:SM6:C3	2:A:1:SM6:H19	2.50	0.41
1:B:618:LEU:HD23	1:B:654:LEU:HD22	2.01	0.41
1:B:599:THR:HG22	1:B:600:VAL:N	2.35	0.41
1:B:668:MET:HB3	1:B:674:LEU:HB2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/307 (84%)	238 (92%)	17 (7%)	3 (1%)	15	54
1	B	260/307 (85%)	245 (94%)	10 (4%)	5 (2%)	9	45
All	All	518/614 (84%)	483 (93%)	27 (5%)	8 (2%)	12	49

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	627	MET
1	A	629	ASP
1	B	523	PRO
1	B	594	ASP
1	B	629	ASP
1	B	535	SER
1	B	721	LEU
1	A	721	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/271 (85%)	216 (94%)	14 (6%)	22	59
1	B	232/271 (86%)	214 (92%)	18 (8%)	15	49
All	All	462/542 (85%)	430 (93%)	32 (7%)	18	55

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	491	THR
1	A	495	LEU
1	A	541	LEU
1	A	547	LYS
1	A	577	LEU
1	A	588	LEU
1	A	623	GLU
1	A	625	ILE
1	A	628	GLN
1	A	658	ASN
1	A	660	ASN
1	A	668	MET
1	A	674	LEU
1	A	684	ASN
1	B	461	GLN
1	B	491	THR
1	B	535	SER
1	B	541	LEU
1	B	547	LYS
1	B	551	ILE
1	B	575	ARG
1	B	577	LEU
1	B	588	LEU
1	B	623	GLU
1	B	628	GLN
1	B	631	ASN
1	B	656	TYR
1	B	658	ASN
1	B	660	ASN
1	B	668	MET
1	B	674	LEU
1	B	684	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	585	HIS
1	A	628	GLN
1	A	631	ASN
1	A	658	ASN
1	A	660	ASN
1	A	684	ASN
1	B	461	GLN
1	B	562	GLN
1	B	631	ASN
1	B	658	ASN
1	B	660	ASN
1	B	684	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SM6	A	1	-	20,29,29	1.73	5 (25%)	19,41,41	3.69	7 (36%)
2	SM6	B	2	-	20,29,29	1.78	5 (25%)	19,41,41	3.45	5 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SM6	A	1	-	-	0/9/22/22	0/4/4/4
2	SM6	B	2	-	-	0/9/22/22	0/4/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	SM6	C4-N10	-3.08	1.34	1.40
2	B	2	SM6	C4-N10	-2.39	1.35	1.40
2	A	1	SM6	C8-C7	2.35	1.54	1.51
2	A	1	SM6	O23-N12	2.35	1.44	1.39
2	B	2	SM6	C8-C7	2.49	1.55	1.51
2	A	1	SM6	C11-C13	2.54	1.46	1.41
2	B	2	SM6	O23-N12	2.66	1.44	1.39
2	B	2	SM6	C11-C13	2.83	1.47	1.41
2	A	1	SM6	C19-C20	4.02	1.42	1.36
2	B	2	SM6	C19-C20	4.05	1.42	1.36

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	SM6	C19-C13-C11	-11.84	128.60	135.54
2	B	2	SM6	C19-C13-C11	-11.66	128.70	135.54
2	B	2	SM6	O23-N12-C7	-5.17	110.18	120.05
2	A	1	SM6	O23-N12-C7	-4.97	110.56	120.05
2	A	1	SM6	C22-C14-C13	-3.01	120.43	122.82
2	B	2	SM6	C22-C14-C13	-2.63	120.73	122.82
2	A	1	SM6	C20-C19-C13	-2.07	117.90	119.74
2	A	1	SM6	C26-O24-C17	2.50	121.95	116.45
2	B	2	SM6	C8-C7-N12	4.04	126.67	122.38
2	A	1	SM6	C11-C13-C14	5.03	113.18	107.49
2	B	2	SM6	C11-C13-C14	5.53	113.74	107.49
2	A	1	SM6	C8-C7-N12	6.38	129.15	122.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	SM6	1	0
2	B	2	SM6	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	262/307 (85%)	-0.07	9 (3%) 46 41	29, 42, 59, 78	0
1	B	264/307 (85%)	0.09	10 (3%) 41 37	32, 48, 68, 78	0
All	All	526/614 (85%)	0.01	19 (3%) 43 39	29, 44, 64, 78	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	723	LYS	5.1
1	B	722	PRO	4.0
1	A	723	LYS	3.9
1	A	722	PRO	3.5
1	B	630	LYS	3.3
1	B	587	ASP	2.7
1	A	492	PRO	2.6
1	A	448	ASP	2.6
1	B	448	ASP	2.5
1	A	630	LYS	2.5
1	A	627	MET	2.4
1	A	629	ASP	2.4
1	B	614	SER	2.3
1	B	721	LEU	2.2
1	A	486	ASN	2.2
1	B	454	ASP	2.1
1	B	486	ASN	2.1
1	A	628	GLN	2.0
1	B	629	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SM6	B	2	26/26	0.87	0.26	1.06	60,61,63,65	0
2	SM6	A	1	26/26	0.92	0.22	0.80	51,52,54,55	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.