



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 12, 2017 – 07:09 AM EST

PDB ID : 4PSR
Title : Crystal Structure of alpha-L-fucosidase from Fusarium graminearum in the open form in complex with L-fucose
Authors : Cao, H.; Walton, J.; Brumm, P.; Phillips Jr., G.N.
Deposited on : unknown
Resolution : 1.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2133 (1.40-1.36)
Clashscore	112137	2266 (1.40-1.36)
Ramachandran outliers	110173	2215 (1.40-1.36)
Sidechain outliers	110143	2214 (1.40-1.36)
RSRZ outliers	101464	2141 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	585	<div> <div>2%</div> <div> <div></div> <div>93%</div> <div>6%</div> </div> </div>
1	B	585	<div> <div>2%</div> <div> <div></div> <div>93%</div> <div>6%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MAN	A	606	-	-	-	X
4	MAN	A	607	-	-	-	X
4	MAN	A	610[A]	-	-	-	X
4	MAN	A	610[B]	-	-	-	X
4	MAN	A	611	-	-	-	X
4	MAN	A	612	-	-	-	X
5	TRS	A	616	-	-	-	X
5	TRS	A	618	-	-	-	X

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 21517 atoms, of which 9640 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-fucosidase GH29.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	581	Total	C	H	N	O	S	0	55	0
			9531	3091	4646	833	951	10			
1	B	583	Total	C	H	N	O	S	0	62	0
			9601	3110	4679	837	965	10			

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O		0	0
			27	8	13	1	5			
2	A	1	Total	C	H	N	O		0	0
			27	8	13	1	5			
2	A	1	Total	C	H	N	O		0	0
			28	8	14	1	5			
2	A	1	Total	C	H	N	O		0	0
			28	8	14	1	5			

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
2	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
2	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		

- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			20	6	9	5		
3	A	1	Total	C	H	O	0	0
			20	6	9	5		

- Molecule 4 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			21	6	10	5		
4	A	1	Total	C	H	O	0	0
			20	6	9	5		
4	A	1	Total	C	H	O	0	0
			20	6	9	5		
4	A	1	Total	C	H	O	0	0
			21	6	10	5		
4	A	1	Total	C	H	O	0	1
			40	12	18	10		
4	A	1	Total	C	H	O	0	0
			21	6	10	5		
4	A	1	Total	C	H	O	0	0
			20	6	9	5		

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	0	0
			20	4	12	1	3		
5	A	1	Total	C	H	N	O	0	0
			20	4	12	1	3		
5	A	1	Total	C	H	N	O	0	0
			20	4	12	1	3		
5	B	1	Total	C	H	N	O	0	0
			20	4	12	1	3		
5	B	1	Total	C	H	N	O	0	0
			20	4	12	1	3		
5	B	1	Total	C	H	N	O	0	0
			20	4	12	1	3		

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	2	Total	Na	0	0
			2	2		
6	A	1	Total	Na	0	0
			1	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			14	3	8	3		
7	A	1	Total	C	H	O	0	0
			14	3	8	3		
7	B	1	Total	C	H	O	0	0
			14	3	8	3		
7	B	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 8 is BETA-L-FUCOSE (three-letter code: FUL) (formula: $C_6H_{12}O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	H	O	0	0
			23	6	12	5		
8	B	1	Total	C	H	O	0	0
			23	6	12	5		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	870	Total	O	0	0
			870	870		
9	B	895	Total	O	0	0
			895	895		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-fucosidase GH29



- Molecule 1: Alpha-fucosidase GH29



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.15Å 75.99Å 80.97Å 105.69° 107.32° 106.93°	Depositor
Resolution (Å)	43.28 – 1.38 43.28 – 1.38	Depositor EDS
% Data completeness (in resolution range)	96.0 (43.28-1.38) 79.5 (43.28-1.38)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 1.38Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.138 , 0.166 0.137 , 0.164	Depositor DCC
R_{free} test set	10215 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	21517	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BMA, NAG, NA, TRS, FUL, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.53	0/5247	0.73	5/7108 (0.1%)
1	B	0.56	0/5287	0.79	9/7164 (0.1%)
All	All	0.54	0/10534	0.76	14/14272 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	212[B]	ARG	NE-CZ-NH1	-14.53	113.03	120.30
1	B	212[C]	ARG	NE-CZ-NH1	-14.53	113.03	120.30
1	B	212[B]	ARG	NE-CZ-NH2	8.21	124.40	120.30
1	B	212[C]	ARG	NE-CZ-NH2	8.21	124.40	120.30
1	B	310[A]	SER	N-CA-CB	-7.26	99.61	110.50
1	B	310[B]	SER	N-CA-CB	-7.26	99.61	110.50
1	A	154[A]	LEU	CA-CB-CG	6.70	130.71	115.30
1	A	154[B]	LEU	CA-CB-CG	6.70	130.71	115.30
1	A	25	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	B	249	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	B	15	ASP	CB-CG-OD1	5.26	123.04	118.30
1	A	249	ARG	NE-CZ-NH1	5.24	122.92	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	414	ASP	CB-CG-OD1	5.08	122.88	118.30
1	A	15	ASP	CB-CG-OD1	5.08	122.87	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	122	PRO	Peptide
1	B	122	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4885	4646	4432	27	0
1	B	4922	4679	4462	22	1
2	A	84	80	74	0	0
2	B	14	14	13	0	0
3	A	22	18	17	0	0
4	A	88	75	75	1	0
5	A	24	36	36	0	0
5	B	24	36	36	0	0
6	A	1	0	0	0	0
6	B	2	0	0	0	0
7	A	12	16	16	0	0
7	B	12	16	16	0	0
8	A	11	12	12	0	0
8	B	11	12	12	0	0
9	A	870	0	0	16	3
9	B	895	0	0	11	7
All	All	11877	9640	9201	51	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160[C]:ARG:NH2	9:A:1527:HOH:O	2.15	0.80
1:A:5:GLY:N	9:A:1488:HOH:O	2.16	0.77
1:B:427[A]:ASN:ND2	9:B:1511:HOH:O	2.22	0.72
1:B:353:GLU:OE1	9:B:1325:HOH:O	2.09	0.70
1:A:188[A]:ASP:OD2	9:A:1114:HOH:O	2.10	0.69
1:A:537:GLU:OE2	9:A:1098:HOH:O	2.10	0.67
1:B:585:ILE:OXT	9:B:1570:HOH:O	2.13	0.67
1:B:3:ALA:N	9:B:1578:HOH:O	2.29	0.65
1:A:163[B]:ASP:OD1	9:A:940:HOH:O	2.16	0.61
1:B:279[B]:ARG:NH1	9:B:1135:HOH:O	2.24	0.61
1:A:289:ARG:HH12	1:A:308[A]:ALA:HA	1.67	0.60
1:B:382:TYR:OH	1:B:388[A]:GLN:NE2	2.34	0.59
1:A:308[A]:ALA:HB2	1:A:320:TYR:CE2	2.39	0.57
1:A:118[A]:ARG:NH2	9:A:1101:HOH:O	2.27	0.56
1:B:212[B]:ARG:NH1	9:B:1589:HOH:O	2.37	0.56
1:B:308[A]:ALA:HB2	1:B:320:TYR:OH	2.06	0.55
1:B:389[B]:VAL:HG12	1:B:391:GLY:H	1.72	0.55
1:B:308[A]:ALA:HB2	1:B:320:TYR:CZ	2.42	0.55
1:A:412:GLU:OE1	9:A:938:HOH:O	2.19	0.53
1:B:212[B]:ARG:HG2	1:B:255:TYR:CE1	2.46	0.50
1:B:535:LYS:NZ	1:B:538:GLU:OE2	2.37	0.49
1:B:307[B]:SER:HB3	9:B:1582:HOH:O	2.11	0.49
1:A:307[B]:SER:HB3	9:A:1526:HOH:O	2.12	0.49
1:A:308[A]:ALA:HB2	1:A:320:TYR:HE2	1.79	0.48
1:A:247[B]:GLN:NE2	9:A:1335:HOH:O	2.45	0.48
1:A:212[B]:ARG:HG3	1:A:255:TYR:CZ	2.48	0.47
1:A:432:LYS:NZ	1:A:469[B]:ASP:OD1	2.47	0.47
1:A:188[B]:ASP:OD2	9:A:831:HOH:O	2.20	0.47
1:A:461:VAL:HG22	1:A:476:PRO:HD3	1.96	0.46
1:A:153[B]:LYS:HD3	9:A:1520:HOH:O	2.17	0.45
1:B:118[B]:ARG:NH2	9:B:1547:HOH:O	2.48	0.44
1:A:461:VAL:HG11	1:A:473:ILE:HG23	2.00	0.44
1:A:11:TRP:CE3	1:A:279[B]:ARG:HD3	2.53	0.44
1:B:208:PHE:HE2	1:B:212[C]:ARG:CZ	2.30	0.44
1:B:497:VAL:HG22	9:B:1268:HOH:O	2.18	0.44
1:A:212[A]:ARG:NH2	9:A:1533:HOH:O	2.51	0.43
1:A:11:TRP:CD2	1:A:250:LEU:HD21	2.53	0.43
1:A:276:HIS:O	9:A:1536:HOH:O	2.22	0.43
1:B:31:VAL:O	1:B:120:ALA:HA	2.19	0.42
1:A:332[B]:ARG:NH1	9:A:1099:HOH:O	2.52	0.42
1:B:290:GLY:HA2	1:B:332[B]:ARG:NH1	2.34	0.42
1:A:153[A]:LYS:HE3	9:A:894:HOH:O	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:GLN:HA	1:B:263:TRP:CH2	2.55	0.42
1:A:332[B]:ARG:NE	9:A:1099:HOH:O	2.49	0.42
1:A:581:VAL:HG21	4:A:606:MAN:H62	2.01	0.41
1:A:328:SER:O	1:A:332[B]:ARG:HG2	2.20	0.41
1:B:328:SER:O	1:B:332[B]:ARG:HG2	2.21	0.41
1:B:405:ILE:HG22	1:B:420[B]:VAL:HG22	2.04	0.40
1:B:436:SER:OG	1:B:468:LYS:HA	2.21	0.40

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:1354:HOH:O	9:B:1275:HOH:O[1_556]	2.02	0.18
9:B:1363:HOH:O	9:B:1412:HOH:O[1_445]	2.08	0.12
9:A:1122:HOH:O	9:B:1176:HOH:O[1_666]	2.09	0.11
9:B:1412:HOH:O	9:B:1457:HOH:O[1_665]	2.11	0.09
9:B:1036:HOH:O	9:B:1569:HOH:O[1_445]	2.13	0.07
9:A:1487:HOH:O	9:B:1289:HOH:O[1_666]	2.17	0.03
1:B:8[B]:GLU:OE1	9:B:1541:HOH:O[1_655]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	638/585 (109%)	624 (98%)	14 (2%)	0	100	100
1	B	645/585 (110%)	630 (98%)	15 (2%)	0	100	100
All	All	1283/1170 (110%)	1254 (98%)	29 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	545/492 (111%)	537 (98%)	8 (2%)	70	38
1	B	550/492 (112%)	547 (100%)	3 (0%)	91	76
All	All	1095/984 (111%)	1084 (99%)	11 (1%)	85	55

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	TYR
1	A	73	TRP
1	A	160[A]	ARG
1	A	160[B]	ARG
1	A	160[C]	ARG
1	A	171	HIS
1	A	310[A]	SER
1	A	310[B]	SER
1	B	32	TYR
1	B	73	TRP
1	B	171	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 32 ligands modelled in this entry, 3 are monoatomic - leaving 29 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	601	1,2	14,14,15	0.38	0	15,19,21	0.57	0
2	NAG	A	602	3,2	14,14,15	0.25	0	15,19,21	0.54	0
3	BMA	A	603	2,4	11,11,12	1.01	1 (9%)	13,15,17	0.86	0
4	MAN	A	604	3	11,11,12	1.21	2 (18%)	13,15,17	1.16	2 (15%)
4	MAN	A	605	3,4	11,11,12	0.88	1 (9%)	13,15,17	1.31	2 (15%)
4	MAN	A	606	4	11,11,12	1.61	4 (36%)	13,15,17	1.03	1 (7%)
4	MAN	A	607	4	11,11,12	1.66	3 (27%)	13,15,17	1.14	2 (15%)
2	NAG	A	608	1	14,14,15	0.13	0	15,19,21	0.48	0
2	NAG	A	609	1	14,14,15	0.36	0	15,19,21	0.50	0
4	MAN	A	610[A]	3,4	11,11,12	0.98	1 (9%)	13,15,17	1.13	1 (7%)
4	MAN	A	610[B]	3,4	11,11,12	0.79	1 (9%)	13,15,17	0.96	1 (7%)
4	MAN	A	611	4	11,11,12	1.17	1 (9%)	13,15,17	1.04	1 (7%)
4	MAN	A	612	4	11,11,12	0.83	1 (9%)	13,15,17	1.07	1 (7%)
2	NAG	A	613	1,2	14,14,15	0.43	0	15,19,21	0.75	1 (6%)
2	NAG	A	614	3,2	14,14,15	0.22	0	15,19,21	0.54	0
3	BMA	A	615	2,4	11,11,12	1.11	0	13,15,17	0.95	0
5	TRS	A	616	-	7,7,7	0.71	0	9,9,9	0.70	0
5	TRS	A	617	-	7,7,7	0.69	0	9,9,9	0.82	0
5	TRS	A	618	-	7,7,7	0.83	1 (14%)	9,9,9	0.76	0
7	GOL	A	620	-	5,5,5	0.29	0	5,5,5	0.18	0
7	GOL	A	621	-	5,5,5	0.32	0	5,5,5	0.23	0
8	FUL	A	622	-	11,11,11	0.52	0	15,16,16	1.09	1 (6%)
2	NAG	B	601	1	14,14,15	0.22	0	15,19,21	0.46	0
5	TRS	B	602	-	7,7,7	0.65	0	9,9,9	0.83	1 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	TRS	B	603	-	7,7,7	0.65	0	9,9,9	0.79	0
5	TRS	B	604	-	7,7,7	0.58	0	9,9,9	0.57	0
7	GOL	B	607	-	5,5,5	0.28	0	5,5,5	0.28	0
7	GOL	B	608	-	5,5,5	0.29	0	5,5,5	0.51	0
8	FUL	B	609	-	11,11,11	0.53	0	15,16,16	0.95	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	601	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	602	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	603	2,4	-	0/2/19/22	0/1/1/1
4	MAN	A	604	3	-	0/2/19/22	0/1/1/1
4	MAN	A	605	3,4	-	0/2/19/22	0/1/1/1
4	MAN	A	606	4	-	0/2/19/22	0/1/1/1
4	MAN	A	607	4	-	0/2/19/22	0/1/1/1
2	NAG	A	608	1	-	0/6/23/26	0/1/1/1
2	NAG	A	609	1	-	0/6/23/26	0/1/1/1
4	MAN	A	610[A]	3,4	-	0/2/19/22	0/1/1/1
4	MAN	A	610[B]	3,4	-	0/2/19/22	0/1/1/1
4	MAN	A	611	4	-	0/2/19/22	0/1/1/1
4	MAN	A	612	4	-	0/2/19/22	0/1/1/1
2	NAG	A	613	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	614	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	615	2,4	-	0/2/19/22	0/1/1/1
5	TRS	A	616	-	-	0/9/9/9	0/0/0/0
5	TRS	A	617	-	-	0/9/9/9	0/0/0/0
5	TRS	A	618	-	-	0/9/9/9	0/0/0/0
7	GOL	A	620	-	-	0/4/4/4	0/0/0/0
7	GOL	A	621	-	-	0/4/4/4	0/0/0/0
8	FUL	A	622	-	-	0/0/20/20	0/1/1/1
2	NAG	B	601	1	-	0/6/23/26	0/1/1/1
5	TRS	B	602	-	-	0/9/9/9	0/0/0/0
5	TRS	B	603	-	-	0/9/9/9	0/0/0/0
5	TRS	B	604	-	-	0/9/9/9	0/0/0/0
7	GOL	B	607	-	-	0/4/4/4	0/0/0/0
7	GOL	B	608	-	-	0/4/4/4	0/0/0/0
8	FUL	B	609	-	-	0/0/20/20	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	606	MAN	O5-C1	-2.95	1.38	1.43
3	A	603	BMA	O5-C1	-2.51	1.39	1.43
4	A	610[A]	MAN	O5-C1	-2.32	1.39	1.43
4	A	611	MAN	O5-C1	-2.29	1.40	1.43
4	A	605	MAN	O5-C1	-2.21	1.40	1.43
5	A	618	TRS	C-N	-2.16	1.47	1.50
4	A	607	MAN	O5-C1	-2.15	1.40	1.43
4	A	610[B]	MAN	O5-C1	-2.15	1.40	1.43
4	A	606	MAN	O2-C2	-2.08	1.38	1.43
4	A	604	MAN	O5-C5	2.11	1.47	1.43
4	A	606	MAN	C2-C3	2.19	1.55	1.52
4	A	612	MAN	O5-C5	2.31	1.48	1.43
4	A	606	MAN	C1-C2	2.38	1.57	1.52
4	A	604	MAN	C2-C3	2.47	1.55	1.52
4	A	607	MAN	C1-C2	2.61	1.58	1.52
4	A	607	MAN	O5-C5	2.88	1.49	1.43

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	605	MAN	O2-C2-C3	-3.14	104.01	110.17
8	A	622	FUL	C1-C2-C3	-2.60	105.96	110.65
4	A	612	MAN	O2-C2-C3	-2.58	105.10	110.17
4	A	606	MAN	O2-C2-C3	-2.42	105.43	110.17
8	B	609	FUL	C1-C2-C3	-2.36	106.38	110.65
4	A	610[B]	MAN	O2-C2-C3	-2.31	105.64	110.17
4	A	607	MAN	O2-C2-C3	-2.23	105.79	110.17
4	A	604	MAN	O2-C2-C3	-2.22	105.81	110.17
2	A	613	NAG	O5-C1-C2	-2.07	108.59	111.47
5	B	602	TRS	C2-C-C1	-2.00	106.92	110.71
4	A	611	MAN	C1-O5-C5	2.49	115.59	112.17
4	A	604	MAN	C1-O5-C5	2.59	115.74	112.17
4	A	605	MAN	C1-O5-C5	2.65	115.82	112.17
4	A	607	MAN	C1-O5-C5	2.66	115.83	112.17
4	A	610[A]	MAN	C1-O5-C5	3.00	116.30	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	606	MAN	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	581/585 (99%)	-0.12	10 (1%)	70 73	9, 15, 33, 73	0
1	B	583/585 (99%)	-0.26	12 (2%)	64 67	9, 14, 30, 77	0
All	All	1164/1170 (99%)	-0.19	22 (1%)	67 70	9, 15, 32, 77	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	556	PHE	5.1
1	A	391	GLY	4.1
1	B	392	GLY	4.0
1	A	585	ILE	3.6
1	A	308[A]	ALA	3.5
1	B	391	GLY	3.4
1	B	574	ASP	3.4
1	B	395	THR	3.3
1	B	585	ILE	3.1
1	B	394	PHE	2.9
1	B	393	SER	2.8
1	A	392	GLY	2.7
1	B	308[A]	ALA	2.7
1	A	394	PHE	2.6
1	B	575	GLY	2.5
1	A	462	SER	2.5
1	A	428	LEU	2.4
1	A	510	ALA	2.2
1	B	573	GLU	2.0
1	B	556	PHE	2.0
1	A	576	SER	2.0
1	B	309[A]	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	MAN	A	607	11/12	0.80	0.24	13.79	31,49,62,69	0
5	TRS	A	616	8/8	0.85	0.17	10.80	21,28,38,38	0
4	MAN	A	612	11/12	0.93	0.17	7.89	16,23,34,34	0
4	MAN	A	606	11/12	0.87	0.12	4.96	21,32,40,42	0
5	TRS	A	618	8/8	0.91	0.19	2.32	29,39,45,49	0
4	MAN	A	611	11/12	0.92	0.19	2.20	23,37,58,70	0
4	MAN	A	610[B]	11/12	0.95	0.09	2.14	14,18,30,31	20
4	MAN	A	610[A]	11/12	0.95	0.09	2.02	14,18,38,38	20
5	TRS	B	603	8/8	0.93	0.12	1.78	16,21,46,48	0
5	TRS	A	617	8/8	0.92	0.12	1.78	17,24,44,53	0
7	GOL	A	620	6/6	0.72	0.14	1.63	36,44,57,59	0
5	TRS	B	604	8/8	0.93	0.11	1.42	16,22,26,29	0
7	GOL	B	607	6/6	0.81	0.15	1.36	33,48,63,65	0
7	GOL	A	621	6/6	0.84	0.15	0.52	40,49,56,65	0
8	FUL	B	609	11/11	0.98	0.07	0.40	12,15,18,18	0
5	TRS	B	602	8/8	0.93	0.09	0.33	21,27,42,51	0
6	NA	B	605	1/1	0.98	0.08	0.26	17,17,17,17	0
2	NAG	A	608	14/15	0.96	0.10	0.08	16,23,56,56	0
2	NAG	A	613	14/15	0.97	0.07	-0.07	11,14,22,34	0
4	MAN	A	605	11/12	0.97	0.07	-0.13	14,18,38,38	0
2	NAG	A	601	14/15	0.97	0.06	-0.30	12,16,20,32	0
2	NAG	A	609	14/15	0.95	0.06	-0.43	17,25,34,40	0
6	NA	B	606	1/1	0.99	0.06	-0.51	13,13,13,13	0
2	NAG	B	601	14/15	0.94	0.08	-0.53	14,25,38,47	0
6	NA	A	619	1/1	1.00	0.06	-0.61	15,15,15,15	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
8	FUL	A	622	11/11	0.97	0.06	-1.49	12,15,18,19	0
7	GOL	B	608	6/6	0.87	0.18	-	39,55,66,76	0
3	BMA	A	603	11/12	0.97	0.06	-	15,18,22,22	0
2	NAG	A	614	14/15	0.96	0.07	-	13,18,29,29	0
3	BMA	A	615	11/12	0.94	0.09	-	15,18,25,26	0
4	MAN	A	604	11/12	0.77	0.19	-	32,51,62,72	0
2	NAG	A	602	14/15	0.96	0.07	-	14,18,30,30	0

6.5 Other polymers

There are no such residues in this entry.