



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 26, 2017 – 02:23 PM EDT

PDB ID : 5PWF  
Title : PanDDA analysis group deposition – Crystal Structure of SP100 after initial refinement with no ligand modelled (structure 2)  
Authors : Pearce, N.M.; Krojer, T.; Talon, R.; Bradley, A.R.; Fairhead, M.; Sethi, R.; Wright, N.; MacLean, E.; Collins, P.; Brandao-Neto, J.; Douangamath, A.; Renjie, Z.; Dias, A.; Ng, J.; Brennan, P.E.; Cox, O.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; von Delft, F.  
Deposited on : unknown  
Resolution : 1.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3517 (1.50-1.46)
Clashscore	112137	3795 (1.50-1.46)
Ramachandran outliers	110173	3721 (1.50-1.46)
Sidechain outliers	110143	3719 (1.50-1.46)
RSRZ outliers	101464	3549 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	<div> <div>6%</div> <div>86%</div> <div>12%</div> <div>..</div> </div>
1	B	180	<div> <div>12%</div> <div>86%</div> <div>11%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ZN	A	903	-	-	-	X
3	MES	A	904	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3620 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nuclear autoantigen Sp-100.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	178	Total	C	N	O	S	0	10	0
			1536	971	268	278	19			
1	B	177	Total	C	N	O	S	0	7	0
			1480	940	258	263	19			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Zn	0	0
			2	2		
2	A	3	Total	Zn	0	0
			3	3		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	313	Total	O	0	3
			313	313		
5	B	266	Total	O	0	0
			266	266		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nuclear autoantigen Sp-100



- Molecule 1: Nuclear autoantigen Sp-100



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	127.59Å 45.33Å 83.18Å 90.00° 102.22° 90.00°	Depositor
Resolution (Å)	28.40 – 1.48 28.40 – 1.48	Depositor EDS
% Data completeness (in resolution range)	92.1 (28.40-1.48) 92.2 (28.40-1.48)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.14 (at 1.48Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.178 , 0.206 0.188 , 0.218	Depositor DCC
$R_{free}$ test set	3583 reflections (5.25%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.0	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 50.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3620	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EDO, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.26	7/1603 (0.4%)	1.15	12/2154 (0.6%)
1	B	1.26	4/1538 (0.3%)	1.20	5/2071 (0.2%)
All	All	1.26	11/3141 (0.4%)	1.18	17/4225 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	742	CYS	CB-SG	-10.99	1.63	1.82
1	A	763	GLU	CD-OE1	-8.51	1.16	1.25
1	A	797	SER	CB-OG	-6.63	1.33	1.42
1	B	843	ASN	N-CA	-5.66	1.35	1.46
1	A	723	SER	CA-CB	5.46	1.61	1.52
1	A	821	SER	CB-OG	5.40	1.49	1.42
1	B	732	SER	CB-OG	-5.31	1.35	1.42
1	B	837	MET	N-CA	5.21	1.56	1.46
1	B	838	ARG	CZ-NH2	-5.06	1.26	1.33
1	A	803[A]	ARG	CZ-NH2	-5.01	1.26	1.33
1	A	803[B]	ARG	CZ-NH2	-5.01	1.26	1.33

All (17) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	829[A]	ARG	NE-CZ-NH2	-13.20	113.70	120.30
1	B	829[B]	ARG	NE-CZ-NH2	-13.20	113.70	120.30
1	B	789	ASP	CB-CG-OD2	-8.11	111.00	118.30
1	A	851	ASP	CB-CG-OD1	7.56	125.11	118.30
1	A	803[A]	ARG	NE-CZ-NH1	6.78	123.69	120.30
1	A	803[B]	ARG	NE-CZ-NH1	6.78	123.69	120.30
1	B	855	ARG	NE-CZ-NH2	-6.73	116.93	120.30
1	A	718	ASP	CB-CG-OD2	6.68	124.31	118.30
1	B	855	ARG	NE-CZ-NH1	6.52	123.56	120.30
1	A	742	CYS	CA-CB-SG	6.07	124.93	114.00
1	A	787	TYR	CB-CG-CD2	5.59	124.35	121.00
1	A	801	TYR	CB-CG-CD1	-5.55	117.67	121.00
1	A	718	ASP	CB-CG-OD1	-5.44	113.40	118.30
1	A	868	PHE	CB-CG-CD2	-5.42	117.01	120.80
1	A	803[A]	ARG	NE-CZ-NH2	-5.16	117.72	120.30
1	A	803[B]	ARG	NE-CZ-NH2	-5.16	117.72	120.30
1	A	713	ARG	NE-CZ-NH2	5.11	122.86	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	755	PRO	Peptide
1	B	759	SER	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1536	0	1497	12	0
1	B	1480	0	1414	9	0
2	A	3	0	0	0	0
2	B	2	0	0	0	0
3	A	12	0	12	2	0
4	A	4	0	6	2	0
4	B	4	0	6	1	0
5	A	313	0	0	12	1
5	B	266	0	0	7	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3620	0	2935	25	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:803[A]:ARG:NH2	5:A:1001:HOH:O	1.73	1.20
1:A:803[B]:ARG:NH1	5:A:1003:HOH:O	1.99	0.95
1:B:716[A]:CYS:SG	5:B:1166:HOH:O	2.28	0.89
1:A:829[A]:ARG:NH1	5:A:1002:HOH:O	1.97	0.88
3:A:904:MES:H81	5:A:1237:HOH:O	1.80	0.82
1:A:756:GLU:HG3	5:A:1112:HOH:O	1.84	0.76
1:B:750:ILE:HD11	5:B:1001:HOH:O	1.91	0.71
5:A:1001:HOH:O	1:B:773:PRO:HG2	1.92	0.68
1:A:862[A]:ASP:OD1	5:A:1004:HOH:O	2.10	0.67
1:B:826[B]:MET:SD	5:B:1018:HOH:O	2.51	0.67
4:B:903:EDO:O2	5:B:1001:HOH:O	2.12	0.66
1:B:826[B]:MET:HE2	5:B:1195:HOH:O	1.96	0.65
1:A:802:ASN:O	3:A:904:MES:H32	2.03	0.58
1:A:824:GLU:OE1	1:A:826[B]:MET:HE2	2.04	0.57
1:B:849:ARG:O	1:B:854[B]:THR:HG21	2.05	0.57
1:A:824:GLU:OE1	1:A:826[B]:MET:CE	2.57	0.53
4:A:905:EDO:C2	5:A:1065:HOH:O	2.57	0.52
1:A:762:GLN:NE2	5:A:1014:HOH:O	2.43	0.50
1:B:817:LYS:NZ	5:B:1009:HOH:O	2.46	0.49
1:A:734:GLU:OE1	1:A:740:TRP:NE1	2.33	0.48
1:A:748:LYS:HE2	5:A:1238:HOH:O	2.15	0.47
1:B:757:SER:C	1:B:759:SER:H	2.18	0.46
4:A:905:EDO:C1	5:A:1229:HOH:O	2.66	0.43
1:A:701:ASN:N	5:A:1020:HOH:O	2.53	0.42
1:B:785:LYS:HE3	5:B:1221:HOH:O	2.19	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:1024:HOH:O	5:B:1024:HOH:O[2_556]	1.33	0.87
5:B:1197:HOH:O	5:B:1197:HOH:O[2_555]	1.94	0.26

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1134:HOH:O	5:A:1239:HOH:O[1_545]	2.01	0.19

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/180 (103%)	180 (97%)	6 (3%)	0	100	100
1	B	182/180 (101%)	172 (94%)	8 (4%)	2 (1%)	17	3
All	All	368/360 (102%)	352 (96%)	14 (4%)	2 (0%)	28	9

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	757	SER
1	B	758	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/171 (104%)	178 (100%)	0	100	100
1	B	165/171 (96%)	160 (97%)	5 (3%)	46	13
All	All	343/342 (100%)	338 (98%)	5 (2%)	82	40

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	711	TRP
1	B	850[A]	GLU
1	B	850[B]	GLU
1	B	862[A]	ASP
1	B	862[B]	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	751	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	MES	A	904	-	12,12,12	0.85	0	14,16,16	3.15	7 (50%)
4	EDO	A	905	-	3,3,3	0.42	0	2,2,2	0.30	0
4	EDO	B	903	-	3,3,3	0.72	0	2,2,2	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MES	A	904	-	-	0/6/14/14	0/1/1/1
4	EDO	A	905	-	-	0/1/1/1	0/0/0/0
4	EDO	B	903	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	904	MES	O2S-S-C8	-4.36	103.05	106.79
3	A	904	MES	O1S-S-C8	-3.03	104.19	106.79
3	A	904	MES	C7-N4-C3	3.58	120.44	111.26
3	A	904	MES	C6-O1-C2	3.75	122.57	109.89
3	A	904	MES	C2-C3-N4	4.35	116.20	110.11
3	A	904	MES	C6-C5-N4	4.50	116.41	110.11
3	A	904	MES	C5-N4-C3	6.10	122.68	108.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	904	MES	2	0
4	A	905	EDO	2	0
4	B	903	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	178/180 (98%)	0.12	11 (6%) 21 24	12, 20, 54, 93	0
1	B	177/180 (98%)	0.57	22 (12%) 4 5	13, 23, 75, 139	0
All	All	355/360 (98%)	0.34	33 (9%) 9 11	12, 22, 66, 139	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	757	SER	13.6
1	A	735	ALA	9.7
1	B	806	SER	9.5
1	B	758	GLN	8.4
1	B	759	SER	7.2
1	B	805	GLY	6.3
1	B	800	TYR	6.2
1	B	760	GLY	5.4
1	B	876	GLU	5.1
1	B	804	GLU	5.0
1	A	736	ASN	4.9
1	B	803	ARG	4.5
1	B	735	ALA	4.5
1	B	736	ASN	4.1
1	B	711	TRP	4.0
1	B	756	GLU	3.6
1	B	875	GLN	3.5
1	A	703	ASN	3.1
1	A	878	SER	3.0
1	A	737	LYS	2.9
1	B	783	LEU	2.6
1	A	701	ASN	2.6
1	B	808	GLY	2.6
1	B	810	GLN	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	782	LEU	2.5
1	B	738	ASN	2.4
1	A	738	ASN	2.3
1	B	802	ASN	2.3
1	A	739	PRO	2.2
1	B	755	PRO	2.1
1	B	850[A]	GLU	2.1
1	A	783	LEU	2.1
1	A	849	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	A	903	1/1	0.15	0.26	6.90	93,93,93,93	1
3	MES	A	904	12/12	0.97	0.17	6.12	14,16,20,20	12
2	ZN	B	902	1/1	0.98	0.05	-0.94	21,21,21,21	0
2	ZN	A	902	1/1	0.92	0.06	-1.29	28,28,28,28	0
4	EDO	B	903	4/4	0.98	0.05	-1.40	27,27,28,28	0
2	ZN	B	901	1/1	0.99	0.06	-	18,18,18,18	0
4	EDO	A	905	4/4	0.91	0.40	-	42,49,53,64	0
2	ZN	A	901	1/1	0.97	0.06	-	21,21,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.