



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:00 am GMT

PDB ID : 3PYF  
Title : Mycobacterium tuberculosis 4-diphosphocytidyl-2-C-methyl-D-erythritol kinase (IspE) in complex with AMP-PNP  
Authors : Shan, S.; Chen, X.H.; Liu, T.; Zhao, H.C.; Rao, Z.H.; Lou, Z.Y.  
Deposited on : 2010-12-13  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3453 (1.70-1.70)
Clashscore	112137	3876 (1.70-1.70)
Ramachandran outliers	110173	3815 (1.70-1.70)
Sidechain outliers	110143	3815 (1.70-1.70)
RSRZ outliers	101464	3491 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ANP	A	307	-	-	-	X

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2324 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-diphosphocytidylyl-2-C-methyl-D-erythritol kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	296	2134	1329	384	412	9	0	0	0

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula:  $C_{10}H_{17}N_6O_{12}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	31	10	6	12	3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	159	Total	O	0	0
			159	159		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-diphosphocytidyl-2-C-methyl-D-erythritol kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.96Å 73.48Å 106.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.42 – 1.70 33.42 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (33.42-1.70) 99.9 (33.42-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.06 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.229 , 0.249 0.227 , 0.248	Depositor DCC
$R_{free}$ test set	2347 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	22.3	Xtriage
Anisotropy	0.403	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 39.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2324	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.86% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.34	9/2168 (0.4%)	1.40	28/2959 (0.9%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	246	GLU	CG-CD	7.49	1.63	1.51
1	A	85	ARG	CD-NE	-6.91	1.34	1.46
1	A	124	ASN	CB-CG	6.89	1.66	1.51
1	A	5	SER	N-CA	6.63	1.59	1.46
1	A	68	GLU	CG-CD	6.28	1.61	1.51
1	A	51	VAL	CB-CG1	-6.21	1.39	1.52
1	A	255	SER	CB-OG	6.04	1.50	1.42
1	A	246	GLU	CB-CG	5.98	1.63	1.52
1	A	81	GLU	CG-CD	5.72	1.60	1.51

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	ARG	NE-CZ-NH1	15.90	128.25	120.30
1	A	85	ARG	NE-CZ-NH2	-15.02	112.79	120.30
1	A	190	ARG	NE-CZ-NH2	-13.78	113.41	120.30
1	A	298	ARG	NE-CZ-NH2	-9.16	115.72	120.30
1	A	298	ARG	NE-CZ-NH1	8.88	124.74	120.30
1	A	172	LEU	CB-CG-CD1	8.26	125.04	111.00
1	A	190	ARG	CG-CD-NE	-7.68	95.67	111.80
1	A	254	VAL	CG1-CB-CG2	7.45	122.82	110.90
1	A	163	LEU	CB-CG-CD1	7.42	123.61	111.00
1	A	202	GLU	C-N-CD	7.09	143.30	128.40
1	A	200	LEU	CB-CG-CD1	7.01	122.91	111.00
1	A	288	ARG	NE-CZ-NH1	6.96	123.78	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	85	ARG	CG-CD-NE	-6.77	97.59	111.80
1	A	163	LEU	CB-CG-CD2	6.74	122.46	111.00
1	A	124	ASN	CB-CA-C	6.61	123.61	110.40
1	A	145	LEU	CB-CG-CD1	6.49	122.04	111.00
1	A	85	ARG	CD-NE-CZ	6.19	132.26	123.60
1	A	17	TYR	CZ-CE2-CD2	-5.97	114.42	119.80
1	A	88	ASP	CB-CG-OD2	5.88	123.59	118.30
1	A	85	ARG	CB-CG-CD	-5.73	96.70	111.60
1	A	190	ARG	CD-NE-CZ	5.59	131.42	123.60
1	A	187	GLU	OE1-CD-OE2	5.42	129.81	123.30
1	A	46	VAL	CG1-CB-CG2	-5.28	102.45	110.90
1	A	9	ARG	NE-CZ-NH2	-5.27	117.66	120.30
1	A	13	LYS	CD-CE-NZ	-5.20	99.73	111.70
1	A	145	LEU	CA-CB-CG	5.16	127.17	115.30
1	A	261	CYS	CA-CB-SG	-5.15	104.73	114.00
1	A	88	ASP	CB-CG-OD1	-5.07	113.73	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2134	0	2141	63	0
2	A	31	0	13	8	0
3	A	159	0	0	14	0
All	All	2324	0	2154	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:ARG:HB2	3:A:448:HOH:O	1.36	1.21

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:GLU:HA	1:A:27:GLY:H	1.04	1.20
1:A:46:VAL:CG1	1:A:89:VAL:HG21	1.78	1.13
1:A:25:GLU:HA	1:A:27:GLY:N	1.72	1.01
1:A:47:ARG:HD2	3:A:429:HOH:O	1.66	0.94
1:A:298:ARG:HH11	1:A:298:ARG:HG2	1.31	0.92
1:A:46:VAL:CG1	1:A:89:VAL:CG2	2.47	0.92
1:A:7:THR:HB	3:A:450:HOH:O	0.72	0.89
1:A:85:ARG:HD2	3:A:390:HOH:O	1.73	0.88
1:A:7:THR:HG23	3:A:400:HOH:O	1.78	0.82
1:A:46:VAL:HG12	1:A:89:VAL:HG21	1.59	0.82
1:A:85:ARG:CD	3:A:390:HOH:O	2.27	0.81
1:A:7:THR:HG22	1:A:45:THR:OG1	1.80	0.80
1:A:46:VAL:HG12	1:A:89:VAL:CG2	2.10	0.80
1:A:176:ASP:N	1:A:176:ASP:OD1	2.17	0.73
1:A:47:ARG:O	1:A:89:VAL:HG23	1.89	0.73
1:A:46:VAL:HG13	1:A:89:VAL:CG2	2.18	0.72
1:A:242:ARG:O	1:A:246:GLU:HG3	1.90	0.72
1:A:33:THR:HG21	3:A:464:HOH:O	1.90	0.72
1:A:71:LEU:H	2:A:307:ANP:H2	1.56	0.70
1:A:36:HIS:HE1	1:A:170:TRP:HE1	1.39	0.69
1:A:227:GLN:HG3	1:A:254:VAL:HG13	1.77	0.67
1:A:104:ALA:H	2:A:307:ANP:HNB1	1.44	0.65
1:A:5:SER:HB2	1:A:47:ARG:CG	2.27	0.65
1:A:29:HIS:HE1	1:A:189:ASP:OD1	1.80	0.63
1:A:71:LEU:H	2:A:307:ANP:C2	2.13	0.61
1:A:46:VAL:HG13	1:A:89:VAL:HG21	1.71	0.61
1:A:31:LEU:H	1:A:155:ARG:HH22	1.47	0.61
1:A:82:HIS:HD2	3:A:453:HOH:O	1.83	0.60
1:A:298:ARG:HH11	1:A:298:ARG:CG	2.07	0.60
1:A:5:SER:HB2	1:A:47:ARG:HG2	1.83	0.60
1:A:33:THR:HG23	3:A:330:HOH:O	2.00	0.60
1:A:127:ARG:NH2	3:A:442:HOH:O	2.35	0.59
1:A:70:ASN:HD22	1:A:72:ALA:H	1.49	0.58
1:A:78:LEU:HG	1:A:133:LEU:HD22	1.86	0.57
1:A:259:PRO:HA	3:A:347:HOH:O	2.03	0.57
1:A:85:ARG:HD3	3:A:390:HOH:O	2.00	0.57
1:A:298:ARG:HG2	1:A:298:ARG:NH1	2.07	0.56
1:A:25:GLU:HB2	1:A:26:ASP:HA	1.88	0.55
1:A:24:ARG:HH11	1:A:28:TYR:HB3	1.71	0.55
1:A:111:ALA:HB1	1:A:145:LEU:HD13	1.88	0.55
1:A:36:HIS:CE1	1:A:170:TRP:HE1	2.23	0.52

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:VAL:HG13	1:A:89:VAL:HG22	1.92	0.52
1:A:61:ALA:HA	1:A:64:LEU:HD22	1.92	0.52
2:A:307:ANP:O1B	2:A:307:ANP:O5'	2.24	0.52
1:A:153:THR:HG23	1:A:157:GLU:HG3	1.92	0.50
1:A:78:LEU:HD11	1:A:136:ARG:CZ	2.41	0.50
1:A:47:ARG:CD	3:A:429:HOH:O	2.40	0.50
1:A:127:ARG:HD3	3:A:309:HOH:O	2.12	0.49
1:A:46:VAL:HG11	1:A:89:VAL:HG21	1.86	0.47
1:A:298:ARG:NH1	1:A:298:ARG:CG	2.71	0.47
1:A:70:ASN:ND2	2:A:307:ANP:N1	2.63	0.46
1:A:181:THR:HG22	1:A:257:SER:OG	2.15	0.46
1:A:5:SER:HB2	1:A:47:ARG:HG3	1.98	0.44
1:A:70:ASN:HD21	2:A:307:ANP:HN62	1.64	0.44
1:A:64:LEU:HG	2:A:307:ANP:N7	2.32	0.44
1:A:25:GLU:CB	1:A:26:ASP:HA	2.47	0.44
1:A:70:ASN:ND2	1:A:72:ALA:H	2.16	0.44
1:A:36:HIS:HD2	1:A:149:THR:OG1	2.01	0.44
1:A:202:GLU:O	1:A:205:PRO:HD2	2.18	0.43
2:A:307:ANP:H8	2:A:307:ANP:H5'2	2.00	0.43
1:A:227:GLN:NE2	1:A:254:VAL:H	2.17	0.42
1:A:244:GLY:HA3	1:A:263:PHE:CE2	2.54	0.42
1:A:71:LEU:HB3	1:A:110:ALA:HB2	2.03	0.41
1:A:68:GLU:HA	1:A:73:TRP:CD1	2.56	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	294/306 (96%)	286 (97%)	6 (2%)	2 (1%)	25 9

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	201	GLY
1	A	157	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	218/226 (96%)	197 (90%)	21 (10%)	10 2

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	LYS
1	A	24	ARG
1	A	32	THR
1	A	33	THR
1	A	64	LEU
1	A	70	ASN
1	A	85	ARG
1	A	128	ARG
1	A	145	LEU
1	A	153	THR
1	A	155	ARG
1	A	163	LEU
1	A	172	LEU
1	A	176	ASP
1	A	196	ASP
1	A	199	ARG
1	A	200	LEU
1	A	222	LEU
1	A	235	PRO
1	A	254	VAL
1	A	298	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	HIS
1	A	36	HIS
1	A	70	ASN
1	A	82	HIS
1	A	169	HIS
1	A	227	GLN
1	A	277	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ANP	A	307	-	29,33,33	3.39	13 (44%)	28,52,52	3.98	13 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	307	-	-	0/13/38/38	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	307	ANP	C6-C5	2.04	1.53	1.42
2	A	307	ANP	PB-O3A	2.08	1.61	1.59
2	A	307	ANP	C6-N1	2.15	1.46	1.37
2	A	307	ANP	PA-O1A	2.20	1.59	1.50
2	A	307	ANP	PB-N3B	2.40	1.69	1.63
2	A	307	ANP	C8-N7	2.42	1.39	1.34
2	A	307	ANP	O4'-C1'	4.21	1.47	1.41
2	A	307	ANP	C2-N1	4.92	1.43	1.33
2	A	307	ANP	C6-N6	5.04	1.54	1.34
2	A	307	ANP	PG-N3B	5.90	1.79	1.63
2	A	307	ANP	C5-C4	6.36	1.54	1.40
2	A	307	ANP	PB-O1B	6.73	1.53	1.46
2	A	307	ANP	PG-O1G	9.69	1.57	1.46

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	307	ANP	N3-C2-N1	-9.51	120.58	128.86
2	A	307	ANP	PA-O3A-PB	-8.12	103.70	132.38
2	A	307	ANP	C5-C6-N1	-5.97	101.64	119.70
2	A	307	ANP	O1B-PB-N3B	-5.50	103.56	111.79
2	A	307	ANP	C4-C5-N7	-5.35	104.24	109.41
2	A	307	ANP	O2A-PA-O5'	-2.28	97.36	108.14
2	A	307	ANP	C1'-N9-C4	-2.17	122.89	126.64
2	A	307	ANP	O3A-PB-N3B	2.24	112.79	106.59
2	A	307	ANP	O3G-PG-O2G	2.98	116.03	107.69
2	A	307	ANP	O2A-PA-O1A	3.09	128.27	112.28
2	A	307	ANP	C5-C6-N6	3.37	127.34	120.47
2	A	307	ANP	N6-C6-N1	6.17	131.00	118.77
2	A	307	ANP	C2-N1-C6	9.48	135.36	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	307	ANP	8	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	296/306 (96%)	0.35	25 (8%) <b>12</b> <b>14</b>	13, 22, 46, 66	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	155	ARG	8.3
1	A	26	ASP	7.0
1	A	156	GLY	6.8
1	A	283	VAL	6.8
1	A	28	TYR	6.4
1	A	27	GLY	5.8
1	A	280	GLY	5.0
1	A	154	GLY	4.9
1	A	24	ARG	4.3
1	A	281	ALA	4.1
1	A	176	ASP	3.7
1	A	157	GLU	3.7
1	A	25	GLU	3.6
1	A	196	ASP	3.5
1	A	153	THR	2.6
1	A	193	GLU	2.5
1	A	158	GLU	2.4
1	A	246	GLU	2.3
1	A	201	GLY	2.2
1	A	194	VAL	2.1
1	A	30	GLU	2.1
1	A	23	ARG	2.1
1	A	198	PRO	2.1
1	A	177	SER	2.0
1	A	47	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ANP	A	307	31/31	0.78	0.39	19.34	16,46,55,56	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.