



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 03:17 am GMT

PDB ID : 4PZF  
Title : Berberine bridge enzyme G164A variant, a reticuline dehydrogenase  
Authors : Zafred, D.; Wallner, S.; Steiner, B.; Macheroux, P.  
Deposited on : 2014-03-30  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	538	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>6%</div> <div>7%</div> </div> </div>
1	B	538	<div> <div>4%</div> <div> <div></div> <div>85%</div> <div>7%</div> <div>7%</div> </div> </div>
1	C	538	<div> <div>6%</div> <div> <div></div> <div>80%</div> <div>12%</div> <div>8%</div> </div> </div>
1	D	538	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>8%</div> <div>8%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	602	-	-	-	X
5	SO4	B	605	-	-	X	-
5	SO4	D	606	-	-	-	X
5	SO4	D	607	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 16476 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reticuline oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	498	Total	C	N	O	S	0	0	0
			3935	2531	653	738	13			
1	B	498	Total	C	N	O	S	0	0	0
			3935	2531	654	737	13			
1	C	496	Total	C	N	O	S	0	0	0
			3916	2518	651	734	13			
1	D	497	Total	C	N	O	S	0	0	0
			3927	2527	652	735	13			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	GLU	LEU	SEE REMARK 999	UNP P30986
A	23	ALA	GLY	SEE REMARK 999	UNP P30986
A	164	ALA	GLY	ENGINEERED MUTATION	UNP P30986
B	22	GLU	LEU	SEE REMARK 999	UNP P30986
B	23	ALA	GLY	SEE REMARK 999	UNP P30986
B	164	ALA	GLY	ENGINEERED MUTATION	UNP P30986
C	22	GLU	LEU	SEE REMARK 999	UNP P30986
C	23	ALA	GLY	SEE REMARK 999	UNP P30986
C	164	ALA	GLY	ENGINEERED MUTATION	UNP P30986
D	22	GLU	LEU	SEE REMARK 999	UNP P30986
D	23	ALA	GLY	SEE REMARK 999	UNP P30986
D	164	ALA	GLY	ENGINEERED MUTATION	UNP P30986

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	B	2	Total	C	N	O	0	0
			28	16	2	10		
3	C	2	Total	C	N	O	0	0
			28	16	2	10		
3	D	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0

- Molecule 6 is DODECAETHYLENE GLYCOL (three-letter code: 12P) (formula: C<sub>24</sub>H<sub>50</sub>O<sub>13</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	1	Total C O 28 18 10	0	0
6	D	1	Total C O 22 14 8	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	105	Total O 105 105	0	0
7	B	71	Total O 71 71	0	0
7	C	72	Total O 72 72	0	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	60	Total	O	0	0
			60	60		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Reticuline oxidase




- Molecule 1: Reticuline oxidase





- Molecule 1: Reticuline oxidase





● Molecule 1: Reticuline oxidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.82Å 175.44Å 195.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.38 – 2.20 48.78 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.9 (47.38-2.20) 61.9 (48.78-1.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.96 (at 1.50Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, $R_{free}$	0.220 , 0.242 0.220 , 0.242	Depositor DCC
$R_{free}$ test set	7007 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	33.7	Xtriage
Anisotropy	0.406	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	16476	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.00 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.0087e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 12P, NAG, FAD, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/4033	0.43	0/5465
1	B	0.25	0/4033	0.43	0/5465
1	C	0.25	0/4013	0.44	0/5438
1	D	0.25	0/4025	0.43	0/5454
All	All	0.25	0/16104	0.43	0/21822

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3935	0	3861	16	0
1	B	3935	0	3863	28	0
1	C	3916	0	3848	38	1
1	D	3927	0	3857	24	0
2	A	53	0	29	1	0
2	B	53	0	29	0	0
2	C	53	0	29	1	0
2	D	53	0	28	2	0
3	A	28	0	25	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	28	0	25	0	0
3	C	28	0	25	0	0
3	D	28	0	25	1	0
4	A	14	0	13	0	0
4	B	14	0	13	0	0
4	C	14	0	13	0	0
4	D	14	0	13	0	0
5	B	5	0	0	3	0
5	C	10	0	0	0	0
5	D	10	0	0	0	0
6	C	28	0	37	1	0
6	D	22	0	29	1	0
7	A	105	0	0	1	0
7	B	71	0	0	0	0
7	C	72	0	0	0	0
7	D	60	0	0	0	0
All	All	16476	0	15762	108	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:483:ARG:NH2	1:B:495:GLU:OE2	2.11	0.83
1:D:52:ARG:NH2	3:D:603:NAG:O7	2.19	0.76
1:D:65:ASN:O	1:D:67:LEU:N	2.25	0.69
1:D:424:GLN:O	1:D:426:GLU:N	2.24	0.68
1:C:287:GLU:O	1:C:289:GLN:N	2.25	0.68

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:495:GLU:OE1	1:C:499:ARG:NH1[3_556]	2.18	0.02

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	496/538 (92%)	475 (96%)	19 (4%)	2 (0%)	38	41
1	B	496/538 (92%)	468 (94%)	27 (5%)	1 (0%)	51	58
1	C	494/538 (92%)	469 (95%)	22 (4%)	3 (1%)	28	29
1	D	495/538 (92%)	467 (94%)	24 (5%)	4 (1%)	22	21
All	All	1981/2152 (92%)	1879 (95%)	92 (5%)	10 (0%)	32	34

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	68	ILE
1	B	288	LYS
1	C	288	LYS
1	D	66	SER
1	D	424	GLN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	426/460 (93%)	423 (99%)	3 (1%)	87	93
1	B	426/460 (93%)	421 (99%)	5 (1%)	75	86
1	C	424/460 (92%)	418 (99%)	6 (1%)	71	84
1	D	425/460 (92%)	421 (99%)	4 (1%)	82	91
All	All	1701/1840 (92%)	1683 (99%)	18 (1%)	78	88

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	25	ASN
1	C	273	LEU
1	D	289	GLN
1	B	425	SER
1	B	426	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	459	HIS
1	C	55	HIS
1	C	476	ASN
1	B	408	HIS
1	C	459	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	602	1,3	14,14,15	0.27	0	15,19,21	0.46	0
3	NAG	A	603	3	14,14,15	0.40	0	15,19,21	0.36	0
3	NAG	B	602	1,3	14,14,15	0.33	0	15,19,21	0.42	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	B	603	3	14,14,15	0.20	0	15,19,21	0.48	0
3	NAG	C	602	1,3	14,14,15	0.36	0	15,19,21	0.48	0
3	NAG	C	603	3	14,14,15	0.28	0	15,19,21	0.37	0
3	NAG	D	602	1,3	14,14,15	0.41	0	15,19,21	0.37	0
3	NAG	D	603	3	14,14,15	0.22	0	15,19,21	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	602	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	603	3	-	0/6/23/26	0/1/1/1
3	NAG	B	602	1,3	-	0/6/23/26	0/1/1/1
3	NAG	B	603	3	-	0/6/23/26	0/1/1/1
3	NAG	C	602	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	603	3	-	0/6/23/26	0/1/1/1
3	NAG	D	602	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	603	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	603	NAG	1	0

## 5.6 Ligand geometry

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected



value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FAD	A	601	1	51,58,58	1.39	5 (9%)	54,89,89	1.97	10 (18%)
4	NAG	A	604	1	14,14,15	0.20	0	15,19,21	0.50	0
2	FAD	B	601	1	51,58,58	1.37	5 (9%)	54,89,89	1.99	9 (16%)
4	NAG	B	604	1	14,14,15	0.29	0	15,19,21	0.38	0
5	SO4	B	605	-	4,4,4	0.14	0	6,6,6	0.07	0
2	FAD	C	601	1	51,58,58	1.36	5 (9%)	54,89,89	1.99	10 (18%)
4	NAG	C	604	1	14,14,15	0.40	0	15,19,21	0.40	0
6	12P	C	605	-	27,27,36	0.40	0	26,26,35	0.63	0
5	SO4	C	606	-	4,4,4	0.16	0	6,6,6	0.07	0
5	SO4	C	607	-	4,4,4	0.15	0	6,6,6	0.06	0
2	FAD	D	601	1	51,58,58	1.38	5 (9%)	54,89,89	2.02	10 (18%)
4	NAG	D	604	1	14,14,15	0.20	0	15,19,21	0.47	0
6	12P	D	605	-	21,21,36	0.42	0	20,20,35	0.48	0
5	SO4	D	606	-	4,4,4	0.15	0	6,6,6	0.07	0
5	SO4	D	607	-	4,4,4	0.14	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	601	1	-	0/28/50/50	0/6/6/6
4	NAG	A	604	1	-	0/6/23/26	0/1/1/1
2	FAD	B	601	1	-	0/28/50/50	0/6/6/6
4	NAG	B	604	1	-	0/6/23/26	0/1/1/1
5	SO4	B	605	-	-	0/0/0/0	0/0/0/0
2	FAD	C	601	1	-	0/28/50/50	0/6/6/6
4	NAG	C	604	1	-	0/6/23/26	0/1/1/1
6	12P	C	605	-	-	0/25/25/34	0/0/0/0
5	SO4	C	606	-	-	0/0/0/0	0/0/0/0
5	SO4	C	607	-	-	0/0/0/0	0/0/0/0
2	FAD	D	601	1	-	0/28/50/50	0/6/6/6
4	NAG	D	604	1	-	0/6/23/26	0/1/1/1
6	12P	D	605	-	-	0/19/19/34	0/0/0/0
5	SO4	D	606	-	-	0/0/0/0	0/0/0/0
5	SO4	D	607	-	-	0/0/0/0	0/0/0/0

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	FAD	C5A-C4A	2.99	1.47	1.40
2	D	601	FAD	C5A-C4A	3.08	1.47	1.40
2	A	601	FAD	C5A-C4A	3.09	1.47	1.40
2	C	601	FAD	C5A-C4A	3.14	1.47	1.40
2	D	601	FAD	C8-C7	3.21	1.49	1.41

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	FAD	N3A-C2A-N1A	-6.32	123.36	128.86
2	C	601	FAD	N3A-C2A-N1A	-6.04	123.60	128.86
2	D	601	FAD	N3A-C2A-N1A	-6.01	123.62	128.86
2	A	601	FAD	N3A-C2A-N1A	-5.81	123.80	128.86
2	D	601	FAD	C4-C4X-C10	-4.47	116.35	119.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	FAD	1	0
5	B	605	SO4	3	0
2	C	601	FAD	1	0
6	C	605	12P	1	0
2	D	601	FAD	2	0
6	D	605	12P	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	498/538 (92%)	-0.14	17 (3%) 46 43	28, 45, 76, 111	0
1	B	498/538 (92%)	0.16	23 (4%) 33 32	29, 56, 86, 114	0
1	C	496/538 (92%)	0.16	31 (6%) 21 19	30, 52, 80, 107	0
1	D	497/538 (92%)	0.06	28 (5%) 25 24	28, 50, 87, 114	0
All	All	1989/2152 (92%)	0.06	99 (4%) 30 28	28, 51, 84, 114	0

The worst 5 of 99 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	286	ASP	10.2
1	B	351	PHE	8.5
1	D	351	PHE	7.9
1	C	351	PHE	7.9
1	A	351	PHE	7.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	A	602	14/15	0.85	0.18	2.08	50,55,60,65	0
3	NAG	B	602	14/15	0.81	0.17	1.73	56,63,69,70	0
3	NAG	C	602	14/15	0.84	0.17	1.43	54,63,67,68	0
3	NAG	D	602	14/15	0.67	0.26	1.40	63,73,79,80	0
3	NAG	B	603	14/15	0.80	0.18	-	68,73,81,84	0
3	NAG	C	603	14/15	0.85	0.23	-	63,70,78,80	0
3	NAG	D	603	14/15	0.81	0.30	-	72,86,97,100	0
3	NAG	A	603	14/15	0.81	0.25	-	50,69,79,80	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	SO4	D	607	5/5	0.84	0.19	4.99	46,50,55,64	5
5	SO4	D	606	5/5	0.81	0.19	3.62	48,52,57,59	5
5	SO4	C	606	5/5	0.97	0.17	1.33	60,62,73,73	0
5	SO4	C	607	5/5	0.92	0.17	0.56	56,57,63,69	5
2	FAD	C	601	53/53	0.95	0.18	0.08	28,38,50,53	0
2	FAD	A	601	53/53	0.96	0.16	0.05	24,31,41,44	0
2	FAD	B	601	53/53	0.96	0.18	-0.03	28,36,45,49	0
6	12P	D	605	22/37	0.82	0.10	-0.25	39,51,56,59	0
6	12P	C	605	28/37	0.85	0.11	-0.26	36,50,60,68	0
2	FAD	D	601	53/53	0.96	0.14	-0.33	26,34,47,50	0
5	SO4	B	605	5/5	0.97	0.09	-0.56	60,66,72,73	0
4	NAG	C	604	14/15	0.58	0.43	-	83,93,99,103	0
4	NAG	A	604	14/15	0.80	0.36	-	78,82,87,87	0
4	NAG	D	604	14/15	0.74	0.40	-	90,95,96,98	0
4	NAG	B	604	14/15	0.74	0.46	-	93,97,102,106	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.