



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 08:11 am GMT

PDB ID : 1Q17  
Title : Structure of the yeast Hst2 protein deacetylase in ternary complex with 2'-O-acetyl ADP ribose and histone peptide  
Authors : Zhao, K.; Chai, X.; Marmorstein, R.  
Deposited on : 2003-07-18  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	<div> <div>2%</div> <div>69%</div> <div>24%</div> <div>• •</div> </div>
1	B	300	<div> <div>%</div> <div>64%</div> <div>29%</div> <div>• • •</div> </div>
1	C	300	<div> <div>3%</div> <div>64%</div> <div>31%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	A	703	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7261 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HST2 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	290	Total	C	N	O	S	0	0	0
			2301	1486	379	424	12			
1	B	294	Total	C	N	O	S	0	0	0
			2337	1508	388	429	12			
1	C	295	Total	C	N	O	S	0	0	0
			2346	1514	390	430	12			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	EXPRESSION TAG	UNP P53686
A	-4	HIS	-	EXPRESSION TAG	UNP P53686
A	-3	GLY	-	EXPRESSION TAG	UNP P53686
A	-2	MET	-	EXPRESSION TAG	UNP P53686
A	-1	ALA	-	EXPRESSION TAG	UNP P53686
A	0	SER	-	EXPRESSION TAG	UNP P53686
B	-5	HIS	-	EXPRESSION TAG	UNP P53686
B	-4	HIS	-	EXPRESSION TAG	UNP P53686
B	-3	GLY	-	EXPRESSION TAG	UNP P53686
B	-2	MET	-	EXPRESSION TAG	UNP P53686
B	-1	ALA	-	EXPRESSION TAG	UNP P53686
B	0	SER	-	EXPRESSION TAG	UNP P53686
C	-5	HIS	-	EXPRESSION TAG	UNP P53686
C	-4	HIS	-	EXPRESSION TAG	UNP P53686
C	-3	GLY	-	EXPRESSION TAG	UNP P53686
C	-2	MET	-	EXPRESSION TAG	UNP P53686
C	-1	ALA	-	EXPRESSION TAG	UNP P53686
C	0	SER	-	EXPRESSION TAG	UNP P53686


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Cl	0	0
			2	2		
3	A	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		

- Molecule 4 is ADENOSINE-5-DIPHOSPHORIBOSE (three-letter code: APR) (formula: C<sub>15</sub>H<sub>23</sub>N<sub>5</sub>O<sub>14</sub>P<sub>2</sub>).




- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	56	Total 56	O 56	0	0
5	B	51	Total 51	O 51	0	0
5	C	55	Total 55	O 55	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: HST2 protein




#### • Molecule 1: HST2 protein



#### • Molecule 1: HST2 protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	146.16Å 109.68Å 85.83Å 90.00° 104.90° 90.00°	Depositor
Resolution (Å)	28.88 – 2.70 28.88 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.0 (28.88-2.70) 97.0 (28.88-2.70)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.39 (at 2.68Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.186 , 0.247 0.191 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	48.0	Xtriage
Anisotropy	0.246	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 58.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7261	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: APR, ZN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/2360	0.61	0/3198
1	B	0.36	0/2399	0.60	0/3251
1	C	0.36	0/2408	0.60	0/3263
All	All	0.36	0/7167	0.60	0/9712

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2301	0	2284	61	0
1	B	2337	0	2314	84	0
1	C	2346	0	2326	79	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	1	0	0	2	0
3	B	2	0	0	1	0
3	C	1	0	0	0	0
4	A	36	0	18	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	36	0	17	0	0
4	C	36	0	19	1	0
5	A	56	0	0	3	0
5	B	51	0	0	2	0
5	C	55	0	0	1	0
All	All	7261	0	6978	217	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 217 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:162:HIS:HB2	1:B:163:PRO:HD3	1.34	1.02
1:C:104:GLN:HE22	1:C:127:LYS:H	1.06	1.01
1:C:162:HIS:HB2	1:C:163:PRO:HD3	1.45	0.99
1:A:104:GLN:HE22	1:A:127:LYS:H	1.01	0.99
1:B:104:GLN:HE21	1:B:130:LEU:HD12	1.34	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/300 (95%)	263 (92%)	18 (6%)	5 (2%)	11	27
1	B	290/300 (97%)	255 (88%)	28 (10%)	7 (2%)	7	17
1	C	291/300 (97%)	260 (89%)	26 (9%)	5 (2%)	11	27
All	All	867/900 (96%)	778 (90%)	72 (8%)	17 (2%)	9	22

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	165	LYS
1	B	165	LYS
1	B	172	VAL
1	C	257	ALA
1	A	164	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/264 (97%)	236 (92%)	19 (8%)	16	36
1	B	259/264 (98%)	243 (94%)	16 (6%)	21	46
1	C	260/264 (98%)	239 (92%)	21 (8%)	14	31
All	All	774/792 (98%)	718 (93%)	56 (7%)	17	39

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	122	ARG
1	B	264	LEU
1	C	272	GLU
1	B	153	GLN
1	B	204	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	140	HIS
1	B	248	ASN
1	C	268	GLN
1	B	142	HIS
1	B	215	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 7 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	APR	A	1001	-	34,39,39	3.83	11 (32%)	36,60,60	2.57	14 (38%)
4	APR	B	1002	-	34,39,39	3.82	10 (29%)	36,60,60	2.67	14 (38%)
4	APR	C	1003	-	34,39,39	3.77	12 (35%)	36,60,60	2.61	13 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	APR	A	1001	-	-	0/18/54/54	0/4/4/4
4	APR	B	1002	-	-	0/18/54/54	0/4/4/4
4	APR	C	1003	-	-	0/18/54/54	0/4/4/4

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1001	APR	C1D-C2D	-14.12	1.37	1.52
4	B	1002	APR	C1D-C2D	-13.73	1.37	1.52
4	C	1003	APR	O2D-C2D	-13.15	1.12	1.43
4	B	1002	APR	O2D-C2D	-11.97	1.15	1.43
4	C	1003	APR	C1D-C2D	-11.90	1.39	1.52

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1002	APR	O1D-C1D-O4D	-6.22	102.78	111.14
4	C	1003	APR	O5D-PB-O1B	-5.21	88.25	109.25
4	A	1001	APR	O5D-PB-O1B	-4.79	89.93	109.25
4	C	1003	APR	O1D-C1D-O4D	-4.67	104.87	111.14
4	B	1002	APR	O5D-PB-O1B	-4.62	90.63	109.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1003	APR	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	290/300 (96%)	-0.38	6 (2%) 64 65	23, 44, 75, 120	0
1	B	294/300 (98%)	-0.21	4 (1%) 75 76	26, 47, 84, 145	0
1	C	295/300 (98%)	-0.31	8 (2%) 55 55	26, 47, 88, 127	0
All	All	879/900 (97%)	-0.30	18 (2%) 65 66	23, 46, 85, 145	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	162	HIS	4.8
1	A	162	HIS	3.8
1	A	165	LYS	3.8
1	B	165	LYS	3.5
1	A	294	GLN	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	APR	C	1003	36/36	0.97	0.14	0.18	25,36,45,49	0
4	APR	B	1002	36/36	0.97	0.15	0.06	22,36,48,50	0
4	APR	A	1001	36/36	0.97	0.14	-0.08	24,35,48,57	0
2	ZN	A	401	1/1	0.99	0.08	-0.89	53,53,53,53	0
2	ZN	C	401	1/1	1.00	0.10	-1.04	63,63,63,63	0
2	ZN	B	401	1/1	0.99	0.05	-3.25	61,61,61,61	0
3	CL	A	703	1/1	0.98	0.12	-	46,46,46,46	0
3	CL	B	702	1/1	0.98	0.17	-	48,48,48,48	0
3	CL	C	701	1/1	0.96	0.13	-	56,56,56,56	0
3	CL	B	704	1/1	0.97	0.17	-	43,43,43,43	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.