



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2017 – 06:20 AM EDT

PDB ID : 3Q5F
Title : Crystal structure of the Salmonella transcriptional regulator SlyA in complex with DNA
Authors : Dolan, K.T.; Duguid, E.M.
Deposited on : unknown
Resolution : 2.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2395 (3.00-2.92)
Clashscore	112137	2773 (3.00-2.92)
Ramachandran outliers	110173	2680 (3.00-2.92)
Sidechain outliers	110143	2683 (3.00-2.92)
RSRZ outliers	101464	2421 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	147	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 57%, yellow 37%, grey 6%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 100% 57% 37% • • </div> </div>
1	B	147	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 59%, yellow 34%, grey 6%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 100% 59% 34% • 5% </div> </div>
2	C	23	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 22%, yellow 43%, orange 35%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 22% 43% 35% </div> </div>
3	D	23	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 26%, yellow 52%, orange 22%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 26% 52% 22% </div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3186 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator slyA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	141	Total	C	N	O	S	0	0	0
			1129	711	206	209	3			
1	B	140	Total	C	N	O	S	0	0	0
			1120	706	205	206	3			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP P40676
A	-1	ASN	-	EXPRESSION TAG	UNP P40676
A	0	ALA	-	EXPRESSION TAG	UNP P40676
A	97	GLU	ASP	SEE REMARK 999	UNP P40676
A	98	PRO	ALA	SEE REMARK 999	UNP P40676
B	-2	SER	-	EXPRESSION TAG	UNP P40676
B	-1	ASN	-	EXPRESSION TAG	UNP P40676
B	0	ALA	-	EXPRESSION TAG	UNP P40676
B	97	GLU	ASP	SEE REMARK 999	UNP P40676
B	98	PRO	ALA	SEE REMARK 999	UNP P40676

- Molecule 2 is a DNA chain called DNA (5'-D(*AP*AP*TP*AP*AP*CP*TP*TP*AP*GP*CP*AP*AP*GP*CP*TP*AP*AP*TP*TP*AP*TP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	23	Total	C	N	O	P	0	0	0
			469	227	88	132	22			


- Molecule 3 is a DNA chain called DNA (5'-D(*TP*TP*AP*TP*AP*AP*TP*TP*AP*GP*CP*TP*TP*GP*CP*TP*AP*AP*GP*TP*TP*AP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	23	Total	C	N	O	P	0	0	0
			468	228	78	140	22			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcriptional regulator slyA



- Molecule 1: Transcriptional regulator slyA



- Molecule 2: DNA (5'-D(*AP*AP*TP*AP*AP*CP*TP*TP*AP*GP*CP*AP*AP*GP*CP*TP*AP*AP*TP*TP*AP*TP*A)-3')



- Molecule 3: DNA (5'-D(*TP*TP*AP*TP*AP*AP*TP*TP*AP*GP*CP*TP*TP*GP*CP*TP*AP*AP*GP*TP*TP*AP*T)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	59.57Å 59.57Å 291.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.10 – 2.96 46.10 – 2.95	Depositor EDS
% Data completeness (in resolution range)	89.4 (46.10-2.96) 89.9 (46.10-2.95)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 2.96Å)	Xtriage
Refinement program	PHENIX 1.6.1 _357	Depositor
R, R_{free}	0.213 , 0.266 0.206 , 0.258	Depositor DCC
R_{free} test set	1057 reflections (10.02%)	DCC
Wilson B-factor (Å ²)	45.1	Xtriage
Anisotropy	1.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 34.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3186	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1145	0.56	0/1546
1	B	0.38	0/1136	0.56	0/1534
2	C	0.93	0/527	1.76	15/811 (1.8%)
3	D	0.87	0/523	1.91	21/806 (2.6%)
All	All	0.60	0/3331	1.17	36/4697 (0.8%)

There are no bond length outliers.

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	7	DT	O4'-C4'-C3'	-12.32	98.61	106.00
2	C	10	DG	O4'-C1'-N9	11.93	116.35	108.00
2	C	23	DA	O4'-C4'-C3'	-9.06	100.56	106.00
2	C	12	DA	O4'-C1'-N9	-8.36	102.15	108.00
2	C	22	DT	N3-C2-O2	-8.07	117.46	122.30
3	D	7	DT	C1'-O4'-C4'	-7.89	102.21	110.10
3	D	16	DT	C4'-C3'-C2'	-7.18	96.64	103.10
3	D	10	DG	O4'-C1'-N9	6.86	112.80	108.00
3	D	14	DG	O4'-C1'-N9	-6.86	103.20	108.00
3	D	17	DA	C4'-C3'-C2'	-6.76	97.01	103.10
2	C	11	DC	O4'-C1'-N1	6.50	112.55	108.00
2	C	6	DC	O4'-C4'-C3'	-6.43	101.93	104.50
3	D	7	DT	C5-C4-O4	-6.33	120.47	124.90
3	D	8	DT	N3-C4-O4	6.29	123.67	119.90
3	D	7	DT	O4'-C1'-N1	-6.26	103.62	108.00
3	D	7	DT	N3-C4-O4	6.20	123.62	119.90
3	D	14	DG	C4'-C3'-C2'	-6.20	97.52	103.10
2	C	14	DG	C4'-C3'-C2'	-6.12	97.60	103.10
2	C	13	DA	O4'-C1'-C2'	-6.07	101.04	105.90
2	C	10	DG	C3'-C2'-C1'	-5.83	95.50	102.50
3	D	17	DA	O4'-C4'-C3'	-5.82	102.17	104.50
3	D	19	DG	N1-C6-O6	5.79	123.37	119.90
3	D	18	DA	O4'-C1'-N9	5.63	111.94	108.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	8	DT	C5-C4-O4	-5.59	120.98	124.90
2	C	19	DT	P-O5'-C5'	-5.58	111.98	120.90
3	D	20	DT	O4'-C1'-N1	5.54	111.88	108.00
3	D	8	DT	O4'-C1'-N1	-5.54	104.12	108.00
3	D	22	DA	N1-C6-N6	5.48	121.89	118.60
2	C	19	DT	C4'-C3'-C2'	-5.45	98.20	103.10
2	C	7	DT	N3-C4-O4	5.27	123.06	119.90
3	D	18	DA	C3'-C2'-C1'	-5.26	96.19	102.50
3	D	17	DA	C3'-C2'-C1'	-5.18	96.28	102.50
2	C	20	DT	C4'-C3'-C2'	-5.16	98.45	103.10
2	C	8	DT	N3-C4-O4	5.06	122.94	119.90
2	C	20	DT	N3-C4-O4	5.05	122.93	119.90
3	D	2	DT	C5-C4-O4	-5.04	121.37	124.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1129	0	1183	52	0
1	B	1120	0	1177	49	0
2	C	469	0	262	23	0
3	D	468	0	266	12	0
All	All	3186	0	2888	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (113) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:15:DC:H2''	3:D:16:DT:H5'	1.38	1.06
2:C:23:DA:H5'	2:C:23:DA:C8	1.92	1.05
1:A:86:ARG:NH1	2:C:23:DA:O4'	2.08	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:22:DT:H2''	2:C:23:DA:C5'	2.13	0.79
3:D:15:DC:C2'	3:D:16:DT:H5'	2.12	0.78
1:A:3:SER:H	1:B:111:ARG:HH12	1.35	0.75
2:C:23:DA:H5'	2:C:23:DA:H8	1.49	0.69
1:A:2:GLU:HG2	1:B:111:ARG:NH1	2.08	0.69
1:B:105:GLU:OE1	1:B:105:GLU:HA	1.95	0.67
1:A:86:ARG:HD3	2:C:22:DT:O2	1.95	0.66
1:A:37:LEU:HD21	1:A:70:LEU:HD22	1.76	0.66
1:A:97:GLU:N	1:A:98:PRO:HD2	2.11	0.66
1:A:105:GLU:OE1	1:A:105:GLU:HA	1.96	0.65
1:B:97:GLU:N	1:B:98:PRO:HD2	2.12	0.65
1:A:50:ILE:HD12	1:A:60:GLN:HE22	1.61	0.64
1:B:50:ILE:HD12	1:B:60:GLN:HE22	1.63	0.62
1:B:99:LEU:O	1:B:103:MET:HB2	2.01	0.60
2:C:19:DT:H1'	2:C:20:DT:H5'	1.84	0.60
1:A:99:LEU:O	1:A:103:MET:HB2	2.01	0.60
1:A:48:SER:OG	1:A:51:GLN:HG3	2.03	0.59
1:A:2:GLU:HG2	1:A:3:SER:H	1.68	0.59
1:B:25:LYS:N	1:B:26:PRO:HD2	2.18	0.58
1:B:48:SER:OG	1:B:51:GLN:HG3	2.03	0.58
1:B:67:LEU:HB2	1:B:78:ARG:HH21	1.69	0.58
1:A:25:LYS:N	1:A:26:PRO:HD2	2.18	0.58
1:A:67:LEU:HB2	1:A:78:ARG:HH21	1.69	0.57
1:A:125:LEU:CD2	1:B:125:LEU:HD21	2.35	0.57
1:A:3:SER:O	1:B:111:ARG:NH1	2.38	0.56
1:B:37:LEU:HD21	1:B:70:LEU:HD22	1.85	0.56
1:B:124:GLU:O	1:B:128:LYS:HG3	2.06	0.56
1:B:126:LEU:O	1:B:130:ILE:HG12	2.06	0.56
2:C:15:DC:H2'	2:C:16:DT:C6	2.41	0.56
1:A:34:TRP:HD1	1:A:103:MET:CE	2.19	0.55
1:B:86:ARG:HH12	3:D:23:DT:H4'	1.73	0.54
2:C:23:DA:C2	3:D:3:DA:C2	2.96	0.54
1:A:125:LEU:HD21	1:B:125:LEU:CD2	2.39	0.53
2:C:14:DG:H2''	2:C:15:DC:H5'	1.91	0.53
3:D:6:DA:H2''	3:D:7:DT:O5'	2.09	0.53
1:B:105:GLU:O	1:B:106:VAL:C	2.48	0.52
1:B:30:THR:HG1	1:B:33:HIS:CG	2.28	0.52
2:C:22:DT:C2'	2:C:23:DA:C5'	2.87	0.52
1:A:95:LYS:O	1:A:95:LYS:HG2	2.09	0.52
3:D:15:DC:H2'	3:D:16:DT:C6	2.45	0.52
1:B:34:TRP:HD1	1:B:103:MET:CE	2.22	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:22:DT:H1'	2:C:23:DA:H5''	1.91	0.51
1:B:86:ARG:NH1	3:D:23:DT:C4'	2.74	0.51
1:B:86:ARG:NH1	3:D:23:DT:H4'	2.25	0.51
1:B:95:LYS:HG2	1:B:95:LYS:O	2.09	0.51
1:A:124:GLU:O	1:A:128:LYS:HG3	2.11	0.51
3:D:9:DA:O5'	3:D:9:DA:H2'	2.11	0.51
1:A:126:LEU:O	1:A:130:ILE:HG12	2.11	0.50
2:C:18:DA:H2''	2:C:19:DT:H71	1.94	0.50
1:B:39:ASN:O	1:B:43:LEU:HG	2.12	0.50
1:A:125:LEU:HD23	1:B:125:LEU:HD21	1.93	0.49
1:A:30:THR:HG1	1:A:33:HIS:CG	2.30	0.49
1:A:2:GLU:HG2	1:B:111:ARG:HH12	1.73	0.49
2:C:22:DT:C2'	2:C:23:DA:H5''	2.43	0.49
1:A:137:ILE:HG21	1:B:15:ILE:HG13	1.96	0.48
1:A:84:ASP:O	1:A:86:ARG:N	2.47	0.48
2:C:14:DG:C2'	2:C:15:DC:H5'	2.44	0.48
1:A:105:GLU:O	1:A:106:VAL:C	2.52	0.47
1:B:84:ASP:O	1:B:86:ARG:N	2.48	0.47
1:A:23:ARG:NH2	1:B:140:LEU:HD22	2.30	0.47
1:A:132:LYS:O	1:A:135:HIS:HB3	2.14	0.47
2:C:1:DA:H1'	2:C:2:DA:H5'	1.96	0.47
2:C:6:DC:H2''	2:C:7:DT:O5'	2.14	0.47
1:A:46:ASP:CG	1:A:89:ARG:HD3	2.35	0.47
1:B:46:ASP:CG	1:B:89:ARG:HD3	2.35	0.47
1:A:125:LEU:HD21	1:B:125:LEU:HD23	1.97	0.46
1:A:39:ASN:O	1:A:43:LEU:HG	2.16	0.46
2:C:2:DA:H2''	2:C:3:DT:OP2	2.14	0.46
3:D:13:DT:H1'	3:D:14:DG:C8	2.51	0.46
1:A:23:ARG:HG3	1:A:23:ARG:HH11	1.81	0.46
1:A:84:ASP:C	1:A:86:ARG:H	2.20	0.45
1:A:3:SER:HB3	1:B:111:ARG:HH22	1.81	0.45
1:A:3:SER:N	1:B:111:ARG:HH12	2.09	0.45
3:D:16:DT:H1'	3:D:17:DA:H5'	1.97	0.45
1:A:23:ARG:HH22	1:B:140:LEU:HD22	1.82	0.45
2:C:15:DC:H2'	2:C:16:DT:H71	1.99	0.44
1:A:40:ILE:C	1:A:42:GLN:H	2.20	0.44
1:B:24:LEU:C	1:B:26:PRO:HD2	2.37	0.44
3:D:7:DT:H2'	3:D:8:DT:C6	2.52	0.44
1:A:23:ARG:HG3	1:A:23:ARG:NH1	2.32	0.44
1:B:132:LYS:O	1:B:135:HIS:HB3	2.18	0.44
1:B:34:TRP:HD1	1:B:103:MET:HE3	1.83	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82:ALA:H	1:A:89:ARG:NH2	2.16	0.44
1:B:23:ARG:O	1:B:106:VAL:HG11	2.18	0.44
1:A:86:ARG:NH1	2:C:23:DA:C4'	2.81	0.44
1:A:59:GLU:HG3	2:C:16:DT:OP2	2.17	0.44
1:A:8:ASP:CG	1:A:11:ARG:HH22	2.22	0.43
2:C:19:DT:H5'	2:C:19:DT:H6	1.82	0.43
1:A:97:GLU:N	1:A:98:PRO:CD	2.81	0.43
1:B:23:ARG:HG3	1:B:23:ARG:HH11	1.83	0.43
1:B:41:HIS:O	1:B:41:HIS:CG	2.71	0.43
2:C:22:DT:H2''	2:C:23:DA:H5'	1.94	0.43
1:B:40:ILE:C	1:B:42:GLN:H	2.21	0.43
1:B:23:ARG:HG3	1:B:23:ARG:NH1	2.33	0.43
1:B:84:ASP:C	1:B:86:ARG:H	2.21	0.43
1:A:95:LYS:O	1:A:98:PRO:HG2	2.18	0.43
1:A:41:HIS:O	1:A:41:HIS:CG	2.71	0.43
1:B:30:THR:OG1	1:B:32:THR:HG22	2.19	0.42
1:B:82:ALA:H	1:B:89:ARG:NH2	2.17	0.42
1:A:24:LEU:C	1:A:26:PRO:HD2	2.39	0.42
1:A:134:GLU:CD	1:B:11:ARG:HE	2.23	0.42
1:A:23:ARG:O	1:A:106:VAL:HG11	2.20	0.42
1:A:69:GLN:O	1:A:73:LYS:HG3	2.20	0.42
2:C:7:DT:H2''	2:C:8:DT:O5'	2.20	0.42
1:B:97:GLU:N	1:B:98:PRO:CD	2.82	0.41
1:A:125:LEU:CD2	1:B:125:LEU:CD2	2.97	0.41
1:A:27:LEU:O	1:A:28:GLU:HB2	2.21	0.41
1:B:8:ASP:CG	1:B:11:ARG:HH22	2.24	0.41
1:A:34:TRP:CD1	1:A:103:MET:CE	3.03	0.40
1:B:69:GLN:O	1:B:73:LYS:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/147 (95%)	120 (86%)	16 (12%)	3 (2%)	8	33
1	B	138/147 (94%)	121 (88%)	14 (10%)	3 (2%)	8	33
All	All	277/294 (94%)	241 (87%)	30 (11%)	6 (2%)	8	33

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	85	ARG
1	A	120	SER
1	B	85	ARG
1	B	120	SER
1	A	41	HIS
1	B	41	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/131 (96%)	120 (95%)	6 (5%)	30	66
1	B	125/131 (95%)	119 (95%)	6 (5%)	30	66
All	All	251/262 (96%)	239 (95%)	12 (5%)	30	66

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	SER
1	A	17	ARG
1	A	32	THR
1	A	81	CYS
1	A	85	ARG
1	A	142	SER
1	B	7	SER
1	B	17	ARG
1	B	32	THR
1	B	81	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	85	ARG
1	B	142	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	GLN
1	B	49	GLN
1	B	60	GLN
1	B	136	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	141/147 (95%)	0.22	1 (0%) 87 74	41, 70, 105, 122	0
1	B	140/147 (95%)	0.18	2 (1%) 75 58	40, 71, 105, 124	0
2	C	23/23 (100%)	-0.28	0 100 100	43, 54, 64, 68	0
3	D	23/23 (100%)	-0.23	0 100 100	44, 53, 62, 65	0
All	All	327/340 (96%)	0.14	3 (0%) 84 69	40, 68, 104, 124	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	102	GLU	2.4
1	A	99	LEU	2.2
1	B	98	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.