



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 09:36 pm GMT

PDB ID : 4Q7U
Title : Crystal structure of photoswitchable fluorescent protein PSmOrange2
Authors : Malashkevich, V.N.; Pletnev, S.; Almo, S.C.
Deposited on : 2014-04-25
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1131 (1.32-1.28)
Clashscore	112137	1185 (1.32-1.28)
Ramachandran outliers	110173	1138 (1.32-1.28)
Sidechain outliers	110143	1138 (1.32-1.28)
RSRZ outliers	101464	1133 (1.32-1.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	<div> <div>4%</div> <div>74%</div> <div>14%</div> <div>11%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	500	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2141 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PSmOrange2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	218	Total	C	N	O	S	0	26	0
			1924	1239	314	362	9			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-16	MET	-	EXPRESSION TAG	UNP D0VWW2
A	-15	GLY	-	EXPRESSION TAG	UNP D0VWW2
A	-14	SER	-	EXPRESSION TAG	UNP D0VWW2
A	-13	HIS	-	EXPRESSION TAG	UNP D0VWW2
A	-12	HIS	-	EXPRESSION TAG	UNP D0VWW2
A	-11	HIS	-	EXPRESSION TAG	UNP D0VWW2
A	-10	HIS	-	EXPRESSION TAG	UNP D0VWW2
A	-9	HIS	-	EXPRESSION TAG	UNP D0VWW2
A	-8	HIS	-	EXPRESSION TAG	UNP D0VWW2
A	-7	GLY	-	EXPRESSION TAG	UNP D0VWW2
A	-6	ARG	-	EXPRESSION TAG	UNP D0VWW2
A	-5	SER	-	EXPRESSION TAG	UNP D0VWW2
A	17	HIS	ARG	ENGINEERED MUTATION	UNP D0VWW2
A	21	THR	SER	ENGINEERED MUTATION	UNP D0VWW2
A	36	HIS	ARG	ENGINEERED MUTATION	UNP D0VWW2
A	64	LEU	GLN	ENGINEERED MUTATION	UNP D0VWW2
A	66	OIM	PHE	CHROMOPHORE	UNP D0VWW2
A	66	OIM	GLY	CHROMOPHORE	UNP D0VWW2
A	66	OIM	TYR	CHROMOPHORE	UNP D0VWW2
A	66	OIM	GLY	CHROMOPHORE	UNP D0VWW2
A	99	TYR	PHE	ENGINEERED MUTATION	UNP D0VWW2
A	124	MET	LEU	ENGINEERED MUTATION	UNP D0VWW2
A	162	ARG	LYS	ENGINEERED MUTATION	UNP D0VWW2
A	186	SER	PRO	ENGINEERED MUTATION	UNP D0VWW2
A	188	LEU	GLN	ENGINEERED MUTATION	UNP D0VWW2
A	217	SER	ALA	ENGINEERED MUTATION	UNP D0VWW2
A	219	ALA	GLY	ENGINEERED MUTATION	UNP D0VWW2

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	211	Total	O	0	0
			211	211		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PSmOrange2

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.72Å 43.87Å 52.54Å 90.00° 94.37° 90.00°	Depositor
Resolution (Å)	52.39 – 1.30 33.16 – 1.30	Depositor EDS
% Data completeness (in resolution range)	96.3 (52.39-1.30) 96.3 (33.16-1.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 1.30Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.148 , 0.177 0.147 , 0.176	Depositor DCC
R_{free} test set	2525 reflections (5.40%)	DCC
Wilson B-factor (Å ²)	10.9	Xtriage
Anisotropy	0.225	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2141	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, OIM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	2/1995 (0.1%)	0.65	0/2681

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	215[A]	GLU	CD-OE2	-6.94	1.18	1.25
1	A	215[B]	GLU	CD-OE2	-6.94	1.18	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1924	0	1912	30	2
2	A	6	0	8	0	0
3	A	211	0	0	3	1
All	All	2141	0	1920	30	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:164[B]:ARG:HH11	1:A:172[B]:HIS:CD2	1.52	1.28
1:A:164[B]:ARG:NH1	1:A:172[B]:HIS:CD2	2.29	0.99
1:A:164[B]:ARG:NH1	1:A:172[B]:HIS:NE2	2.17	0.91
1:A:215[A]:GLU:OE1	3:A:673:HOH:O	1.93	0.87
1:A:164[B]:ARG:HH11	1:A:172[B]:HIS:CG	1.94	0.85
1:A:164[B]:ARG:HH11	1:A:172[B]:HIS:CE1	1.99	0.80
1:A:151[B]:TYR:OH	3:A:803:HOH:O	2.02	0.76
1:A:139:LYS:HA	1:A:168[A]:LYS:HD2	1.67	0.73
1:A:40:GLY:HA2	1:A:72[B]:TYR:O	1.90	0.71
1:A:164[B]:ARG:NH1	1:A:172[B]:HIS:CE1	2.57	0.71
1:A:174[A]:THR:HG23	3:A:788:HOH:O	1.94	0.67
1:A:141:MET:SD	1:A:168[A]:LYS:CG	2.84	0.66
1:A:141:MET:SD	1:A:168[A]:LYS:HG2	2.37	0.64
1:A:149:ARG:HG2	1:A:194[B]:ILE:CD1	2.27	0.64
1:A:162:ARG:CZ	1:A:176:GLU:OE1	2.51	0.59
1:A:96[A]:VAL:HG22	1:A:106[A]:THR:HG22	1.84	0.58
1:A:19[A]:GLU:HG2	1:A:28:GLU:HG2	1.88	0.56
1:A:21[A]:THR:HG22	1:A:26:GLU:HG2	1.87	0.55
1:A:141:MET:SD	1:A:168[A]:LYS:HG3	2.47	0.53
1:A:164[A]:ARG:HG2	1:A:174[A]:THR:HG22	1.89	0.53
1:A:149:ARG:HG2	1:A:194[B]:ILE:HD11	1.90	0.52
1:A:78[B]:ASP:OD1	1:A:221:HIS:NE2	2.42	0.51
1:A:53:PRO:HB3	1:A:207:ASP:HB2	1.93	0.49
1:A:47:LYS:HG2	1:A:210[A]:ILE:HD13	1.95	0.49
1:A:162:ARG:NH2	1:A:176:GLU:OE1	2.47	0.48
1:A:151[B]:TYR:HE2	1:A:160:GLU:HG3	1.82	0.43
1:A:164[B]:ARG:HH11	1:A:172[B]:HIS:NE2	1.57	0.42
1:A:47:LYS:HG2	1:A:210[A]:ILE:CD1	2.50	0.41
1:A:62[B]:SER:OG	1:A:201:ILE:HD11	2.21	0.40
1:A:15:LYS:CD	1:A:32:GLU:HG2	2.51	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13[B]:ARG:NH1	1:A:169:ASP:O[1_545]	2.01	0.19
1:A:166[B]:LYS:NZ	3:A:644:HOH:O[2_656]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/245 (98%)	236 (99%)	2 (1%)	1 (0%)	38	11

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/208 (101%)	207 (99%)	3 (1%)	71	34

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	154	ASP
1	A	215[A]	GLU
1	A	215[B]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	GLN
1	A	128	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	137	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	OIM	A	66	1	28,32,33	5.19	7 (25%)	29,47,49	2.84	9 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OIM	A	66	1	-	0/12/53/54	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	OIM	C0-N1	-21.35	1.23	1.44
1	A	66	OIM	O0-C0	-5.41	1.31	1.38
1	A	66	OIM	CA2-C2	-4.44	1.44	1.48
1	A	66	OIM	CB1-CA1	-4.35	1.38	1.53
1	A	66	OIM	OH-CZ	-2.01	1.32	1.37
1	A	66	OIM	CA1-N1	6.62	1.33	1.28
1	A	66	OIM	CB2-CA2	12.98	1.46	1.35

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	OIM	O2-C2-CA2	-8.84	126.21	130.97
1	A	66	OIM	CG1-CB1-CA1	-6.17	100.93	114.70
1	A	66	OIM	O0-C0-N1	-5.60	99.39	110.13
1	A	66	OIM	O3-C3-CA3	-4.67	110.94	126.38
1	A	66	OIM	N3-C1-N2	-2.73	109.80	113.31
1	A	66	OIM	C2-CA2-N2	2.18	110.52	108.93
1	A	66	OIM	CD1-CG2-CD2	2.48	121.36	117.63
1	A	66	OIM	OG1-CB1-CG1	2.65	112.39	109.95
1	A	66	OIM	CA2-C2-N3	4.02	105.09	103.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	500	-	5,5,5	0.18	0	5,5,5	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	500	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/245 (88%)	0.10	10 (4%) 33 32	7, 11, 21, 51	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	224	GLY	11.1
1	A	225	GLY	8.2
1	A	6	ALA	5.7
1	A	154	ASP	5.0
1	A	155	GLY	4.0
1	A	7	ILE	3.6
1	A	223	THR	3.4
1	A	188	LEU	3.3
1	A	226	MET	2.7
1	A	162	ARG	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	OIM	A	66	30/31	0.88	0.12	-	11,14,20,24	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GOL	A	500	6/6	0.93	0.17	11.00	13,19,22,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.