



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 07:32 pm GMT

PDB ID : 1Q9G  
Title : NMR STRUCTURE OF THE OUTER MEMBRANE PROTEIN OMPX IN  
DHPC MICELLES  
Authors : Fernandez, C.; Hilty, C.; Wider, G.; Guntert, P.; Wuthrich, K.  
Deposited on : 2003-08-25

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : trunk28760  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*


The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	148	

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:93, A:101-A:148 (141)	1.34	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 2 single-model clusters were found.

Cluster number	Models
1	2, 5, 6, 9, 12, 13, 19
2	1, 3, 7, 10, 11, 16
3	4, 15, 17
4	14, 18
Single-model clusters	8; 20

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2223 atoms, of which 1066 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Outer membrane protein X.

Mol	Chain	Residues	Atoms						Trace
1	A	148	Total	C	H	N	O	S	0
			2223	731	1066	190	233	3	

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ASN	HIS	ENGINEERED	UNP P0A917

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Outer membrane protein X




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Outer membrane protein X




#### 4.2.2 Score per residue for model 2

- Molecule 1: Outer membrane protein X




### 4.2.3 Score per residue for model 3

- Molecule 1: Outer membrane protein X




### 4.2.4 Score per residue for model 4

- Molecule 1: Outer membrane protein X




### 4.2.5 Score per residue for model 5

- Molecule 1: Outer membrane protein X




### 4.2.6 Score per residue for model 6


- Molecule 1: Outer membrane protein X



### 4.2.7 Score per residue for model 7



- Molecule 1: Outer membrane protein X







#### 4.2.8 Score per residue for model 8

- Molecule 1: Outer membrane protein X





#### 4.2.9 Score per residue for model 9

- Molecule 1: Outer membrane protein X




#### 4.2.10 Score per residue for model 10

- Molecule 1: Outer membrane protein X




#### 4.2.11 Score per residue for model 11

- Molecule 1: Outer membrane protein X




#### 4.2.12 Score per residue for model 12

- Molecule 1: Outer membrane protein X




#### 4.2.13 Score per residue for model 13

- Molecule 1: Outer membrane protein X




#### 4.2.14 Score per residue for model 14

- Molecule 1: Outer membrane protein X



#### 4.2.15 Score per residue for model 15


- Molecule 1: Outer membrane protein X




#### 4.2.16 Score per residue for model 16

- Molecule 1: Outer membrane protein X



Chain A: 



#### 4.2.17 Score per residue for model 17

- Molecule 1: Outer membrane protein X


Chain A: 



#### 4.2.18 Score per residue for model 18


- Molecule 1: Outer membrane protein X


Chain A: 



#### 4.2.19 Score per residue for model 19

- Molecule 1: Outer membrane protein X

Chain A: 



#### 4.2.20 Score per residue for model 20

- Molecule 1: Outer membrane protein X

Chain A: 





## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *TORSION ANGLE DYNAMICS (CANDID)*.

Of the 400 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
OPALp	refinement	2.6
CYANA	structure solution	1.0

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.72±0.01	0±0/1120 (0.0±0.0%)	1.07±0.03	1±1/1516 (0.0±0.1%)
All	All	0.72	0/22400 (0.0%)	1.07	13/30320 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.0±1.6
All	All	0	59

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	147	ARG	NE-CZ-NH2	-6.85	116.88	120.30	3	2
1	A	72	ARG	NE-CZ-NH2	-6.65	116.97	120.30	3	2
1	A	29	ARG	NE-CZ-NH2	-6.50	117.05	120.30	15	2
1	A	62	TYR	CB-CG-CD2	-6.15	117.31	121.00	13	1
1	A	127	TYR	CB-CG-CD2	-5.22	117.87	121.00	11	1
1	A	135	VAL	CA-CB-CG2	5.19	118.69	110.90	6	1
1	A	105	TYR	CB-CG-CD2	-5.18	117.89	121.00	15	1
1	A	147	ARG	NE-CZ-NH1	5.16	122.88	120.30	3	1
1	A	127	TYR	CB-CG-CD1	-5.11	117.93	121.00	20	1
1	A	131	ARG	NE-CZ-NH2	-5.02	117.79	120.30	18	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	133	ARG	Sidechain	4
1	A	72	ARG	Sidechain	4
1	A	9	TYR	Sidechain	3
1	A	105	TYR	Sidechain,Peptide	3
1	A	30	TYR	Sidechain	3
1	A	29	ARG	Sidechain	3
1	A	50	ARG	Sidechain	2
1	A	62	TYR	Sidechain,Peptide	2
1	A	80	TYR	Sidechain	2
1	A	146	TYR	Sidechain	2
1	A	74	ASN	Peptide	2
1	A	147	ARG	Sidechain	2
1	A	28	TYR	Sidechain	2
1	A	121	VAL	Peptide	2
1	A	87	TYR	Sidechain	2
1	A	45	TYR	Sidechain	2
1	A	33	ASP	Peptide	1
1	A	48	LYS	Peptide	1
1	A	139	THR	Peptide	1
1	A	88	GLY	Peptide	1
1	A	77	ALA	Peptide	1
1	A	127	TYR	Sidechain	1
1	A	64	GLY	Peptide	1
1	A	111	ALA	Peptide	1
1	A	61	GLN	Peptide	1
1	A	56	ASP	Peptide	1
1	A	63	TYR	Sidechain	1
1	A	37	LEU	Peptide	1
1	A	124	ASP	Peptide	1
1	A	49	SER	Peptide	1
1	A	109	TYR	Sidechain	1
1	A	131	ARG	Sidechain	1
1	A	71	TYR	Peptide	1
1	A	82	VAL	Peptide	1
1	A	20	LYS	Peptide	1

## 6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1093	1009	1011	1±1
All	All	21860	20180	20220	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:80:TYR:HA	1:A:111:ALA:HB3	0.55	1.78	14	1
1:A:73:ILE:HG22	1:A:74:ASN:H	0.51	1.65	12	1
1:A:80:TYR:CD1	1:A:113:LEU:HD11	0.47	2.44	12	1
1:A:73:ILE:HD11	1:A:79:ILE:HD11	0.46	1.85	15	1
1:A:132:ILE:O	1:A:135:VAL:HG22	0.46	2.10	6	1
1:A:80:TYR:CD1	1:A:113:LEU:HD21	0.43	2.48	14	1
1:A:65:ILE:H	1:A:65:ILE:HD12	0.43	1.74	9	1
1:A:62:TYR:CD2	1:A:89:LYS:HE3	0.42	2.48	16	1
1:A:79:ILE:HG22	1:A:113:LEU:CD1	0.42	2.44	1	1
1:A:4:THR:H	1:A:29:ARG:HD2	0.42	1.75	12	1
1:A:73:ILE:HG21	1:A:77:ALA:HB3	0.41	1.93	8	1
1:A:46:THR:HG22	1:A:62:TYR:H	0.40	1.77	15	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/148 (94%)	114±3 (82±2%)	20±3 (15±2%)	5±2 (4±2%)	7	37
All	All	2780/2960 (94%)	2277 (82%)	405 (15%)	98 (4%)	7	37

All 35 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	15	GLN	8
1	A	51	THR	6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	111	ALA	6
1	A	52	ALA	5
1	A	81	GLY	5
1	A	55	GLY	4
1	A	74	ASN	4
1	A	90	PHE	4
1	A	133	ARG	4
1	A	53	SER	3
1	A	119	GLU	3
1	A	54	SER	3
1	A	73	ILE	3
1	A	76	TRP	3
1	A	16	GLY	3
1	A	75	ASP	3
1	A	101	ASP	3
1	A	22	GLY	2
1	A	120	ASN	2
1	A	33	ASP	2
1	A	38	GLY	2
1	A	103	SER	2
1	A	60	ASN	2
1	A	89	LYS	2
1	A	17	GLN	2
1	A	8	GLY	2
1	A	93	THR	2
1	A	21	MET	1
1	A	34	ASN	1
1	A	137	VAL	1
1	A	118	MET	1
1	A	36	PRO	1
1	A	136	ASP	1
1	A	18	MET	1
1	A	108	SER	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	111/118 (94%)	92±4 (83±4%)	19±4 (17±4%)	6	41
All	All	2220/2360 (94%)	1843 (83%)	377 (17%)	6	41

All 99 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	26	LEU	11
1	A	115	PHE	9
1	A	24	PHE	9
1	A	120	ASN	8
1	A	48	LYS	8
1	A	147	ARG	8
1	A	50	ARG	7
1	A	18	MET	7
1	A	27	LYS	7
1	A	9	TYR	7
1	A	114	GLN	6
1	A	34	ASN	6
1	A	91	GLN	6
1	A	63	TYR	6
1	A	29	ARG	6
1	A	124	ASP	6
1	A	60	ASN	6
1	A	140	TRP	6
1	A	71	TYR	6
1	A	109	TYR	6
1	A	144	VAL	5
1	A	19	ASN	5
1	A	116	ASN	5
1	A	21	MET	5
1	A	33	ASP	5
1	A	73	ILE	5
1	A	51	THR	5
1	A	93	THR	5
1	A	30	TYR	5
1	A	25	ASN	4
1	A	132	ILE	4
1	A	113	LEU	4
1	A	108	SER	4
1	A	101	ASP	4
1	A	134	SER	4
1	A	80	TYR	4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	40	ILE	4
1	A	47	GLU	4
1	A	3	SER	4
1	A	11	GLN	4
1	A	133	ARG	4
1	A	65	ILE	4
1	A	125	PHE	4
1	A	57	TYR	4
1	A	37	LEU	4
1	A	61	GLN	4
1	A	32	GLU	4
1	A	72	ARG	4
1	A	105	TYR	4
1	A	78	SER	4
1	A	28	TYR	4
1	A	87	TYR	3
1	A	62	TYR	3
1	A	74	ASN	3
1	A	43	PHE	3
1	A	79	ILE	3
1	A	6	THR	3
1	A	15	GLN	3
1	A	66	THR	3
1	A	46	THR	3
1	A	45	TYR	3
1	A	123	LEU	3
1	A	127	TYR	3
1	A	75	ASP	3
1	A	102	THR	3
1	A	118	MET	3
1	A	130	SER	3
1	A	59	LYS	3
1	A	76	TRP	3
1	A	49	SER	3
1	A	148	PHE	3
1	A	54	SER	2
1	A	35	SER	2
1	A	103	SER	2
1	A	89	LYS	2
1	A	90	PHE	2
1	A	83	VAL	2
1	A	119	GLU	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	146	TYR	2
1	A	139	THR	2
1	A	13	ASP	2
1	A	107	PHE	2
1	A	128	GLU	2
1	A	17	GLN	2
1	A	129	GLN	2
1	A	2	THR	2
1	A	104	ASP	1
1	A	56	ASP	1
1	A	39	VAL	1
1	A	136	ASP	1
1	A	44	THR	1
1	A	92	THR	1
1	A	53	SER	1
1	A	4	THR	1
1	A	82	VAL	1
1	A	141	ILE	1
1	A	20	LYS	1
1	A	131	ARG	1
1	A	42	SER	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided