



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 05:35 am GMT

PDB ID : 4QBA  
Title : Crystal structure of the effector-binding domain of S. aureus CcpE  
Authors : Liu, X.; Lan, L.; Yang, C.G.  
Deposited on : 2014-05-06  
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4744 (2.24-2.20)
Clashscore	112137	5509 (2.24-2.20)
Ramachandran outliers	110173	5427 (2.24-2.20)
Sidechain outliers	110143	5428 (2.24-2.20)
RSRZ outliers	101464	4776 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	<div> <div>5%</div> <div> <div></div> <div>82%</div> <div>16%</div> <div>.</div> </div> </div>
1	B	204	<div> <div>2%</div> <div> <div></div> <div>78%</div> <div>17%</div> <div>..</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3295 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LysR family regulatory protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	204	Total	C	N	O	S	Se	0	0	0
			1647	1050	271	315	2	9			
1	B	196	Total	C	N	O	S	Se	0	0	0
			1579	1006	259	304	2	8			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cl	0	0
			1	1		
2	A	1	Total	Cl	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	34	Total	O	0	0
			34	34		
3	B	33	Total	O	0	0
			33	33		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LysR family regulatory protein



- Molecule 1: LysR family regulatory protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.61Å 77.18Å 93.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.21 36.80 – 2.21	Depositor EDS
% Data completeness (in resolution range)	98.8 (30.00-2.21) 98.9 (36.80-2.21)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.15 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.202 , 0.243 0.215 , 0.243	Depositor DCC
$R_{free}$ test set	1371 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.8	Xtriage
Anisotropy	0.454	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 38.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.033 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3295	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/1673	0.63	0/2253
1	B	0.45	0/1604	0.64	0/2163
All	All	0.46	0/3277	0.63	0/4416

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1647	0	1613	17	0
1	B	1579	0	1547	26	0
2	A	1	0	0	1	0
2	B	1	0	0	1	0
3	A	34	0	0	0	0
3	B	33	0	0	0	0
All	All	3295	0	3160	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:165:ILE:HD11	1:B:234:MSE:HE2	1.39	1.01
1:B:163:TYR:HB2	1:B:234:MSE:HE1	1.52	0.91
1:B:163:TYR:HB2	1:B:234:MSE:CE	2.20	0.70
1:B:104:THR:HG21	1:B:232:ILE:HG12	1.78	0.65
1:B:109:VAL:HG11	1:B:279:MSE:HE2	1.80	0.64
1:A:109:VAL:HG11	1:A:279:MSE:HE2	1.82	0.62
1:B:229:LEU:O	1:B:234:MSE:HE3	2.05	0.56
1:B:229:LEU:HB2	1:B:234:MSE:HE3	1.90	0.54
1:A:97:CYS:SG	1:A:101:ILE:HG23	2.48	0.54
1:A:157:LEU:HD21	1:A:260:MSE:HE2	1.91	0.53
1:A:100:LEU:HB2	2:A:301:CL:CL	2.46	0.53
1:B:97:CYS:SG	1:B:101:ILE:HG23	2.49	0.53
1:A:90:ASN:HD22	1:A:90:ASN:C	2.13	0.52
1:A:123:GLN:HE22	1:B:205:HIS:HE1	1.57	0.52
1:B:101:ILE:HA	1:B:104:THR:HG22	1.92	0.52
1:A:112:LEU:HD23	1:A:278:LEU:HD11	1.94	0.49
1:B:112:LEU:HD23	1:B:278:LEU:HD11	1.94	0.49
1:B:106:LEU:HD22	1:B:110:LEU:HG	1.95	0.48
1:A:106:LEU:HD22	1:A:110:LEU:HG	1.96	0.48
1:B:105:LEU:HD11	1:B:282:PHE:CD2	2.49	0.47
1:B:100:LEU:HB2	2:B:301:CL:CL	2.51	0.47
1:B:157:LEU:HD21	1:B:260:MSE:HE2	1.98	0.45
1:B:192:ILE:HG23	1:B:228:ILE:HD12	1.97	0.45
1:A:222:SER:OG	1:A:224:VAL:HG23	2.16	0.45
1:B:222:SER:OG	1:B:224:VAL:HG23	2.17	0.45
1:A:192:ILE:HG23	1:A:228:ILE:HD12	1.98	0.45
1:B:229:LEU:O	1:B:234:MSE:CE	2.64	0.44
1:A:219:MSE:HE1	1:B:106:LEU:HD13	1.99	0.44
1:B:98:SER:HA	1:B:127:GLY:O	2.17	0.44
1:A:98:SER:HA	1:A:127:GLY:O	2.19	0.43
1:A:263:ASP:HB3	1:A:266:MSE:HE2	2.00	0.43
1:A:185:ASP:OD1	1:A:187:ILE:HB	2.19	0.42
1:B:104:THR:HG21	1:B:232:ILE:CG1	2.47	0.42
1:A:123:GLN:NE2	1:B:205:HIS:HE1	2.17	0.42
1:B:104:THR:HG21	1:B:232:ILE:CD1	2.50	0.41
1:B:206:ALA:HB1	1:B:209:THR:HG22	2.02	0.41
1:A:123:GLN:HE22	1:B:205:HIS:CE1	2.38	0.41
1:A:206:ALA:HB1	1:A:209:THR:HG22	2.03	0.41
1:B:195:TRP:CD1	1:B:247:VAL:HG13	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	202/204 (99%)	198 (98%)	3 (2%)	1 (0%)	32	34
1	B	194/204 (95%)	189 (97%)	4 (2%)	1 (0%)	32	34
All	All	396/408 (97%)	387 (98%)	7 (2%)	2 (0%)	32	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	169	ASN
1	A	169	ASN

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/179 (105%)	176 (94%)	12 (6%)	20	22
1	B	181/179 (101%)	173 (96%)	8 (4%)	33	39
All	All	369/358 (103%)	349 (95%)	20 (5%)	26	29

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	88	GLU
1	A	89	VAL
1	A	90	ASN
1	A	100	LEU

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	106	LEU
1	A	108	GLU
1	A	120	VAL
1	A	150	MSE
1	A	152	LEU
1	A	168	LYS
1	A	170	ARG
1	A	207	THR
1	B	89	VAL
1	B	104	THR
1	B	106	LEU
1	B	120	VAL
1	B	130	GLU
1	B	152	LEU
1	B	207	THR
1	B	266	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	ASN
1	A	116	GLN
1	A	119	ASN
1	A	123	GLN
1	A	169	ASN
1	A	197	ASN
1	B	116	GLN
1	B	119	ASN
1	B	123	GLN
1	B	156	HIS
1	B	197	ASN
1	B	205	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	195/204 (95%)	-0.01	11 (5%) 25 23	34, 51, 96, 118	0
1	B	188/204 (92%)	-0.21	4 (2%) 64 61	35, 50, 77, 100	0
All	All	383/408 (93%)	-0.11	15 (3%) 40 38	34, 50, 92, 118	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	79	PHE	5.4
1	A	83	GLN	4.8
1	A	201	GLU	4.0
1	A	84	ALA	3.9
1	A	172	ASP	3.8
1	B	88	GLU	3.5
1	A	252	GLU	3.1
1	B	201	GLU	3.0
1	B	87	GLY	2.9
1	B	172	ASP	2.8
1	A	251	ASN	2.4
1	A	85	HIS	2.4
1	A	193	LYS	2.3
1	A	80	ASP	2.3
1	A	171	ARG	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CL	A	301	1/1	1.00	0.13	-	38,38,38,38	0
2	CL	B	301	1/1	0.99	0.06	-	49,49,49,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.