



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 10:50 pm GMT

PDB ID : 4QDM  
Title : Crystal structure of N-terminal mutant (V1L) of an alkali thermostable GH10 xylanase from *Bacillus* sp. NG-27  
Authors : Mahanta, P.; Bhardwaj, A.; Reddy, V.S.; Ramakumar, S.  
Deposited on : 2014-05-14  
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	402	-	-	X	X
3	GOL	B	402	-	-	X	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6304 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alkaline thermostable endoxylanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	354	Total	C	N	O	S	0	4	0
			2919	1858	492	564	5			
1	B	354	Total	C	N	O	S	0	4	0
			2935	1865	497	568	5			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	EXPRESSION TAG	UNP O30700
A	1	LEU	VAL	ENGINEERED MUTATION	UNP O30700
B	0	MET	-	EXPRESSION TAG	UNP O30700
B	1	LEU	VAL	ENGINEERED MUTATION	UNP O30700

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		
4	A	1	Total	Na	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	213	Total	O	0	0
			213	213		
5	B	221	Total	O	0	0
			221	221		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alkaline thermostable endoxylanase



- Molecule 1: Alkaline thermostable endoxylanase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.88Å 76.58Å 176.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.19 – 1.96 40.16 – 1.96	Depositor EDS
% Data completeness (in resolution range)	97.1 (40.19-1.96) 97.1 (40.16-1.96)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.76 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.152 , 0.190 0.164 , 0.197	Depositor DCC
$R_{free}$ test set	2682 reflections (5.39%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.3	Xtriage
Anisotropy	0.430	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 54.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6304	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.00	1/3008 (0.0%)	0.94	7/4102 (0.2%)
1	B	0.98	3/3017 (0.1%)	0.94	12/4115 (0.3%)
All	All	0.99	4/6025 (0.1%)	0.94	19/8217 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	299	GLU	CD-OE1	-6.59	1.18	1.25
1	B	59	GLU	CG-CD	6.03	1.60	1.51
1	B	135	GLU	CD-OE2	-5.20	1.20	1.25
1	A	149	GLU	CD-OE1	5.04	1.31	1.25

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	242	ARG	NE-CZ-NH2	-7.39	116.61	120.30
1	A	14	ARG	NE-CZ-NH2	-7.22	116.69	120.30
1	A	344	ARG	NE-CZ-NH1	6.28	123.44	120.30
1	A	215	LEU	CB-CG-CD2	6.23	121.60	111.00
1	A	14	ARG	NE-CZ-NH1	6.19	123.39	120.30
1	A	183	ARG	NE-CZ-NH1	5.90	123.25	120.30
1	B	136	ARG	NE-CZ-NH1	5.85	123.22	120.30
1	B	59	GLU	OE1-CD-OE2	-5.78	116.36	123.30
1	B	340	ASP	CB-CG-OD1	5.73	123.46	118.30
1	A	115	ARG	NE-CZ-NH1	5.72	123.16	120.30
1	B	136	ARG	NE-CZ-NH2	-5.51	117.55	120.30
1	B	261	ASP	CB-CG-OD2	-5.50	113.35	118.30
1	A	69	ASP	CB-CG-OD2	-5.47	113.38	118.30
1	B	176	ARG	NE-CZ-NH1	5.43	123.02	120.30
1	B	340	ASP	CB-CG-OD2	-5.24	113.58	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	323	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	B	100	ASP	CB-CG-OD1	5.14	122.92	118.30
1	B	205	ARG	NE-CZ-NH2	-5.10	117.75	120.30
1	B	140	ASP	CB-CG-OD2	5.06	122.85	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2919	0	2748	18	0
1	B	2935	0	2741	20	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	6	0	7	10	0
3	B	6	0	7	8	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	213	0	0	1	0
5	B	221	0	0	4	0
All	All	6304	0	5503	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:96:TRP:HH2	3:B:402:GOL:H11	1.42	0.83
1:B:96:TRP:HH2	3:B:402:GOL:C1	2.02	0.72
1:A:33:ARG:CB	1:A:343:TYR:CE2	2.75	0.69
1:B:322:ASP:OD1	1:B:325:ARG:NH2	2.30	0.64
1:A:128:THR:CG2	3:A:402:GOL:H2	2.29	0.62
1:A:57:GLN:HE22	3:A:402:GOL:H32	1.66	0.60

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:ILE:HG22	1:A:222[A]:ILE:HD11	1.84	0.60
1:B:125:ARG:HA	3:B:402:GOL:H31	1.83	0.60
1:A:128:THR:HG21	3:A:402:GOL:H2	1.84	0.59
1:A:57:GLN:NE2	3:A:402:GOL:H32	2.21	0.55
1:A:96:TRP:HH2	3:A:402:GOL:C3	2.18	0.55
1:B:57:GLN:HE22	3:B:402:GOL:H11	1.72	0.54
1:B:96:TRP:CH2	3:B:402:GOL:H11	2.32	0.54
1:A:26[B]:GLU:OE1	1:A:28:HIS:HE1	1.92	0.53
1:B:131:LYS:HD2	1:B:131:LYS:C	2.29	0.52
1:A:322:ASP:OD1	1:A:325:ARG:NH2	2.44	0.49
1:A:96:TRP:HH2	3:A:402:GOL:H31	1.77	0.49
1:B:173:GLU:HB2	5:B:721:HOH:O	2.14	0.48
1:B:57:GLN:NE2	3:B:402:GOL:H11	2.28	0.48
1:A:276:TYR:CZ	1:A:341:PRO:HD3	2.48	0.48
1:B:126:LEU:CD1	5:B:630:HOH:O	2.61	0.47
1:A:96:TRP:HH2	3:A:402:GOL:H32	1.81	0.45
1:B:33:ARG:CB	1:B:343:TYR:CE2	3.00	0.45
1:B:328:ASN:HB2	1:B:331:VAL:O	2.17	0.44
1:A:57:GLN:HE22	3:A:402:GOL:C3	2.29	0.44
1:B:267:TRP:HH2	1:B:327:TYR:HH	1.64	0.44
1:A:328:ASN:HB2	1:A:331:VAL:O	2.19	0.43
1:B:34:GLN:HG3	1:B:343:TYR:OH	2.18	0.43
1:A:202:THR:HB	1:A:203:PRO:HA	2.00	0.43
1:A:96:TRP:CH2	3:A:402:GOL:H31	2.53	0.43
1:B:126:LEU:HD12	5:B:630:HOH:O	2.19	0.43
1:B:276:TYR:CZ	1:B:341:PRO:HD3	2.54	0.43
1:B:66:ASP:OD1	5:B:696:HOH:O	2.21	0.43
1:A:283:ARG:HD2	5:A:662:HOH:O	2.20	0.41
1:A:96:TRP:CH2	3:A:402:GOL:C3	3.03	0.41
1:B:96:TRP:CH2	3:B:402:GOL:C1	2.92	0.41
1:B:22:GLY:HA2	1:B:44:SER:O	2.21	0.40
1:B:57:GLN:HE22	3:B:402:GOL:C1	2.33	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	356/355 (100%)	352 (99%)	4 (1%)	0	100	100
1	B	356/355 (100%)	351 (99%)	5 (1%)	0	100	100
All	All	712/710 (100%)	703 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	309/307 (101%)	309 (100%)	0	100	100
1	B	309/307 (101%)	308 (100%)	1 (0%)	94	93
All	All	618/614 (101%)	617 (100%)	1 (0%)	94	94

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	131	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	A	402	-	5,5,5	1.99	1 (20%)	5,5,5	1.62	1 (20%)
3	GOL	B	402	-	5,5,5	1.45	1 (20%)	5,5,5	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	402	-	-	0/4/4/4	0/0/0/0
3	GOL	B	402	-	-	0/4/4/4	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	GOL	O3-C3	-3.08	1.29	1.42
3	B	402	GOL	O1-C1	-2.76	1.30	1.42

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	GOL	O2-C2-C1	2.84	122.23	108.84

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	GOL	10	0
3	B	402	GOL	8	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	354/355 (99%)	-0.20	4 (1%)	80 86	6, 12, 25, 54	1 (0%)
1	B	354/355 (99%)	-0.20	6 (1%)	70 79	6, 12, 28, 42	0
All	All	708/710 (99%)	-0.20	10 (1%)	75 83	6, 12, 27, 54	1 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	267	TRP	4.8
1	A	267	TRP	3.9
1	B	1	LEU	3.6
1	A	270	ARG	2.7
1	A	268	PRO	2.3
1	B	102	GLU	2.2
1	B	110	THR	2.1
1	B	116	GLN	2.1
1	B	113	ALA	2.1
1	A	271	PRO	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	402	6/6	0.89	0.25	7.56	17,19,22,34	0
3	GOL	A	402	6/6	0.88	0.18	2.80	14,18,20,31	0
4	NA	B	403	1/1	0.99	0.06	-3.58	12,12,12,12	0
4	NA	A	403	1/1	0.99	0.06	-6.82	13,13,13,13	0
2	MG	B	401	1/1	0.99	0.09	-	8,8,8,8	0
2	MG	A	401	1/1	0.99	0.07	-	5,5,5,5	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.