



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2017 – 01:10 AM EDT

PDB ID : 3QGM
Title : p-nitrophenyl phosphatase from *Archaeoglobus fulgidus*
Authors : Osipiuk, J.; Zheng, H.; Xu, X.; Savchenko, A.; Edwards, A.; Joachimiak, A.;
Midwest Center for Structural Genomics (MCSG)
Deposited on : unknown
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 13%, green 84%, grey 12%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 100% 84% 13% • </div> </div>
1	B	268	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 13%, green 84%, grey 12%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 100% 84% 13% • • </div> </div>
1	C	268	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 87%, yellow 10%, orange 2%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 87% 10% • • </div> </div>
1	D	268	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 16%, green 80%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 100% 80% 16% • • </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8792 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called p-nitrophenyl phosphatase (Pho2).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	261	Total	C	N	O	S	Se	0	10	0
			2066	1319	347	392	1	7			
1	B	261	Total	C	N	O	S	Se	0	14	0
			2098	1337	355	398	1	7			
1	C	262	Total	C	N	O	S	Se	0	14	0
			2102	1341	354	398	1	8			
1	D	262	Total	C	N	O	S	Se	0	7	0
			2067	1316	352	390	1	8			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP O29873
A	-1	ASN	-	EXPRESSION TAG	UNP O29873
A	0	ALA	-	EXPRESSION TAG	UNP O29873
B	-2	SER	-	EXPRESSION TAG	UNP O29873
B	-1	ASN	-	EXPRESSION TAG	UNP O29873
B	0	ALA	-	EXPRESSION TAG	UNP O29873
C	-2	SER	-	EXPRESSION TAG	UNP O29873
C	-1	ASN	-	EXPRESSION TAG	UNP O29873
C	0	ALA	-	EXPRESSION TAG	UNP O29873
D	-2	SER	-	EXPRESSION TAG	UNP O29873
D	-1	ASN	-	EXPRESSION TAG	UNP O29873
D	0	ALA	-	EXPRESSION TAG	UNP O29873

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Ca	0	0
			1	1		
2	C	1	Total	Ca	0	0
			1	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	92	Total	O	0	0
			92	92		
4	B	122	Total	O	0	0
			122	122		
4	C	123	Total	O	0	1
			123	123		
4	D	114	Total	O	0	0
			114	114		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: p-nitrophenyl phosphatase (Pho2)




• Molecule 1: p-nitrophenyl phosphatase (Pho2)



• Molecule 1: p-nitrophenyl phosphatase (Pho2)



• Molecule 1: p-nitrophenyl phosphatase (Pho2)





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.25Å 68.84Å 70.15Å 74.99° 87.33° 85.76°	Depositor
Resolution (Å)	37.20 – 2.00 37.16 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.0 (37.20-2.00) 88.1 (37.16-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.169 , 0.213 0.172 , 0.213	Depositor DCC
R_{free} test set	3161 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.243	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.009 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8792	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	1/2116 (0.0%)	0.84	3/2845 (0.1%)
1	B	0.86	2/2161 (0.1%)	0.84	4/2904 (0.1%)
1	C	0.86	0/2165	0.85	4/2909 (0.1%)
1	D	0.82	0/2106	0.87	5/2830 (0.2%)
All	All	0.84	3/8548 (0.0%)	0.85	16/11488 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	138	CYS	CB-SG	-5.50	1.72	1.81
1	B	100[A]	GLU	CG-CD	5.40	1.60	1.51
1	B	100[B]	GLU	CG-CD	5.40	1.60	1.51

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	11	ASP	CB-CG-OD1	7.45	125.00	118.30
1	C	124[A]	ARG	NE-CZ-NH2	7.31	123.96	120.30
1	C	124[B]	ARG	NE-CZ-NH2	7.31	123.96	120.30
1	A	11	ASP	CB-CG-OD2	-6.75	112.22	118.30
1	A	140	ARG	NE-CZ-NH2	-6.62	116.99	120.30
1	D	11	ASP	CB-CG-OD2	-6.16	112.75	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	11	ASP	CB-CG-OD1	6.14	123.83	118.30
1	C	140	ARG	NE-CZ-NH2	-5.85	117.38	120.30
1	D	140	ARG	NE-CZ-NH2	-5.41	117.60	120.30
1	B	140	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	B	140	ARG	NE-CZ-NH2	-5.33	117.64	120.30
1	D	49[A]	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	D	49[B]	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	C	52	ARG	NE-CZ-NH2	-5.08	117.76	120.30
1	B	11	ASP	CB-CG-OD1	5.04	122.84	118.30
1	B	83	ARG	NE-CZ-NH2	-5.02	117.79	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	95	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2066	0	2164	21	0
1	B	2098	0	2199	22	0
1	C	2102	0	2208	30	0
1	D	2067	0	2158	27	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	B	4	0	6	0	0
4	A	92	0	0	2	0
4	B	122	0	0	3	0
4	C	123	0	0	2	0
4	D	114	0	0	5	0
All	All	8792	0	8735	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:83[B]:ARG:HB3	1:C:83[B]:ARG:NH1	1.62	1.12
1:A:165:MSE:HE2	1:B:131[A]:MSE:HG2	1.39	0.99
1:C:83[B]:ARG:CB	1:C:83[B]:ARG:HH11	1.82	0.92
1:C:90[A]:VAL:HG23	1:C:109:ILE:HD13	1.56	0.88
1:C:83[B]:ARG:CB	1:C:83[B]:ARG:NH1	2.37	0.84
1:C:83[B]:ARG:HB3	1:C:83[B]:ARG:HH11	1.40	0.83
1:C:83[B]:ARG:HB3	1:C:83[B]:ARG:CZ	2.17	0.73
1:C:58:LEU:HD12	1:C:70:ILE:HD11	1.71	0.72
1:A:240[B]:GLN:HE22	1:A:244:ARG:HH11	1.38	0.70
1:C:75:TYR:CE1	1:C:79[A]:ARG:HD2	2.27	0.69
1:C:96[B]:GLU:OE1	1:C:96[B]:GLU:HA	1.93	0.68
1:D:241:MSE:HE3	4:D:292:HOH:O	1.97	0.64
1:D:229:LEU:HD13	1:D:235[A]:ARG:HG2	1.80	0.64
1:B:227:LEU:HD21	1:B:233:THR:HG21	1.81	0.63
1:C:83[B]:ARG:HB2	1:C:83[B]:ARG:HH11	1.63	0.62
1:A:131[A]:MSE:HG2	1:B:165:MSE:HE2	1.80	0.62
1:D:103:ARG:NH1	4:D:298:HOH:O	2.32	0.61
1:A:51:ARG:HD2	1:A:70[B]:ILE:HG23	1.83	0.60
1:D:235[A]:ARG:HG3	1:D:235[A]:ARG:HH11	1.67	0.59
1:A:153:PHE:HB3	1:A:160[B]:ILE:HG22	1.84	0.59
1:D:72:VAL:CG1	1:D:74:THR:OG1	2.50	0.59
1:D:25[B]:GLU:OE2	1:D:257:LYS:NZ	2.33	0.57
1:D:46:ASN:HD22	1:D:48:THR:H	1.52	0.56
1:C:83[B]:ARG:NH1	4:C:332:HOH:O	2.39	0.55
1:D:79:ARG:HE	1:D:192[A]:GLU:CG	2.19	0.55
1:A:54:LEU:CB	1:A:70[B]:ILE:HD11	2.36	0.55
1:A:5:LYS:NZ	1:A:224:GLU:OE1	2.34	0.55
1:A:54:LEU:HB2	1:A:70[B]:ILE:HD11	1.87	0.55
1:A:34[B]:LYS:NZ	1:A:69:GLU:OE2	2.36	0.54
1:C:257:LYS:NZ	1:C:261:GLU:OE2	2.39	0.54
1:B:90[B]:VAL:HG12	1:B:117:TYR:HB2	1.90	0.54
1:C:196:ILE:HD13	4:C:338:HOH:O	2.07	0.54
1:C:58:LEU:HD12	1:C:70:ILE:CD1	2.37	0.53
1:A:54:LEU:HD12	1:A:72:VAL:HG21	1.92	0.52
1:C:90[A]:VAL:CG2	1:C:109:ILE:HD13	2.35	0.52
1:B:19:SER:OG	1:B:20:VAL:N	2.41	0.52
1:A:52:ARG:HD3	4:A:329:HOH:O	2.10	0.50
1:A:27:VAL:HG13	1:A:63:LEU:HD21	1.92	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:254:ASN:HB2	1:B:258:ASP:OD2	2.11	0.49
1:D:155:ALA:HB3	1:D:158:GLY:O	2.13	0.48
1:C:28[A]:GLU:OE1	1:C:32[A]:LYS:NZ	2.47	0.48
1:B:212:ASP:HB3	1:B:245:HIS:CD2	2.49	0.47
1:B:245:HIS:HD2	4:B:383:HOH:O	1.98	0.47
1:A:153:PHE:CD1	1:A:154:PRO:HD2	2.48	0.47
1:B:219:LYS:HE3	1:B:247:LEU:HD22	1.97	0.47
1:B:248:LYS:NZ	4:B:308:HOH:O	2.47	0.46
1:D:110:VAL:HB	1:D:114:GLU:HG3	1.96	0.46
1:D:32[B]:LYS:HB3	1:D:260:VAL:HG11	1.96	0.46
1:D:4:ASP:OD2	1:D:38:LYS:NZ	2.49	0.46
1:A:96:GLU:HG3	4:A:334:HOH:O	2.14	0.46
1:D:183:GLY:O	1:D:186:SER:HB3	2.16	0.46
1:B:9:ILE:HG23	1:B:43:VAL:CG2	2.46	0.46
1:B:4:ASP:O	1:B:5:LYS:HD2	2.15	0.46
1:B:83:ARG:HD2	4:B:312:HOH:O	2.14	0.46
1:D:47:SER:HA	4:D:441:HOH:O	2.14	0.46
1:D:5:LYS:HE3	1:D:224:GLU:HG3	1.97	0.46
1:D:229:LEU:HD13	1:D:235[B]:ARG:HG3	1.97	0.46
1:D:222:GLY:HA2	4:D:296:HOH:O	2.16	0.45
1:C:187[A]:GLU:HB3	1:C:191:ARG:HD2	1.97	0.45
1:B:90[A]:VAL:HG13	1:B:119:VAL:HG23	1.98	0.45
1:C:90[B]:VAL:HG13	1:C:119:VAL:HG23	1.99	0.45
1:A:41:ILE:HD12	1:A:196:ILE:HG21	1.99	0.45
1:D:87:ASN:HA	1:D:106:GLY:O	2.17	0.44
1:D:22:PRO:O	1:D:24:PRO:HD3	2.16	0.44
1:C:75:TYR:CZ	1:C:79[A]:ARG:HD2	2.52	0.44
1:A:228:VAL:HA	1:A:253:PHE:O	2.18	0.44
1:A:30:VAL:HG12	1:A:34[A]:LYS:HE2	1.99	0.43
1:C:228:VAL:HA	1:C:253:PHE:O	2.19	0.43
1:C:90[A]:VAL:HG12	1:C:117:TYR:HB2	2.01	0.43
1:C:32[A]:LYS:HB3	1:C:260:VAL:HG11	2.01	0.43
1:A:240[B]:GLN:HE22	1:A:244:ARG:NH1	2.09	0.43
1:D:92:THR:HA	1:D:119:VAL:O	2.18	0.43
1:D:188:VAL:O	1:D:192[B]:GLU:HG2	2.18	0.43
1:C:96[B]:GLU:OE1	1:C:96[B]:GLU:CA	2.63	0.43
1:B:126:ILE:HD11	1:B:131[B]:MSE:SE	2.69	0.42
1:C:124[A]:ARG:CG	1:C:124[A]:ARG:HH21	2.32	0.42
1:D:260:VAL:O	1:D:263:LEU:HB2	2.19	0.42
1:A:169:ALA:HB1	1:B:165:MSE:HG2	2.01	0.42
1:D:254:ASN:HB2	1:D:258:ASP:OD2	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:27:VAL:HG13	1:C:63:LEU:HD21	2.01	0.42
1:D:79:ARG:HE	1:D:192[A]:GLU:HG3	1.84	0.42
1:A:218:GLY:O	1:A:221:ILE:HD12	2.20	0.41
1:C:124[A]:ARG:HG2	1:C:124[A]:ARG:HH21	1.85	0.41
1:A:58:LEU:HD12	1:A:70[A]:ILE:CD1	2.51	0.41
1:D:235[A]:ARG:HG3	1:D:235[A]:ARG:NH1	2.34	0.41
1:C:32[B]:LYS:HB3	1:C:260:VAL:HG11	2.03	0.41
1:B:9:ILE:HG23	1:B:43:VAL:HG21	2.03	0.40
1:B:32:LYS:HB3	1:B:260:VAL:HG11	2.02	0.40
1:C:169:ALA:HB1	1:D:165[A]:MSE:HG2	2.02	0.40
1:B:130:LEU:HA	1:B:130:LEU:HD23	1.94	0.40
1:B:124[B]:ARG:NH1	1:B:124[B]:ARG:HG2	2.36	0.40
1:B:18:LYS:HZ3	1:B:23:ILE:HD11	1.85	0.40
1:D:241:MSE:HE1	4:D:404:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/268 (100%)	263 (98%)	6 (2%)	0	100	100
1	B	273/268 (102%)	269 (98%)	4 (2%)	0	100	100
1	C	274/268 (102%)	270 (98%)	2 (1%)	2 (1%)	25	18
1	D	267/268 (100%)	262 (98%)	5 (2%)	0	100	100
All	All	1083/1072 (101%)	1064 (98%)	17 (2%)	2 (0%)	55	48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	96[A]	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	96[B]	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/213 (106%)	221 (98%)	5 (2%)	57	60
1	B	230/213 (108%)	223 (97%)	7 (3%)	46	46
1	C	231/213 (108%)	222 (96%)	9 (4%)	37	34
1	D	224/213 (105%)	208 (93%)	16 (7%)	17	11
All	All	911/852 (107%)	874 (96%)	37 (4%)	41	31

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ASP
1	A	35	GLU
1	A	36	LEU
1	A	64	GLU
1	A	144	TYR
1	B	79	ARG
1	B	130	LEU
1	B	144	TYR
1	B	148	ASN
1	B	157[A]	ASP
1	B	157[B]	ASP
1	B	235	ARG
1	C	5	LYS
1	C	13[A]	ASP
1	C	13[B]	ASP
1	C	32[A]	LYS
1	C	32[B]	LYS
1	C	52	ARG
1	C	124[A]	ARG
1	C	124[B]	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	144	TYR
1	D	2	MSE
1	D	13	ASP
1	D	18[A]	LYS
1	D	18[B]	LYS
1	D	46	ASN
1	D	49[A]	ARG
1	D	49[B]	ARG
1	D	52	ARG
1	D	55	LEU
1	D	60	SER
1	D	72	VAL
1	D	144	TYR
1	D	202	LYS
1	D	235[A]	ARG
1	D	235[B]	ARG
1	D	263	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	148	ASN
1	B	245	HIS
1	D	46	ASN
1	D	210	GLN
1	D	240	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	B	505	-	3,3,3	0.76	0	2,2,2	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	505	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	255/268 (95%)	-0.37	2 (0%) 86 85	22, 33, 49, 57	0
1	B	255/268 (95%)	-0.34	2 (0%) 86 85	19, 30, 47, 56	0
1	C	255/268 (95%)	-0.32	1 (0%) 92 92	20, 30, 48, 62	0
1	D	255/268 (95%)	-0.32	3 (1%) 79 78	21, 33, 48, 59	0
All	All	1020/1072 (95%)	-0.34	8 (0%) 86 85	19, 31, 49, 62	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	157	ASP	3.1
1	A	235	ARG	2.9
1	B	235	ARG	2.6
1	C	3	PRO	2.6
1	D	235[A]	ARG	2.4
1	B	260	VAL	2.4
1	D	156	GLU	2.3
1	A	3	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CA	D	502	1/1	0.99	0.09	0.36	24,24,24,24	0
2	CA	B	504	1/1	1.00	0.06	-1.50	23,23,23,23	0
2	CA	C	501	1/1	1.00	0.07	-1.63	23,23,23,23	0
2	CA	A	503	1/1	0.99	0.03	-3.88	26,26,26,26	0
3	EDO	B	505	4/4	0.81	0.13	-	46,47,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.