



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 26, 2018 – 06:41 PM EST

PDB ID : 4QI6
Title : Cellobiose dehydrogenase from *Myriococcum thermophilum*, MtCDH
Authors : Tan, T.C.; Gandini, R.; Sygmund, C.; Kittl, R.; Haltrich, D.; Ludwig, R.;
Hallberg, B.M.; Divne, C.
Deposited on : 2014-05-30
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1015 (3.22-3.18)
Clashscore	112137	1009 (3.20-3.20)
Ramachandran outliers	110173	1118 (3.22-3.18)
Sidechain outliers	110143	1117 (3.22-3.18)
RSRZ outliers	101464	1020 (3.22-3.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	807	<div> <div>10%</div> <div>83%</div> <div>16%</div> <div>.</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6262 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellobiose dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	800	Total	C	N	O	S	0	0	0
			6055	3807	1023	1201	24			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	PCA	-	SEE REMARK 999	UNP A9XK88

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O		0	0
			14	8	1	5			
4	A	1	Total	C	N	O		0	0
			14	8	1	5			
4	A	1	Total	C	N	O		0	0
			14	8	1	5			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cellobiose dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	156.15Å 156.15Å 85.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.35 – 3.20 46.35 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.35-3.20) 99.9 (46.35-3.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 3.19Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.3_1477)	Depositor
R, R_{free}	0.215 , 0.273 0.215 , 0.275	Depositor DCC
R_{free} test set	1792 reflections (10.00%)	DCC
Wilson B-factor (Å ²)	91.7	Xtriage
Anisotropy	0.142	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 81.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6262	wwPDB-VP
Average B, all atoms (Å ²)	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, PCA, NAG, FAD, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.22	0/6202	0.43	0/8469

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	163	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6055	0	5745	80	0
2	A	43	0	30	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	53	0	31	3	0
4	A	56	0	52	0	0
5	A	55	0	50	2	0
All	All	6262	0	5908	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:MET:HE1	1:A:173:LEU:H	1.52	0.73
1:A:443:VAL:HG12	1:A:453:VAL:HG12	1.73	0.70
1:A:403:PHE:HB3	1:A:582:PHE:HB3	1.76	0.67
1:A:471:LYS:O	1:A:472:VAL:HB	1.96	0.64
1:A:642:ASP:OD2	1:A:647:LYS:NZ	2.29	0.64
1:A:582:PHE:CD1	1:A:598:TRP:HB2	2.34	0.62
1:A:38:ALA:HB3	1:A:52:ILE:HB	1.80	0.62
1:A:34:THR:HB	1:A:56:LYS:HB3	1.83	0.59
1:A:176:HIS:NE2	2:A:901:HEM:NC	2.51	0.59
1:A:205:LYS:HE2	5:A:909:MAN:H2	1.86	0.58
1:A:582:PHE:CE1	1:A:598:TRP:HB2	2.39	0.58
1:A:491:ARG:HD3	1:A:636:LEU:HD13	1.86	0.57
1:A:391:THR:HB	1:A:394:ASN:HB2	1.85	0.57
1:A:582:PHE:CE1	1:A:598:TRP:CB	2.88	0.57
1:A:532:LEU:HD13	1:A:645:TYR:HB3	1.87	0.56
1:A:498:ASP:OD2	1:A:629:ARG:NH2	2.37	0.56
1:A:530:ASP:OD1	1:A:531:HIS:N	2.39	0.56
1:A:595:GLN:HG2	1:A:746:THR:HG23	1.87	0.56
1:A:33:PHE:HZ	1:A:68:ILE:HD11	1.71	0.55
1:A:747:THR:HG22	1:A:748:ASN:H	1.71	0.55
1:A:334:ASP:O	1:A:342:LYS:NZ	2.41	0.53
1:A:441:LYS:HE3	1:A:454:GLU:HG2	1.90	0.53
1:A:540:HIS:HB3	1:A:543:VAL:HG23	1.90	0.53
1:A:706:ASN:O	1:A:740:ILE:HA	2.10	0.52
1:A:326:PHE:CG	1:A:747:THR:HG21	2.45	0.51
1:A:573:GLN:NE2	1:A:577:ASN:OD1	2.45	0.50
1:A:701:HIS:HB3	1:A:749:PRO:HG3	1.93	0.50
1:A:274:PRO:HD2	1:A:277:LEU:HD12	1.94	0.50
1:A:146:THR:HG22	1:A:151:LEU:HD11	1.95	0.49
1:A:731:ASP:OD2	1:A:732:ASN:N	2.46	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:ARG:HB3	1:A:102:PRO:HB2	1.94	0.49
1:A:105:TYR:CZ	1:A:107:GLY:HA3	2.48	0.48
1:A:85:HIS:HB3	1:A:90:TYR:HE2	1.79	0.48
1:A:472:VAL:HG12	1:A:473:THR:N	2.26	0.48
1:A:273:GLY:HA2	1:A:284:ARG:HB3	1.95	0.48
1:A:267:ASN:O	1:A:427:LYS:NZ	2.46	0.47
1:A:444:ILE:HD13	1:A:466:ILE:HD13	1.97	0.46
1:A:94:ARG:HD3	1:A:104:VAL:HA	1.97	0.46
1:A:293:GLN:HG3	1:A:552:TRP:CE3	2.51	0.46
1:A:274:PRO:HG3	1:A:290:LEU:HD13	1.98	0.45
1:A:315:GLY:O	1:A:318:THR:OG1	2.26	0.45
1:A:478:LEU:HB2	1:A:735:VAL:HG22	1.98	0.45
1:A:85:HIS:HB3	1:A:90:TYR:CE2	2.52	0.45
1:A:451:THR:O	1:A:469:VAL:HG22	2.17	0.45
1:A:435:TRP:CH2	1:A:467:VAL:HG11	2.52	0.45
1:A:15:ILE:HG23	1:A:199:TRP:CE3	2.52	0.44
1:A:587:VAL:HG22	1:A:593:VAL:HG22	1.99	0.44
1:A:198:ASP:OD2	5:A:907:MAN:O6	2.34	0.44
1:A:30:GLN:H	1:A:178:ASN:HD21	1.64	0.44
1:A:748:ASN:HB3	3:A:902:FAD:O2	2.18	0.44
1:A:579:GLY:N	1:A:600:ALA:O	2.50	0.44
1:A:440:VAL:HG22	1:A:455:VAL:HG12	2.00	0.44
1:A:748:ASN:HB3	3:A:902:FAD:C2	2.47	0.44
1:A:259:GLU:HG3	1:A:261:GLY:H	1.83	0.43
1:A:536:THR:OG1	1:A:676:PHE:O	2.30	0.43
1:A:79:LEU:HD22	2:A:901:HEM:HAC	1.99	0.43
1:A:482:THR:HA	1:A:739:SER:HB3	1.99	0.43
1:A:232:TYR:HB2	1:A:255:VAL:HG22	2.00	0.43
1:A:557:GLU:O	1:A:561:ASN:ND2	2.50	0.42
1:A:581:MET:HE2	1:A:599:THR:HG23	2.00	0.42
1:A:328:PRO:HB3	1:A:333:TRP:HE1	1.85	0.42
1:A:531:HIS:CD2	1:A:621:GLY:HA2	2.54	0.42
1:A:741:PHE:HA	1:A:742:PRO:HD3	1.89	0.42
1:A:238:GLY:HA3	3:A:902:FAD:O5B	2.19	0.42
1:A:101:MET:HB2	1:A:101:MET:HE2	1.89	0.42
1:A:435:TRP:CZ3	1:A:467:VAL:HG11	2.55	0.42
1:A:113:GLN:HB2	1:A:205:LYS:HB2	2.01	0.42
1:A:54:TYR:HD1	1:A:126:ILE:HG13	1.84	0.42
1:A:74:MET:HB2	2:A:901:HEM:C1D	2.55	0.42
1:A:399:LYS:HA	1:A:399:LYS:HD3	1.84	0.41
1:A:337:PHE:HA	1:A:338:PRO:HD3	1.92	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:TRP:CZ2	1:A:173:LEU:HD23	2.56	0.41
1:A:64:GLY:N	1:A:161:ASP:OD1	2.49	0.41
1:A:536:THR:HG22	1:A:616:MET:O	2.21	0.41
1:A:694:PRO:O	1:A:698:ARG:HB3	2.21	0.41
1:A:450:ILE:O	1:A:732:ASN:ND2	2.53	0.40
1:A:663:GLN:O	1:A:667:GLN:HB3	2.21	0.40
1:A:75:THR:HG21	1:A:307:ASP:HA	2.02	0.40
1:A:409:MET:O	1:A:416:ASN:HB3	2.21	0.40
1:A:676:PHE:HA	1:A:677:PRO:HA	1.97	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	796/807 (99%)	766 (96%)	30 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	648/654 (99%)	643 (99%)	5 (1%)	85	95

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	TRP
1	A	129	CYS
1	A	204	THR
1	A	498	ASP
1	A	509	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	573	GLN
1	A	577	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	A	1	1	8,8,9	1.58	1 (12%)	9,10,12	1.94	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	PCA	CD-N	4.15	1.46	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	PCA	CA-N-CD	-2.78	104.04	113.58
1	A	1	PCA	OE-CD-CG	-2.17	122.87	126.86
1	A	1	PCA	CG-CD-N	2.29	114.85	108.33
1	A	1	PCA	CB-CA-N	2.74	111.17	103.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	901	1	28,50,50	3.29	11 (39%)	17,82,82	2.80	7 (41%)
3	FAD	A	902	-	51,58,58	1.30	6 (11%)	54,89,89	1.86	4 (7%)
4	NAG	A	903	1	14,14,15	0.18	0	15,19,21	0.43	0
4	NAG	A	904	1	14,14,15	0.23	0	15,19,21	0.45	0
4	NAG	A	905	1	14,14,15	0.33	0	15,19,21	0.45	0
4	NAG	A	906	1	14,14,15	0.33	0	15,19,21	0.42	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	MAN	A	907	1	11,11,12	0.72	0	13,15,17	1.10	2 (15%)
5	MAN	A	908	1	11,11,12	0.80	0	13,15,17	1.08	2 (15%)
5	MAN	A	909	1	11,11,12	0.81	0	13,15,17	1.05	2 (15%)
5	MAN	A	910	1	11,11,12	0.71	0	13,15,17	1.05	2 (15%)
5	MAN	A	911	1	11,11,12	0.81	0	13,15,17	1.09	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	901	1	-	0/6/54/54	0/0/8/8
3	FAD	A	902	-	-	0/28/50/50	0/6/6/6
4	NAG	A	903	1	-	0/6/23/26	0/1/1/1
4	NAG	A	904	1	-	0/6/23/26	0/1/1/1
4	NAG	A	905	1	-	0/6/23/26	0/1/1/1
4	NAG	A	906	1	-	0/6/23/26	0/1/1/1
5	MAN	A	907	1	-	0/2/19/22	0/1/1/1
5	MAN	A	908	1	-	0/2/19/22	0/1/1/1
5	MAN	A	909	1	-	0/2/19/22	0/1/1/1
5	MAN	A	910	1	-	0/2/19/22	0/1/1/1
5	MAN	A	911	1	-	0/2/19/22	0/1/1/1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	902	FAD	C1'-N10	2.07	1.50	1.48
2	A	901	HEM	C4A-CHB	2.10	1.45	1.40
2	A	901	HEM	C4B-CHC	2.24	1.46	1.40
3	A	902	FAD	C2A-N1A	2.49	1.38	1.33
2	A	901	HEM	C1D-CHD	2.50	1.46	1.40
2	A	901	HEM	C3D-C2D	2.85	1.46	1.37
3	A	902	FAD	C4-N3	3.06	1.38	1.33
3	A	902	FAD	C4X-N5	3.46	1.38	1.33
2	A	901	HEM	C2A-C3A	3.59	1.48	1.37
3	A	902	FAD	C10-N1	3.61	1.38	1.33
3	A	902	FAD	C2A-N3A	3.83	1.38	1.32
2	A	901	HEM	C3B-C2B	4.04	1.45	1.40
2	A	901	HEM	C1A-NA	4.97	1.46	1.36
2	A	901	HEM	C4A-NA	5.28	1.47	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	HEM	C3C-C2C	6.25	1.48	1.40
2	A	901	HEM	C4C-NC	8.31	1.46	1.36
2	A	901	HEM	C1C-NC	9.09	1.47	1.36

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	902	FAD	N3A-C2A-N1A	-9.94	120.20	128.86
2	A	901	HEM	C1D-C2D-C3D	-7.50	101.78	107.00
3	A	902	FAD	C4X-C4-N3	-3.15	119.00	123.48
2	A	901	HEM	CBD-CAD-C3D	-2.44	107.82	112.47
2	A	901	HEM	C4A-C3A-C2A	-2.41	105.32	107.00
5	A	911	MAN	O2-C2-C3	-2.32	105.62	110.17
5	A	910	MAN	O2-C2-C3	-2.26	105.73	110.17
5	A	908	MAN	O2-C2-C3	-2.24	105.77	110.17
5	A	907	MAN	O2-C2-C3	-2.23	105.80	110.17
5	A	909	MAN	O2-C2-C3	-2.04	106.16	110.17
2	A	901	HEM	CBA-CAA-C2A	-2.02	108.62	112.48
5	A	910	MAN	C1-O5-C5	2.10	115.07	112.17
5	A	911	MAN	C1-O5-C5	2.11	115.08	112.17
5	A	909	MAN	C1-O5-C5	2.16	115.14	112.17
5	A	907	MAN	C1-O5-C5	2.19	115.18	112.17
5	A	908	MAN	C1-O5-C5	2.33	115.37	112.17
2	A	901	HEM	CMC-C2C-C3C	2.50	129.53	124.89
2	A	901	HEM	CMB-C2B-C3B	2.86	130.19	124.89
3	A	902	FAD	C5X-C9A-N10	4.43	120.95	117.66
3	A	902	FAD	C4-N3-C2	5.19	119.70	115.16
2	A	901	HEM	C3B-C4B-NB	5.78	116.68	109.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	HEM	3	0
3	A	902	FAD	3	0
5	A	907	MAN	1	0
5	A	909	MAN	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	799/807 (99%)	0.56	81 (10%) 8 5	39, 76, 161, 258	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	132	CYS	6.5
1	A	114	VAL	5.5
1	A	113	GLN	5.4
1	A	90	TYR	5.3
1	A	198	ASP	5.1
1	A	89	VAL	5.1
1	A	129	CYS	4.7
1	A	91	THR	4.6
1	A	54	TYR	4.6
1	A	149	GLY	4.5
1	A	16	THR	4.4
1	A	17	PHE	4.3
1	A	200	ALA	4.3
1	A	278	GLU	4.3
1	A	146	THR	4.2
1	A	126	ILE	4.2
1	A	115	SER	4.0
1	A	148	GLY	4.0
1	A	276	TRP	3.9
1	A	39	LEU	3.9
1	A	127	PHE	3.9
1	A	111	ILE	3.8
1	A	165	PRO	3.7
1	A	51	PHE	3.7
1	A	277	LEU	3.7
1	A	37	VAL	3.5
1	A	204	THR	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	556	ILE	3.3
1	A	199	TRP	3.3
1	A	53	GLY	3.3
1	A	695	SER	3.3
1	A	52	ILE	3.2
1	A	87	ASP	3.2
1	A	31	GLY	3.2
1	A	543	VAL	3.2
1	A	279	GLY	3.1
1	A	83	TRP	3.1
1	A	11	PRO	3.1
1	A	205	LYS	3.1
1	A	282	LEU	3.1
1	A	281	ASP	3.0
1	A	45	THR	3.0
1	A	196	TYR	2.9
1	A	86	GLU	2.8
1	A	796	CYS	2.8
1	A	555	PRO	2.8
1	A	557	GLU	2.7
1	A	44	LEU	2.7
1	A	696	ASN	2.7
1	A	580	PRO	2.7
1	A	128	ARG	2.7
1	A	14	GLY	2.6
1	A	203	ALA	2.6
1	A	150	VAL	2.6
1	A	13	SER	2.6
1	A	124	SER	2.5
1	A	202	GLN	2.5
1	A	542	ASP	2.5
1	A	125	LEU	2.5
1	A	147	SER	2.5
1	A	194	PRO	2.4
1	A	85	HIS	2.4
1	A	700	ASN	2.3
1	A	25	ASP	2.3
1	A	193	SER	2.3
1	A	8	PHE	2.2
1	A	112	THR	2.2
1	A	461	GLY	2.2
1	A	807	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	84	PRO	2.2
1	A	151	LEU	2.2
1	A	7	THR	2.2
1	A	12	ASP	2.1
1	A	38	ALA	2.1
1	A	220	VAL	2.1
1	A	275	GLU	2.1
1	A	546	TYR	2.1
1	A	460	ASP	2.1
1	A	15	ILE	2.0
1	A	312	CYS	2.0
1	A	559	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	PCA	A	1	8/9	0.71	0.41	-	110,123,135,141	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	NAG	A	905	14/15	0.85	0.30	1.57	76,76,76,76	0
4	NAG	A	904	14/15	0.91	0.36	0.45	76,76,76,76	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	HEM	A	901	43/43	0.93	0.26	-0.18	36,38,44,48	0
3	FAD	A	902	53/53	0.89	0.25	-0.85	42,56,76,81	0
4	NAG	A	906	14/15	0.87	0.26	-	76,76,76,76	0
5	MAN	A	910	11/12	0.77	0.52	-	88,90,91,92	0
4	NAG	A	903	14/15	0.94	0.22	-	76,76,76,76	0
5	MAN	A	907	11/12	0.71	0.47	-	86,93,96,96	0
5	MAN	A	909	11/12	0.73	0.75	-	92,97,99,100	0
5	MAN	A	911	11/12	0.71	0.35	-	86,93,96,96	0
5	MAN	A	908	11/12	0.83	0.42	-	79,84,86,87	0

6.5 Other polymers [i](#)

There are no such residues in this entry.