



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:16 am GMT

PDB ID : 1QNO
Title : The 3-D structure of a Trichoderma reesei b-mannanase from glycoside hydrolase family 5
Authors : Sabini, E.; Schubert, H.; Murshudov, G.; Wilson, K.S.; Siika-Aho, M.; Penttila, M.
Deposited on : 1999-10-20
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	344	 88% 10%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3147 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENDO-1,4-B-D-MANNANASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	344	Total	C	N	O	S	0	4	0
			2677	1692	447	528	10			


- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code:

TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	406	Total	O	0	0
			406	406		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: ENDO-1,4-B-D-MANNANASE

Chain A:  88% 10% .



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.01Å 54.27Å 60.24Å 90.00° 111.31° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00	Depositor
% Data completeness (in resolution range)	97.8 (20.00-2.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.03	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.126 , 0.200	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3147	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.79	0/2777	1.47	33/3799 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	298	ARG	NE-CZ-NH1	15.18	127.89	120.30
1	A	54	ARG	NE-CZ-NH1	11.60	126.10	120.30
1	A	306	TRP	O-C-N	-9.44	107.60	122.70
1	A	145	ARG	CD-NE-CZ	8.21	135.10	123.60
1	A	298	ARG	CD-NE-CZ	8.18	135.05	123.60
1	A	15	ASP	CB-CG-OD1	8.10	125.59	118.30
1	A	213	ASP	CB-CG-OD1	7.99	125.49	118.30
1	A	165	GLU	OE1-CD-OE2	7.71	132.55	123.30
1	A	154	ARG	NE-CZ-NH2	-7.59	116.51	120.30
1	A	213	ASP	CB-CG-OD2	-7.51	111.54	118.30
1	A	236	ASP	CB-CG-OD1	-7.39	111.65	118.30
1	A	63[A]	THR	CA-CB-CG2	-7.35	102.11	112.40
1	A	63[B]	THR	CA-CB-CG2	-7.35	102.11	112.40
1	A	236	ASP	CB-CG-OD2	7.15	124.73	118.30
1	A	87	ASP	CB-CG-OD1	6.78	124.40	118.30
1	A	298	ARG	NH1-CZ-NH2	-6.55	112.20	119.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	306	TRP	N-CA-CB	6.49	122.29	110.60
1	A	42	PHE	CB-CG-CD2	-6.45	116.28	120.80
1	A	157	ASN	CB-CG-OD1	6.35	134.31	121.60
1	A	130	ASN	CB-CG-OD1	6.27	134.13	121.60
1	A	97	GLN	CB-CG-CD	6.14	127.56	111.60
1	A	117	TYR	CB-CG-CD2	5.88	124.53	121.00
1	A	123	TYR	CB-CG-CD1	5.80	124.48	121.00
1	A	123	TYR	CB-CG-CD2	-5.78	117.53	121.00
1	A	39	ASP	CB-CG-OD1	5.66	123.40	118.30
1	A	130	ASN	OD1-CG-ND2	-5.57	109.08	121.90
1	A	117	TYR	CB-CG-CD1	-5.50	117.70	121.00
1	A	145	ARG	NE-CZ-NH2	-5.47	117.56	120.30
1	A	307	GLN	N-CA-CB	5.41	120.34	110.60
1	A	33	THR	CA-CB-CG2	-5.27	105.02	112.40
1	A	306	TRP	CA-C-N	5.13	128.48	117.20
1	A	37	ASP	CB-CG-OD2	-5.09	113.72	118.30
1	A	178	ASP	CB-CG-OD2	-5.01	113.79	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	306	TRP	Mainchain,Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2677	0	2455	13	0
2	A	56	0	52	0	0
3	A	8	0	12	0	0
4	A	406	0	0	3	0
All	All	3147	0	2519	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:THR:HG23	1:A:70:ILE:HD11	1.70	0.73
1:A:33:THR:CG2	1:A:70:ILE:HD11	2.24	0.68
1:A:252:THR:HG23	4:A:2312:HOH:O	2.01	0.60
1:A:65:PRO:HB2	1:A:69:GLN:HB2	1.89	0.55
1:A:97:GLN:HG3	4:A:2153:HOH:O	2.07	0.54
1:A:2:SER:O	1:A:3:SER:OG	2.22	0.54
1:A:343:ILE:O	1:A:344:ASN:C	2.51	0.48
1:A:109:PRO:HA	1:A:165:GLU:O	2.15	0.47
1:A:101:GLN:NE2	4:A:2158:HOH:O	2.45	0.43
1:A:2:SER:O	1:A:3:SER:CB	2.66	0.43
1:A:321:ASP:HB2	1:A:322:PRO:CD	2.49	0.42
1:A:55:VAL:O	1:A:109:PRO:HD2	2.21	0.41
1:A:58:PHE:CE1	1:A:112:ASN:HB2	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/344 (101%)	336 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	286/282 (101%)	282 (99%)	4 (1%)	71 76

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	THR
1	A	56	TRP
1	A	272	CYS
1	A	344	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	GLN
1	A	331	ASN
1	A	344	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	430	1	14,14,15	0.94	1 (7%)	15,19,21	1.66	4 (26%)
2	NAG	A	431	1	14,14,15	1.00	1 (7%)	15,19,21	1.65	4 (26%)
2	NAG	A	432	1	14,14,15	1.15	1 (7%)	15,19,21	2.28	2 (13%)
2	NAG	A	433	1	14,14,15	1.31	2 (14%)	15,19,21	2.07	5 (33%)
3	TRS	A	801	-	7,7,7	1.62	2 (28%)	9,9,9	1.61	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	430	1	-	0/6/23/26	0/1/1/1
2	NAG	A	431	1	-	0/6/23/26	0/1/1/1
2	NAG	A	432	1	-	0/6/23/26	0/1/1/1
2	NAG	A	433	1	-	0/6/23/26	0/1/1/1
3	TRS	A	801	-	-	0/9/9/9	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	432	NAG	O7-C7	-3.21	1.15	1.23
2	A	433	NAG	O7-C7	-3.00	1.16	1.23
2	A	430	NAG	O7-C7	-2.58	1.17	1.23
2	A	431	NAG	O7-C7	-2.37	1.17	1.23
2	A	433	NAG	C2-N2	2.08	1.50	1.46
3	A	801	TRS	C1-C	2.28	1.58	1.52
3	A	801	TRS	O2-C2	2.61	1.50	1.42

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	433	NAG	C2-N2-C7	-4.09	116.98	122.94
2	A	433	NAG	O3-C3-C2	-4.01	100.80	109.39
2	A	430	NAG	C6-C5-C4	-2.71	106.66	113.00
3	A	801	TRS	C1-C-N	-2.62	102.16	107.73
2	A	430	NAG	O3-C3-C2	-2.47	104.09	109.39
2	A	431	NAG	C8-C7-N2	-2.36	111.84	116.11
2	A	433	NAG	O4-C4-C5	-2.23	103.66	109.28
2	A	430	NAG	C4-C3-C2	-2.20	107.80	111.02
3	A	801	TRS	O3-C3-C	-2.17	104.20	110.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	433	NAG	O6-C6-C5	-2.15	104.12	111.34
2	A	431	NAG	C1-C2-N2	-2.09	106.91	110.49
2	A	432	NAG	C1-O5-C5	2.06	115.01	112.17
2	A	430	NAG	O5-C1-C2	2.49	114.94	111.47
3	A	801	TRS	C3-C-N	2.75	113.57	107.73
2	A	431	NAG	O7-C7-C8	2.87	127.29	122.06
2	A	433	NAG	C1-O5-C5	3.31	116.73	112.17
2	A	431	NAG	O5-C1-C2	3.44	116.26	111.47
2	A	432	NAG	O5-C1-C2	7.88	122.44	111.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.