



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 15, 2017 – 03:10 PM EDT

PDB ID : 2QOJ
Title : Coevolution of a homing endonuclease and its host target sequence
Authors : Scalley-Kim, M.; McConnell Smith, A.; Stoddard, B.L.
Deposited on : unknown
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	Z	254	<div> <div>4%</div> <div> <div></div> <div>71%</div> <div>24%</div> <div>• •</div> </div> </div>
2	X	31	<div> <div>6%</div> <div> <div>29%</div> <div>42%</div> <div>23%</div> <div>6%</div> </div> </div>
3	Y	31	<div> <div>10%</div> <div> <div>29%</div> <div>52%</div> <div>16%</div> <div>•</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MG	Z	602	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3344 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Intron-encoded DNA endonuclease I-AniI.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	Z	254	Total	C	N	O	S	Se	0	0	0
			2085	1363	339	379	2	2			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Z	1	GLY	-	EXPRESSION TAG	UNP P03880
Z	2	SER	-	EXPRESSION TAG	UNP P03880
Z	61	ARG	ILE	ENGINEERED	UNP P03880

- Molecule 2 is a DNA chain called I-AniI DNA target seq1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	29	Total	C	N	O	P	0	0	0
			600	284	112	175	29			

- Molecule 3 is a DNA chain called I-AniI DNA target seq2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Y	30	Total	C	N	O	P	0	0	0
			610	289	113	178	30			

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	Z	3	Total	Mg	0	0
			3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	Z	38	Total 38	O 38	0	0
5	X	4	Total 4	O 4	0	0
5	Y	4	Total 4	O 4	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Intron-encoded DNA endonuclease I-AniI



- Molecule 2: I-AniI DNA target seq1



- Molecule 3: I-AniI DNA target seq2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.35Å 72.70Å 61.10Å 90.00° 103.88° 90.00°	Depositor
Resolution (Å)	29.30 – 2.40 29.30 – 2.40	Depositor EDS
% Data completeness (in resolution range)	92.1 (29.30-2.40) 76.7 (29.30-2.40)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	11.96 (at 2.42Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.216 , 0.276 0.212 , 0.269	Depositor DCC
R_{free} test set	778 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	32.6	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.056 for l,-k,h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3344	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	Z	0.66	0/2124	0.79	2/2854 (0.1%)
2	X	1.00	0/673	1.82	24/1038 (2.3%)
3	Y	0.99	1/683 (0.1%)	1.73	23/1050 (2.2%)
All	All	0.81	1/3480 (0.0%)	1.30	49/4942 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Z	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	Y	523	DC	C3'-O3'	-5.12	1.37	1.44

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	X	401	DG	O4'-C1'-N9	-9.22	101.54	108.00
2	X	409	DG	O4'-C1'-N9	8.97	114.28	108.00
2	X	410	DA	O4'-C1'-N9	8.11	113.68	108.00
2	X	404	DC	O4'-C4'-C3'	-7.68	101.39	106.00
2	X	417	DT	O4'-C1'-N1	7.15	113.01	108.00
3	Y	518	DC	O4'-C1'-N1	7.14	113.00	108.00
3	Y	529	DG	O4'-C1'-N9	7.09	112.96	108.00
3	Y	501	DG	P-O3'-C3'	6.98	128.08	119.70
2	X	428	DC	O4'-C1'-N1	6.92	112.84	108.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	Y	516	DA	N1-C2-N3	-6.68	125.96	129.30
3	Y	507	DT	O4'-C1'-N1	6.50	112.55	108.00
2	X	409	DG	C8-N9-C4	-6.46	103.82	106.40
1	Z	62	ASN	N-CA-C	6.45	128.42	111.00
3	Y	519	DT	O4'-C1'-N1	6.39	112.47	108.00
2	X	415	DT	O4'-C1'-N1	-6.37	103.54	108.00
2	X	404	DC	C1'-O4'-C4'	-6.36	103.74	110.10
2	X	419	DT	O4'-C1'-N1	6.28	112.40	108.00
2	X	424	DA	O4'-C1'-N9	6.27	112.39	108.00
2	X	405	DT	N3-C2-O2	-6.17	118.60	122.30
3	Y	514	DA	P-O3'-C3'	6.08	126.99	119.70
3	Y	518	DC	O4'-C1'-C2'	-6.00	101.10	105.90
3	Y	513	DG	O4'-C1'-N9	5.93	112.15	108.00
2	X	410	DA	O4'-C1'-C2'	-5.77	101.28	105.90
3	Y	518	DC	C1'-O4'-C4'	-5.75	104.35	110.10
2	X	427	DG	P-O3'-C3'	5.70	126.54	119.70
2	X	423	DA	O4'-C1'-N9	5.67	111.97	108.00
2	X	410	DA	N1-C2-N3	-5.53	126.53	129.30
2	X	416	DC	N1-C1'-C2'	5.53	123.10	112.60
3	Y	529	DG	P-O3'-C3'	5.45	126.24	119.70
3	Y	522	DT	C4-C5-C7	5.44	122.26	119.00
2	X	406	DG	P-O3'-C3'	5.38	126.16	119.70
3	Y	523	DC	O3'-P-O5'	-5.37	93.80	104.00
2	X	404	DC	O4'-C1'-C2'	-5.37	101.61	105.90
3	Y	505	DT	O4'-C1'-N1	5.34	111.74	108.00
3	Y	516	DA	C4-C5-C6	-5.30	114.35	117.00
2	X	428	DC	C1'-O4'-C4'	-5.30	104.80	110.10
3	Y	526	DC	O4'-C1'-N1	-5.26	104.32	108.00
2	X	406	DG	O4'-C1'-N9	5.25	111.68	108.00
3	Y	520	DC	C5-C4-N4	5.24	123.87	120.20
3	Y	522	DT	O4'-C1'-N1	-5.23	104.34	108.00
2	X	424	DA	C1'-O4'-C4'	-5.19	104.91	110.10
3	Y	520	DC	O4'-C1'-N1	5.19	111.63	108.00
3	Y	504	DC	P-O3'-C3'	5.18	125.91	119.70
1	Z	67	VAL	CB-CA-C	-5.09	101.72	111.40
3	Y	511	DG	O4'-C1'-N9	-5.09	104.43	108.00
3	Y	528	DC	P-O3'-C3'	5.09	125.80	119.70
2	X	429	DA	O4'-C1'-N9	5.05	111.54	108.00
3	Y	520	DC	N3-C4-N4	-5.03	114.48	118.00
2	X	424	DA	O4'-C1'-C2'	-5.01	101.89	105.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	Z	61	ARG	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Z	2085	0	2164	60	0
2	X	600	0	327	8	0
3	Y	610	0	336	8	0
4	Z	3	0	0	0	0
5	X	4	0	0	0	0
5	Y	4	0	0	2	0
5	Z	38	0	0	4	1
All	All	3344	0	2827	70	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Z:239:ARG:HH21	1:Z:251:PRO:HB3	1.33	0.89
1:Z:196:THR:O	1:Z:198:CYS:N	2.04	0.88
1:Z:5:THR:HG22	5:Z:629:HOH:O	1.83	0.77
1:Z:34:ILE:HD12	1:Z:69:LEU:HB3	1.70	0.73
1:Z:5:THR:CG2	5:Z:629:HOH:O	2.37	0.71
1:Z:35:GLU:HG2	1:Z:66:MSE:HE3	1.72	0.70
1:Z:97:ASP:OD1	1:Z:120:ARG:HD3	1.91	0.70
1:Z:60:LYS:O	1:Z:61:ARG:C	2.31	0.69
1:Z:41:VAL:HG22	1:Z:67:VAL:HG21	1.75	0.68
1:Z:95:GLN:HE21	1:Z:95:GLN:HA	1.62	0.64
1:Z:34:ILE:CD1	1:Z:69:LEU:HB3	2.28	0.63
3:Y:516:DA:H2''	3:Y:517:DC:O5'	1.99	0.63
1:Z:41:VAL:CG2	1:Z:67:VAL:HG21	2.29	0.63
1:Z:152:SER:HB2	2:X:419:DT:OP2	1.99	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Z:60:LYS:O	1:Z:62:ASN:HA	1.99	0.62
1:Z:226:ASN:O	1:Z:230:GLN:HG3	2.00	0.62
1:Z:41:VAL:HG22	1:Z:67:VAL:CG2	2.32	0.60
1:Z:251:PRO:O	1:Z:252:SER:HB2	2.01	0.60
1:Z:59:ARG:NH1	2:X:411:DG:O6	2.34	0.59
1:Z:1:GLY:O	1:Z:2:SER:O	2.20	0.59
1:Z:60:LYS:O	1:Z:60:LYS:HG2	2.01	0.59
1:Z:189:THR:HG21	1:Z:202:LYS:O	2.03	0.58
3:Y:526:DC:H2''	3:Y:527:DG:C8	2.39	0.58
1:Z:104:ALA:HA	1:Z:109:ILE:HD12	1.85	0.58
3:Y:521:DC:H2'	3:Y:522:DT:H72	1.85	0.57
1:Z:194:ASP:OD1	1:Z:196:THR:O	2.23	0.57
3:Y:516:DA:C2'	3:Y:517:DC:O5'	2.52	0.57
1:Z:1:GLY:O	1:Z:5:THR:OG1	2.24	0.56
1:Z:95:GLN:HA	1:Z:95:GLN:NE2	2.21	0.55
1:Z:222:LYS:HB2	1:Z:254:TYR:CE2	2.41	0.55
1:Z:204:THR:HG21	3:Y:507:DT:C7	2.36	0.55
3:Y:529:DG:H2''	3:Y:530:DC:C5	2.42	0.54
1:Z:117:ASP:HB2	5:Z:627:HOH:O	2.08	0.53
1:Z:164:ILE:O	1:Z:165:ALA:HB3	2.08	0.53
1:Z:228:LYS:HE3	1:Z:254:TYR:CD2	2.43	0.53
2:X:424:DA:H2''	2:X:425:DG:C8	2.44	0.52
1:Z:225:GLY:O	1:Z:228:LYS:HB3	2.09	0.52
1:Z:204:THR:CG2	5:Y:40:HOH:O	2.58	0.52
1:Z:228:LYS:HE3	1:Z:254:TYR:HD2	1.75	0.52
1:Z:38:ILE:HD12	1:Z:67:VAL:HG22	1.90	0.51
1:Z:128:ILE:HD13	1:Z:176:ILE:HG23	1.93	0.51
2:X:428:DC:H2''	2:X:429:DA:O5'	2.12	0.49
1:Z:150:CYS:HB3	1:Z:168:ASP:OD1	2.14	0.48
1:Z:2:SER:HB3	1:Z:5:THR:HG23	1.96	0.48
1:Z:34:ILE:HD12	1:Z:69:LEU:CB	2.41	0.48
1:Z:228:LYS:CE	1:Z:254:TYR:HD2	2.26	0.47
1:Z:204:THR:HG21	3:Y:507:DT:H72	1.96	0.47
1:Z:61:ARG:NH1	2:X:412:DG:N7	2.62	0.47
1:Z:204:THR:HG23	5:Y:40:HOH:O	2.14	0.46
1:Z:41:VAL:CG2	1:Z:67:VAL:CG2	2.93	0.46
1:Z:13:PHE:CE2	1:Z:34:ILE:HG13	2.51	0.45
1:Z:192:TYR:CE1	3:Y:509:DC:H2'	2.52	0.45
1:Z:80:PHE:O	1:Z:84:ILE:HG12	2.16	0.45
1:Z:100:ARG:HG2	1:Z:118:TYR:CG	2.53	0.43
1:Z:253:ASN:O	1:Z:254:TYR:HB3	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Z:53:ILE:HG13	1:Z:71:ILE:HG12	1.99	0.43
1:Z:249:LYS:HE3	1:Z:249:LYS:HB3	1.73	0.42
1:Z:13:PHE:CE2	1:Z:34:ILE:CG1	3.02	0.42
1:Z:132:ILE:HD12	1:Z:184:TYR:HB2	2.01	0.42
1:Z:132:ILE:HG23	1:Z:184:TYR:HD1	1.85	0.42
2:X:405:DT:H2'	2:X:406:DG:C8	2.55	0.42
2:X:402:DC:H2''	2:X:403:DG:C8	2.55	0.42
1:Z:130:SER:O	1:Z:134:THR:HG23	2.19	0.41
1:Z:58:PHE:CE2	1:Z:67:VAL:HG13	2.55	0.41
1:Z:109:ILE:HD13	1:Z:116:PRO:HD3	2.02	0.41
1:Z:138:SER:O	1:Z:142:VAL:HG23	2.20	0.41
1:Z:89:PRO:HD2	1:Z:136:TYR:OH	2.20	0.41
2:X:401:DG:P	2:X:401:DG:C8	3.14	0.41
1:Z:172:ARG:HA	1:Z:198:CYS:HA	2.02	0.40
1:Z:5:THR:HG21	5:Z:629:HOH:O	2.13	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:Z:610:HOH:O	5:Z:630:HOH:O[2_545]	1.29	0.91

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	Z	252/254 (99%)	236 (94%)	10 (4%)	6 (2%)	7 7

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Z	2	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	Z	62	ASN
1	Z	197	ASN
1	Z	252	SER
1	Z	61	ARG
1	Z	251	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Z	232/230 (101%)	214 (92%)	18 (8%)	15	23

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Z	8	TYR
1	Z	20	SER
1	Z	34	ILE
1	Z	53	ILE
1	Z	67	VAL
1	Z	69	LEU
1	Z	100	ARG
1	Z	111	SER
1	Z	119	THR
1	Z	152	SER
1	Z	189	THR
1	Z	193	LEU
1	Z	196	THR
1	Z	212	ILE
1	Z	235	LEU
1	Z	236	LYS
1	Z	237	GLN
1	Z	238	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	Z	42	GLN
1	Z	95	GLN
1	Z	171	GLN
1	Z	211	ASN
1	Z	218	ASN
1	Z	230	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	Z	252/254 (99%)	0.06	10 (3%) 39 38	12, 24, 41, 52	0
2	X	29/31 (93%)	0.10	2 (6%) 18 16	16, 31, 55, 61	0
3	Y	30/31 (96%)	0.39	3 (10%) 8 7	16, 37, 75, 87	0
All	All	311/316 (98%)	0.10	15 (4%) 31 30	12, 25, 49, 87	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	Y	530	DC	7.5
3	Y	501	DG	5.0
1	Z	62	ASN	3.6
1	Z	253	ASN	3.4
1	Z	1	GLY	3.3
1	Z	195	LYS	2.9
2	X	401	DG	2.6
2	X	429	DA	2.5
1	Z	252	SER	2.4
1	Z	160	ASP	2.4
1	Z	251	PRO	2.3
1	Z	158	LYS	2.2
3	Y	529	DG	2.2
1	Z	159	ASP	2.1
1	Z	17	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	MG	Z	602	1/1	0.80	0.45	5.20	47,47,47,47	0
4	MG	Z	603	1/1	0.96	0.17	-0.47	37,37,37,37	0
4	MG	Z	601	1/1	0.96	0.16	-0.63	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.