



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 01:12 am GMT

PDB ID : 3QQ3  
Title : Crystal structure of swine major histocompatibility complex class I SLA-1 0401 and identification of 2009 pandemic swine-origin influenza A H1N1 virus cytotoxic T lymphocyte epitope peptides  
Authors : Zhang, N.; Qi, J.; Gao, F.; Pan, X.; Chen, R.; Li, Q.; Chen, Z.; Li, X.; Xia, C.; Gao, G.F.  
Deposited on : 2011-02-15  
Resolution : 2.59 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>3%</div> <div> <div></div> <div>73%</div> <div>23%</div> <div>•</div> </div> </div>
1	D	275	<div> <div>3%</div> <div> <div></div> <div>68%</div> <div>28%</div> <div>•</div> </div> </div>
2	B	100	<div> <div>%</div> <div> <div></div> <div>70%</div> <div>28%</div> <div>••</div> </div> </div>
2	E	100	<div> <div>5%</div> <div> <div></div> <div>77%</div> <div>19%</div> <div>••</div> </div> </div>
3	C	9	<div> <div></div> <div> <div></div> <div>89%</div> <div>11%</div> </div> </div>
3	F	9	<div> <div></div> <div> <div></div> <div>67%</div> <div>22%</div> <div>11%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6593 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2233	1394	398	432	9			
1	D	275	Total	C	N	O	S	0	0	0
			2233	1394	398	432	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			817	524	142	148	3			
2	E	99	Total	C	N	O	S	0	0	0
			817	524	142	148	3			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	GLU	-	EXPRESSION TAG	UNP Q07717
B	2	PHE	-	EXPRESSION TAG	UNP Q07717
E	1	GLU	-	EXPRESSION TAG	UNP Q07717
E	2	PHE	-	EXPRESSION TAG	UNP Q07717

- Molecule 3 is a protein called 9-mer peptide from Neuraminidase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			75	47	12	16			
3	F	9	Total	C	N	O	0	0	0
			75	47	12	16			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	112	Total 112	O 112	0	0
4	B	72	Total 72	O 72	0	0
4	C	6	Total 6	O 6	0	0
4	D	102	Total 102	O 102	0	0
4	E	43	Total 43	O 43	0	0
4	F	8	Total 8	O 8	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: MHC class I antigen




#### • Molecule 1: MHC class I antigen




#### • Molecule 2: Beta-2-microglobulin




#### • Molecule 2: Beta-2-microglobulin






● Molecule 3: 9-mer peptide from Neuraminidase



● Molecule 3: 9-mer peptide from Neuraminidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.73Å 37.65Å 111.43Å 90.00° 113.06° 90.00°	Depositor
Resolution (Å)	28.31 – 2.59 34.48 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (28.31-2.59) 99.8 (34.48-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.03 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.200 , 0.262 0.195 , 0.196	Depositor DCC
$R_{free}$ test set	1211 reflections (5.13%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.4	Xtriage
Anisotropy	0.205	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 36.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.006 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6593	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.63 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.5453e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/2293	0.48	0/3114
1	D	0.38	1/2293 (0.0%)	0.53	3/3114 (0.1%)
2	B	0.24	0/843	0.41	0/1145
2	E	0.24	0/843	0.42	0/1145
3	C	0.20	0/78	0.31	0/106
3	F	0.24	0/78	0.35	0/106
All	All	0.33	1/6428 (0.0%)	0.48	3/8730 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	145	ARG	CG-CD	5.08	1.64	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	145	ARG	NE-CZ-NH1	7.77	124.19	120.30
1	D	19	ASP	CB-CA-C	-6.11	98.18	110.40
1	D	18	GLY	C-N-CA	-5.39	108.23	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	15	PRO	Mainchain
1	D	16	ASP	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2233	0	2096	58	1
1	D	2233	0	2096	70	0
2	B	817	0	788	18	0
2	E	817	0	788	20	0
3	C	75	0	61	1	0
3	F	75	0	61	2	0
4	A	112	0	0	5	0
4	B	72	0	0	2	1
4	C	6	0	0	0	0
4	D	102	0	0	3	0
4	E	43	0	0	3	0
4	F	8	0	0	0	0
All	All	6593	0	5890	160	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 160 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:14:ARG:HH21	1:D:19:ASP:CG	1.57	1.07
2:E:6:PRO:HA	2:E:87:THR:HG21	1.37	1.04
1:D:14:ARG:NH2	1:D:19:ASP:OD1	1.91	1.02
2:E:3:VAL:HG12	4:E:331:HOH:O	1.69	0.91
1:D:14:ARG:NH2	1:D:19:ASP:CG	2.27	0.87

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:ASP:OD2	4:B:129:HOH:O[1_565]	2.10	0.10

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/275 (99%)	263 (96%)	10 (4%)	0	100	100
1	D	273/275 (99%)	259 (95%)	14 (5%)	0	100	100
2	B	97/100 (97%)	91 (94%)	6 (6%)	0	100	100
2	E	97/100 (97%)	92 (95%)	5 (5%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	F	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
All	All	754/768 (98%)	718 (95%)	36 (5%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/234 (100%)	222 (95%)	12 (5%)	28	52
1	D	234/234 (100%)	217 (93%)	17 (7%)	16	33
2	B	92/93 (99%)	90 (98%)	2 (2%)	57	81
2	E	92/93 (99%)	87 (95%)	5 (5%)	26	49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	8/8 (100%)	8 (100%)	0	100	100
3	F	8/8 (100%)	7 (88%)	1 (12%)	5	10
All	All	668/670 (100%)	631 (94%)	37 (6%)	25	49

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	17	ARG
1	D	88	SER
2	E	95	LYS
1	D	35	ARG
1	D	48	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	49	ASN
1	D	42	ASN
1	D	262	GLN
3	C	1	ASN
1	A	93	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	275/275 (100%)	-0.14	9 (3%)	47	39	11, 23, 49, 63	0
1	D	275/275 (100%)	-0.07	7 (2%)	58	50	16, 27, 47, 66	0
2	B	99/100 (99%)	-0.39	1 (1%)	82	79	13, 18, 30, 81	0
2	E	99/100 (99%)	0.10	5 (5%)	29	22	19, 31, 52, 83	0
3	C	9/9 (100%)	-0.43	0	100	100	17, 22, 25, 27	0
3	F	9/9 (100%)	-0.25	0	100	100	21, 26, 35, 36	0
All	All	766/768 (99%)	-0.12	22 (2%)	52	45	11, 25, 48, 83	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	197	LEU	5.5
2	B	2	PHE	5.4
2	E	2	PHE	5.1
1	D	145	ARG	4.2
1	A	197	LEU	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.