



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 04:02 pm GMT

PDB ID : 2QSS
Title : Bovine hemoglobin at pH 6.3
Authors : Aranda IV, R.; Richards, M.P.; Phillips Jr., G.N.
Deposited on : 2007-07-31
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1762 (1.76-1.76)
Clashscore	112137	1889 (1.76-1.76)
Ramachandran outliers	110173	1868 (1.76-1.76)
Sidechain outliers	110143	1868 (1.76-1.76)
RSRZ outliers	101464	1770 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	<div> <div>4%</div> <div>88%</div> <div>11%</div> <div>•</div> </div>
1	C	141	<div> <div>4%</div> <div>93%</div> <div>6%</div> <div>•</div> </div>
2	B	145	<div> <div>8%</div> <div>83%</div> <div>15%</div> <div>•</div> </div>
2	D	145	<div> <div>5%</div> <div>84%</div> <div>11%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CMO	C	143	-	-	-	X
3	CMO	D	147	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hemoglobin subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	141	Total	C	N	O	S	0	2	0
			1083	693	190	199	1			
1	C	141	Total	C	N	O	S	0	3	0
			1078	692	186	199	1			

- Molecule 2 is a protein called Hemoglobin subunit beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	145	Total	C	N	O	S	0	7	0
			1172	760	199	209	4			
2	D	145	Total	C	N	O	S	0	5	0
			1161	750	199	207	5			

- Molecule 3 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 2 1 1	0	0
3	B	1	Total C O 2 1 1	0	0
3	C	1	Total C O 2 1 1	0	0
3	D	1	Total C O 2 1 1	0	0

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C Fe N O 43 34 1 4 4	0	0
4	B	1	Total C Fe N O 43 34 1 4 4	0	0
4	C	1	Total C Fe N O 43 34 1 4 4	0	0
4	D	1	Total C Fe N O 43 34 1 4 4	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	153	Total O 153 153	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	114	Total 114	O 114	0	0
5	C	155	Total 155	O 155	0	0
5	D	195	Total 195	O 195	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Hemoglobin subunit alpha





- Molecule 1: Hemoglobin subunit alpha



- Molecule 2: Hemoglobin subunit beta



- Molecule 2: Hemoglobin subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.00Å 78.38Å 109.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.63 – 1.75 17.74 – 1.68	Depositor EDS
% Data completeness (in resolution range)	98.4 (63.63-1.75) 92.7 (17.74-1.68)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 1.68Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.177 , 0.225 0.177 , 0.224	Depositor DCC
R_{free} test set	2879 reflections (5.41%)	DCC
Wilson B-factor (Å ²)	24.7	Xtriage
Anisotropy	0.201	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5291	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/1113	0.67	0/1509
1	C	0.68	0/1114	0.65	1/1512 (0.1%)
2	B	0.65	0/1212	0.64	0/1637
2	D	0.70	0/1195	0.68	0/1610
All	All	0.67	0/4634	0.66	1/6268 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	11	LYS	CD-CE-NZ	5.92	125.33	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1083	0	1087	16	0
1	C	1078	0	1087	10	0
2	B	1172	0	1197	24	0
2	D	1161	0	1184	24	0
3	A	2	0	0	1	0
3	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	2	0	0	1	0
3	D	2	0	0	0	0
4	A	43	0	30	0	0
4	B	43	0	30	1	0
4	C	43	0	30	1	0
4	D	43	0	30	0	0
5	A	153	0	0	3	0
5	B	114	0	0	2	0
5	C	155	0	0	2	0
5	D	195	0	0	9	0
All	All	5291	0	4675	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:10:VAL:HG13	2:B:74:MET:HE1	1.46	0.97
1:A:92[A]:ARG:HH11	1:A:92[A]:ARG:HG2	1.32	0.94
2:D:27:LEU:HD11	2:D:62:HIS:HD2	1.33	0.90
1:C:11:LYS:HE3	1:C:73:LEU:HD12	1.58	0.85
2:D:15:GLY:HA3	5:D:180:HOH:O	1.75	0.85
2:B:42[A]:GLU:CD	2:B:42[A]:GLU:H	1.80	0.85
2:B:10:VAL:CG1	2:B:74:MET:CE	2.59	0.81
2:B:10:VAL:HG13	2:B:74:MET:CE	2.14	0.78
2:B:27[A]:LEU:HD12	2:B:59:VAL:HG13	1.67	0.77
1:A:11:LYS:HE2	1:A:73:LEU:HD12	1.72	0.72
2:B:10:VAL:CG1	2:B:74:MET:HE3	2.19	0.71
2:D:15:GLY:CA	5:D:180:HOH:O	2.36	0.70
2:D:27:LEU:HD11	2:D:62:HIS:CD2	2.24	0.67
2:B:5:GLU:HG3	5:B:166:HOH:O	1.95	0.66
1:C:45:HIS:CD2	1:C:45:HIS:H	2.13	0.66
2:B:1:MET:HG3	2:B:2:LEU:N	2.10	0.66
2:B:10:VAL:HG12	2:B:74:MET:HE3	1.77	0.66
1:A:1:VAL:HG11	5:A:234:HOH:O	1.96	0.65
2:D:89:GLU:HG2	5:D:208:HOH:O	1.99	0.62
4:B:146:HEM:HBB2	4:B:146:HEM:HHC	1.81	0.61
2:B:90:LEU:HA	2:B:94:LYS:HD2	1.83	0.60
1:A:39:THR:HG22	1:A:97:ASN:HD22	1.67	0.60
1:A:58:HIS:HE2	3:A:143:CMO:C	2.17	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:141:ALA:O	2:D:144:TYR:HB2	2.04	0.57
2:D:1[B]:MET:HB3	5:D:285:HOH:O	2.05	0.56
2:B:99:PRO:HG3	2:B:144:TYR:CD2	2.42	0.55
1:A:38:THR:HG23	5:A:153:HOH:O	2.06	0.55
2:B:1:MET:HG3	2:B:2:LEU:HD13	1.88	0.55
2:D:10:VAL:HG13	2:D:74:MET:CE	2.38	0.54
1:C:58:HIS:HE2	3:C:143:CMO:C	2.21	0.54
1:A:1:VAL:CG1	5:A:234:HOH:O	2.56	0.53
1:C:45:HIS:HE1	4:C:142:HEM:O1D	1.92	0.52
1:C:11:LYS:CE	1:C:73:LEU:HD12	2.36	0.52
1:A:92[A]:ARG:NH1	1:A:92[A]:ARG:HG2	2.09	0.52
2:D:103:LYS:HE3	5:D:295:HOH:O	2.11	0.51
1:A:1:VAL:HB	1:A:131:ASN:HD21	1.76	0.50
2:B:13:PHE:CG	2:B:125[A]:LEU:HD21	2.47	0.50
1:A:39:THR:HG22	1:A:97:ASN:ND2	2.26	0.49
2:B:21[A]:GLU:CD	2:B:21[A]:GLU:H	2.16	0.49
2:B:13:PHE:CD2	2:B:125[A]:LEU:HD21	2.48	0.49
2:D:27:LEU:CD1	2:D:62:HIS:HD2	2.15	0.49
2:D:107:ASN:ND2	2:D:130:GLN:HE22	2.11	0.48
1:A:140:TYR:HB3	2:D:35:PRO:HB2	1.96	0.47
2:B:98:ASP:CG	1:C:96[A]:VAL:HG21	2.35	0.46
1:C:87:HIS:HA	1:C:91:LEU:HD12	1.99	0.45
1:C:11:LYS:HD2	5:C:227:HOH:O	2.16	0.45
1:A:7:LYS:HB3	1:A:11:LYS:HE3	1.99	0.45
1:A:119:PRO:HG2	2:B:54:MET:HG3	1.98	0.45
1:A:140:TYR:CE2	2:D:36:TRP:HB3	2.51	0.44
2:D:2:LEU:HB3	2:D:6:GLU:HB2	2.00	0.44
2:D:18[B]:LYS:HZ2	2:D:18[B]:LYS:HG2	1.67	0.44
2:D:27:LEU:O	2:D:31:LEU:HG	2.17	0.44
1:A:33:PHE:CE2	1:A:48:LEU:HD22	2.53	0.43
1:C:140:TYR:HD1	5:C:286:HOH:O	2.01	0.43
1:C:80:LEU:HD22	1:C:83:LEU:HD12	2.01	0.43
2:D:10:VAL:HG13	2:D:74:MET:HE1	2.00	0.43
2:B:121:PHE:HB3	5:B:179:HOH:O	2.17	0.43
2:D:1[B]:MET:HE1	5:D:304:HOH:O	2.19	0.42
1:A:86:LEU:HD12	1:A:90:LYS:HD3	2.02	0.42
2:D:13:PHE:CG	2:D:125:LEU:HD21	2.54	0.42
2:D:64[B]:LYS:HE2	5:D:221:HOH:O	2.20	0.41
2:D:64[A]:LYS:NZ	2:D:65[A]:LYS:HD3	2.35	0.41
2:D:119:LYS:HE2	5:D:240:HOH:O	2.22	0.40
2:D:1[A]:MET:HB2	5:D:285:HOH:O	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:18:LYS:HA	2:B:18:LYS:HD2	1.83	0.40
2:B:1:MET:HB3	2:B:77:LEU:O	2.20	0.40
2:D:107:ASN:HD21	2:D:130:GLN:HE22	1.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/141 (100%)	140 (99%)	1 (1%)	0	100	100
1	C	142/141 (101%)	140 (99%)	2 (1%)	0	100	100
2	B	150/145 (103%)	145 (97%)	5 (3%)	0	100	100
2	D	147/145 (101%)	144 (98%)	3 (2%)	0	100	100
All	All	580/572 (101%)	569 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/112 (102%)	110 (96%)	4 (4%)	41	16
1	C	115/112 (103%)	115 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	B	125/118 (106%)	114 (91%)	11 (9%)	12 1
2	D	123/118 (104%)	113 (92%)	10 (8%)	14 2
All	All	477/460 (104%)	452 (95%)	25 (5%)	37 7

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	THR
1	A	92[A]	ARG
1	A	92[B]	ARG
1	A	141	ARG
2	B	1	MET
2	B	2	LEU
2	B	42[A]	GLU
2	B	42[B]	GLU
2	B	60	LYS
2	B	66[A]	VAL
2	B	66[B]	VAL
2	B	87	LEU
2	B	109[A]	LEU
2	B	109[B]	LEU
2	B	143	ARG
2	D	1[A]	MET
2	D	1[B]	MET
2	D	10	VAL
2	D	18[A]	LYS
2	D	18[B]	LYS
2	D	27	LEU
2	D	64[A]	LYS
2	D	64[B]	LYS
2	D	89	GLU
2	D	144	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	97	ASN
2	B	107	ASN
1	C	9	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	45	HIS
2	D	76	HIS
2	D	107	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	HEM	A	142	1,3	28,50,50	2.10	8 (28%)	17,82,82	2.44	7 (41%)
3	CMO	A	143	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	B	146	3,2	28,50,50	2.44	11 (39%)	17,82,82	2.44	9 (52%)
3	CMO	B	147	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	C	142	1,3	28,50,50	2.19	10 (35%)	17,82,82	2.33	8 (47%)
3	CMO	C	143	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	D	146	3,2	28,50,50	2.21	11 (39%)	17,82,82	2.25	8 (47%)
3	CMO	D	147	4	0,1,1	0.00	-	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEM	A	142	1,3	-	0/6/54/54	0/0/8/8
3	CMO	A	143	4	-	0/0/0/0	0/0/0/0
4	HEM	B	146	3,2	-	0/6/54/54	0/0/8/8
3	CMO	B	147	4	-	0/0/0/0	0/0/0/0
4	HEM	C	142	1,3	-	0/6/54/54	0/0/8/8
3	CMO	C	143	4	-	0/0/0/0	0/0/0/0
4	HEM	D	146	3,2	-	0/6/54/54	0/0/8/8
3	CMO	D	147	4	-	0/0/0/0	0/0/0/0

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	146	HEM	C3B-C2B	-5.52	1.33	1.40
4	C	142	HEM	C3C-C2C	-5.23	1.33	1.40
4	A	142	HEM	C3B-C2B	-4.71	1.34	1.40
4	D	146	HEM	C3B-C2B	-4.37	1.34	1.40
4	B	146	HEM	C3C-C2C	-3.93	1.35	1.40
4	D	146	HEM	C3C-C2C	-3.55	1.35	1.40
4	A	142	HEM	C3C-C2C	-3.33	1.35	1.40
4	C	142	HEM	C3B-C2B	-3.31	1.36	1.40
4	D	146	HEM	CMC-C2C	2.02	1.55	1.51
4	C	142	HEM	C1B-NB	2.07	1.39	1.36
4	B	146	HEM	CAD-C3D	2.13	1.56	1.52
4	D	146	HEM	CAA-C2A	2.14	1.55	1.52
4	C	142	HEM	CAD-C3D	2.14	1.56	1.52
4	A	142	HEM	CMB-C2B	2.19	1.56	1.51
4	D	146	HEM	C4B-NB	2.23	1.40	1.36
4	C	142	HEM	C3C-CAC	2.31	1.52	1.47
4	B	146	HEM	C4B-NB	2.37	1.41	1.36
4	B	146	HEM	CMB-C2B	2.37	1.56	1.51
4	C	142	HEM	CAA-C2A	2.39	1.56	1.52
4	D	146	HEM	C3C-CAC	2.43	1.52	1.47
4	C	142	HEM	CMB-C2B	2.50	1.56	1.51
4	A	142	HEM	C4D-ND	2.53	1.39	1.36
4	A	142	HEM	C3C-CAC	2.64	1.53	1.47
4	B	146	HEM	C1B-NB	2.65	1.39	1.36
4	D	146	HEM	CAD-C3D	2.81	1.57	1.52
4	A	142	HEM	CAA-C2A	2.81	1.56	1.52
4	D	146	HEM	C1B-NB	2.91	1.40	1.36
4	B	146	HEM	CAA-C2A	3.14	1.57	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	142	HEM	C4D-ND	3.22	1.40	1.36
4	B	146	HEM	C3C-CAC	3.35	1.54	1.47
4	D	146	HEM	C4D-ND	3.49	1.40	1.36
4	B	146	HEM	C4D-ND	4.01	1.41	1.36
4	A	142	HEM	C3B-CAB	4.03	1.55	1.47
4	B	146	HEM	C3B-CAB	4.11	1.56	1.47
4	D	146	HEM	C3B-CAB	4.28	1.56	1.47
4	C	142	HEM	C3D-C2D	4.45	1.50	1.37
4	A	142	HEM	C3D-C2D	4.51	1.51	1.37
4	D	146	HEM	C3D-C2D	4.64	1.51	1.37
4	C	142	HEM	C3B-CAB	4.67	1.57	1.47
4	B	146	HEM	C3D-C2D	5.11	1.52	1.37

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	142	HEM	CBD-CAD-C3D	-5.13	102.68	112.47
4	A	142	HEM	CAD-CBD-CGD	-4.97	104.17	112.66
4	B	146	HEM	CMA-C3A-C4A	-4.65	121.32	128.46
4	A	142	HEM	CBD-CAD-C3D	-4.59	103.71	112.47
4	C	142	HEM	CBA-CAA-C2A	-4.28	104.31	112.48
4	B	146	HEM	CBD-CAD-C3D	-4.07	104.70	112.47
4	D	146	HEM	CMA-C3A-C4A	-4.01	122.30	128.46
4	D	146	HEM	C1D-C2D-C3D	-4.00	104.22	107.00
4	A	142	HEM	C1D-C2D-C3D	-3.82	104.34	107.00
4	B	146	HEM	CAD-CBD-CGD	-3.47	106.73	112.66
4	A	142	HEM	CMA-C3A-C4A	-3.46	123.14	128.46
4	C	142	HEM	CMA-C3A-C4A	-3.26	123.46	128.46
4	D	146	HEM	CBA-CAA-C2A	-3.22	106.33	112.48
4	B	146	HEM	C3C-C4C-NC	-2.78	105.70	110.94
4	D	146	HEM	CBD-CAD-C3D	-2.76	107.21	112.47
4	D	146	HEM	CAD-CBD-CGD	-2.75	107.96	112.66
4	B	146	HEM	CBA-CAA-C2A	-2.71	107.31	112.48
4	C	142	HEM	C3C-C4C-NC	-2.56	106.10	110.94
4	D	146	HEM	C3C-C4C-NC	-2.54	106.14	110.94
4	C	142	HEM	C1D-C2D-C3D	-2.30	105.40	107.00
4	B	146	HEM	C1D-C2D-C3D	-2.19	105.47	107.00
4	A	142	HEM	C3C-C4C-NC	-2.01	107.15	110.94
4	C	142	HEM	C4C-C3C-C2C	2.05	108.33	106.90
4	D	146	HEM	CMC-C2C-C3C	2.05	128.70	124.89
4	B	146	HEM	C4A-C3A-C2A	2.09	108.45	107.00
4	C	142	HEM	CMB-C2B-C3B	2.20	128.98	124.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	142	HEM	CMA-C3A-C2A	2.27	129.22	124.94
4	A	142	HEM	CMC-C2C-C3C	2.48	129.50	124.89
4	C	142	HEM	C4A-C3A-C2A	2.56	108.78	107.00
4	B	146	HEM	CMA-C3A-C2A	2.80	130.23	124.94
4	B	146	HEM	CMC-C2C-C3C	2.82	130.13	124.89
4	D	146	HEM	CMA-C3A-C2A	2.83	130.28	124.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	143	CMO	1	0
4	B	146	HEM	1	0
4	C	142	HEM	1	0
3	C	143	CMO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	141/141 (100%)	0.09	5 (3%)	44	51	13, 18, 31, 44	0
1	C	141/141 (100%)	-0.03	6 (4%)	36	43	13, 16, 32, 43	0
2	B	145/145 (100%)	0.45	11 (7%)	15	19	15, 23, 40, 44	0
2	D	145/145 (100%)	0.11	7 (4%)	31	37	14, 21, 28, 58	0
All	All	572/572 (100%)	0.16	29 (5%)	29	35	13, 19, 37, 58	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	140	TYR	10.5
1	A	1	VAL	7.9
2	B	87	LEU	7.3
1	A	141	ARG	6.2
2	D	144	TYR	5.6
1	A	140	TYR	5.3
2	B	84	PHE	4.4
2	D	1[A]	MET	4.3
2	D	143	ARG	4.2
2	D	145	HIS	4.2
2	D	141	ALA	4.1
1	C	1	VAL	4.0
2	B	140	LEU	4.0
1	A	137	THR	4.0
2	B	19	VAL	3.7
2	B	59	VAL	3.4
2	B	145	HIS	3.2
2	B	2	LEU	3.1
2	D	142	HIS	3.1
1	C	137	THR	2.9
1	C	141	ARG	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	20	ASP	2.5
2	B	77	LEU	2.3
2	D	48	SER	2.3
2	B	85	ALA	2.2
2	B	142	HIS	2.1
1	A	70	VAL	2.1
1	C	139	LYS	2.1
1	C	78	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	CMO	D	147	2/2	0.98	0.15	2.85	23,23,23,25	0
3	CMO	C	143	2/2	0.97	0.13	2.07	20,20,20,25	0
3	CMO	B	147	2/2	0.94	0.11	0.31	28,28,28,29	0
3	CMO	A	143	2/2	0.96	0.10	-0.07	21,21,21,25	0
4	HEM	B	146	43/43	0.95	0.12	-0.10	21,28,35,40	0
4	HEM	A	142	43/43	0.96	0.09	-0.21	16,20,34,45	0
4	HEM	C	142	43/43	0.97	0.08	-0.50	16,19,31,40	0
4	HEM	D	146	43/43	0.97	0.08	-0.64	18,23,31,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.