



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 06:26 am GMT

PDB ID : 4QSH
Title : Crystal Structure of *L. monocytogenes* Pyruvate Carboxylase in complex with Cyclic-di-AMP
Authors : Choi, P.H.; Tong, L.
Deposited on : 2014-07-04
Resolution : 2.51 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1148	 3% 81% 11% • 7%
1	B	1148	 4% 83% 9% • 6%
1	C	1148	 4% 82% 11% • 6%
1	D	1148	 2% 81% 11% • 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FLC	A	2001	-	-	X	-
2	FLC	B	2001	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 34524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pyruvate carboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1072	8447	5361	1435	1612	39	0	0	0
1	B	1074	8436	5356	1431	1608	41	0	0	0
1	C	1081	8501	5396	1442	1623	40	0	0	0
1	D	1072	8446	5363	1431	1611	41	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	HIS	-	EXPRESSION TAG	UNP W6G6F5
A	0	MET	-	EXPRESSION TAG	UNP W6G6F5
B	-1	HIS	-	EXPRESSION TAG	UNP W6G6F5
B	0	MET	-	EXPRESSION TAG	UNP W6G6F5
C	-1	HIS	-	EXPRESSION TAG	UNP W6G6F5
C	0	MET	-	EXPRESSION TAG	UNP W6G6F5
D	-1	HIS	-	EXPRESSION TAG	UNP W6G6F5
D	0	MET	-	EXPRESSION TAG	UNP W6G6F5

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: C₆H₅O₇).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 13 6 7	0	0
2	B	1	Total C O 13 6 7	0	0
2	C	1	Total C O 13 6 7	0	0
2	D	1	Total C O 13 6 7	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mn 1 1	0	0
3	A	1	Total Mn 1 1	0	0
3	D	1	Total Mn 1 1	0	0
3	C	1	Total Mn 1 1	0	0

- Molecule 4 is (2R,3R,3AS,5R,7AR,9R,10R,10AS,12R,14AR)-2,9-BIS(6-AMINO-9H-PURIN-9-YL)OCTAHYDRO-2H,7H-DIFURO[3,2-D:3',2'-J][1,3,7,9,2,8]TETRAOXADIPHOSPHACYCLODODECINE-3,5,10,12-TETROL 5,12-DIOXIDE (three-letter code: 2BA) (formula: C₂₀H₂₄N₁₀O₁₂P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	N	O	P	0	0
			44	20	10	12	2		
4	A	1	Total	C	N	O	P	0	0
			44	20	10	12	2		
4	C	1	Total	C	N	O	P	0	0
			44	20	10	12	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	163	Total	O	0	0
			163	163		
5	B	79	Total	O	0	0
			79	79		
5	C	82	Total	O	0	0
			82	82		
5	D	182	Total	O	0	0
			182	182		


3 Residue-property plots i


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyruvate carboxylase





- Molecule 1: Pyruvate carboxylase








• Molecule 1: Pyruvate carboxylase



• Molecule 1: Pyruvate carboxylase





4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	96.78Å 153.30Å 221.24Å 90.00° 101.58° 90.00°	Depositor
Resolution (Å)	49.80 – 2.51 49.75 – 2.51	Depositor EDS
% Data completeness (in resolution range)	96.6 (49.80-2.51) 96.7 (49.75-2.51)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.17 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.195 , 0.233 0.200 , 0.236	Depositor DCC
R_{free} test set	10515 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	47.9	Xtrriage
Anisotropy	0.163	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.015 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	34524	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC, MN, 2BA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	3/8614 (0.0%)	0.87	19/11669 (0.2%)
1	B	0.68	1/8603 (0.0%)	0.85	14/11658 (0.1%)
1	C	0.68	1/8670 (0.0%)	0.84	10/11747 (0.1%)
1	D	0.81	2/8613 (0.0%)	0.88	17/11666 (0.1%)
All	All	0.75	7/34500 (0.0%)	0.86	60/46740 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	213	GLU	CD-OE1	7.13	1.33	1.25
1	D	813	TYR	CE1-CZ	6.95	1.47	1.38
1	C	213	GLU	CD-OE1	6.81	1.33	1.25
1	D	213	GLU	CD-OE1	6.52	1.32	1.25
1	A	590	GLU	CG-CD	6.50	1.61	1.51

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	570	MET	CA-CB-CG	-9.90	96.46	113.30
1	A	712	MET	CG-SD-CE	8.35	113.56	100.20
1	B	570	MET	CG-SD-CE	-7.50	88.21	100.20
1	D	989	MET	CG-SD-CE	-7.36	88.43	100.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	413	ARG	NE-CZ-NH1	7.00	123.80	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	682	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8447	0	8377	79	0
1	B	8436	0	8357	76	0
1	C	8501	0	8419	89	0
1	D	8446	0	8392	83	0
2	A	13	0	5	4	0
2	B	13	0	5	5	0
2	C	13	0	5	1	0
2	D	13	0	5	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	88	0	46	0	0
4	C	44	0	23	0	0
5	A	163	0	0	14	0
5	B	79	0	0	5	0
5	C	82	0	0	7	0
5	D	182	0	0	12	0
All	All	34524	0	33634	318	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 318 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:97:ARG:HD3	5:A:2174:HOH:O	1.53	1.08
1:C:125:GLN:OE1	5:C:1359:HOH:O	1.83	0.95
1:C:945:GLY:HA2	1:C:948:MET:HE2	1.51	0.91
1:A:723:ARG:HD3	5:A:2115:HOH:O	1.71	0.90
1:B:582:ASP:OD2	1:B:586:ARG:NH1	2.08	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1066/1148 (93%)	1026 (96%)	40 (4%)	0	100	100
1	B	1068/1148 (93%)	1027 (96%)	41 (4%)	0	100	100
1	C	1075/1148 (94%)	1034 (96%)	41 (4%)	0	100	100
1	D	1066/1148 (93%)	1021 (96%)	45 (4%)	0	100	100
All	All	4275/4592 (93%)	4108 (96%)	167 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	919/983 (94%)	879 (96%)	40 (4%)	33	57
1	B	916/983 (93%)	882 (96%)	34 (4%)	39	66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	924/983 (94%)	889 (96%)	35 (4%)	38	64
1	D	921/983 (94%)	884 (96%)	37 (4%)	36	62
All	All	3680/3932 (94%)	3534 (96%)	146 (4%)	36	62

5 of 146 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	940	LEU
1	C	120	ILE
1	D	915	MET
1	B	941	THR
1	C	-1	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 62 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	824	GLN
1	C	112	HIS
1	D	479	ASN
1	C	43	HIS
1	C	292	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FLC	A	2001	-	3,12,12	1.55	1 (33%)	3,17,17	3.07	2 (66%)
4	2BA	A	2003	-	44,50,50	1.29	1 (2%)	46,78,78	2.61	12 (26%)
4	2BA	A	2004	-	44,50,50	1.51	7 (15%)	46,78,78	2.65	13 (28%)
2	FLC	B	2001	-	3,12,12	1.24	0	3,17,17	2.66	1 (33%)
4	2BA	C	1201	-	44,50,50	1.16	5 (11%)	46,78,78	2.32	11 (23%)
2	FLC	C	1202	-	3,12,12	1.03	0	3,17,17	4.44	2 (66%)
2	FLC	D	2001	-	3,12,12	1.34	1 (33%)	3,17,17	2.97	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FLC	A	2001	-	-	0/6/16/16	0/0/0/0
4	2BA	A	2003	-	-	0/22/62/62	0/6/7/7
4	2BA	A	2004	-	-	0/22/62/62	0/6/7/7
2	FLC	B	2001	-	-	0/6/16/16	0/0/0/0
4	2BA	C	1201	-	-	0/22/62/62	0/6/7/7
2	FLC	C	1202	-	-	0/6/16/16	0/0/0/0
2	FLC	D	2001	-	-	0/6/16/16	0/0/0/0

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	1201	2BA	C5-N7	-2.13	1.32	1.39
4	C	1201	2BA	C2'-C1'	-2.08	1.50	1.53
4	C	1201	2BA	O4'-C1'	2.01	1.44	1.41
2	D	2001	FLC	OHB-CB	2.31	1.46	1.43
4	A	2004	2BA	C51-C41	2.39	1.45	1.40

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2003	2BA	N31-C21-N11	-10.40	119.80	128.86
4	A	2004	2BA	N31-C21-N11	-10.25	119.93	128.86
4	A	2003	2BA	N3-C2-N1	-9.33	120.74	128.86
4	C	1201	2BA	N31-C21-N11	-9.22	120.83	128.86
4	A	2004	2BA	N3-C2-N1	-8.59	121.38	128.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2001	FLC	4	0
2	B	2001	FLC	5	0
2	C	1202	FLC	1	0
2	D	2001	FLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1072/1148 (93%)	-0.08	33 (3%) 49 52	17, 47, 86, 120	0
1	B	1074/1148 (93%)	0.02	43 (4%) 39 41	34, 56, 100, 128	0
1	C	1081/1148 (94%)	0.05	45 (4%) 37 39	30, 57, 98, 130	0
1	D	1072/1148 (93%)	-0.12	22 (2%) 64 66	20, 45, 84, 123	0
All	All	4299/4592 (93%)	-0.03	143 (3%) 47 50	17, 51, 94, 130	0

The worst 5 of 143 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	906	PHE	7.0
1	C	933	VAL	6.5
1	C	202	VAL	6.5
1	C	904	ILE	6.1
1	C	201	TYR	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	MN	B	2002	1/1	0.79	0.16	0.63	94,94,94,94	0
3	MN	C	1203	1/1	0.98	0.15	-0.02	85,85,85,85	0
4	2BA	A	2003	44/44	0.99	0.12	-0.03	21,32,38,44	0
3	MN	D	2002	1/1	0.93	0.16	-0.08	75,75,75,75	0
4	2BA	C	1201	44/44	0.98	0.11	-0.31	33,49,59,65	0
2	FLC	B	2001	13/13	0.91	0.15	-0.39	46,53,59,69	0
2	FLC	A	2001	13/13	0.95	0.12	-0.71	43,55,59,62	0
2	FLC	D	2001	13/13	0.98	0.08	-1.33	41,46,56,59	0
3	MN	A	2002	1/1	0.87	0.14	-1.74	72,72,72,72	0
2	FLC	C	1202	13/13	0.95	0.12	-2.38	47,54,59,62	0
4	2BA	A	2004	44/44	0.89	0.16	-	35,71,86,94	0

6.5 Other polymers [i](#)

There are no such residues in this entry.