



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 13, 2017 – 06:24 AM EST

PDB ID : 4QSZ  
Title : Crystal structure of mouse JMjd7 fused with maltose-binding protein  
Authors : Liu, H.; Wang, C.; Zhang, G.Y.  
Deposited on : unknown  
Resolution : 2.86 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2469 (2.90-2.82)
Clashscore	112137	2749 (2.90-2.82)
Ramachandran outliers	110173	2687 (2.90-2.82)
Sidechain outliers	110143	2690 (2.90-2.82)
RSRZ outliers	101464	2487 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	687	 80% 18% .
1	B	687	 74% 24% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	A	902	-	-	-	X
3	FLC	A	903	-	-	-	X
3	FLC	B	903	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10832 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein, JmjC domain-containing protein 7 chimera.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	686	Total	C	N	O	S	0	0	0
			5395	3472	892	1012	19			
1	B	682	Total	C	N	O	S	0	0	0
			5365	3453	888	1005	19			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	ALA	-	LINKER	UNP P0C872
A	-8	ALA	-	LINKER	UNP P0C872
A	-7	ALA	-	LINKER	UNP P0C872
A	-6	GLN	-	LINKER	UNP P0C872
A	-5	THR	-	LINKER	UNP P0C872
A	-4	ASN	-	LINKER	UNP P0C872
A	-3	ALA	-	LINKER	UNP P0C872
A	-2	ALA	-	LINKER	UNP P0C872
A	-1	ALA	-	LINKER	UNP P0C872
A	0	GLU	-	LINKER	UNP P0C872
A	1	PHE	-	LINKER	UNP P0C872
B	-9	ALA	-	LINKER	UNP P0C872
B	-8	ALA	-	LINKER	UNP P0C872
B	-7	ALA	-	LINKER	UNP P0C872
B	-6	GLN	-	LINKER	UNP P0C872
B	-5	THR	-	LINKER	UNP P0C872
B	-4	ASN	-	LINKER	UNP P0C872
B	-3	ALA	-	LINKER	UNP P0C872
B	-2	ALA	-	LINKER	UNP P0C872
B	-1	ALA	-	LINKER	UNP P0C872
B	0	GLU	-	LINKER	UNP P0C872
B	1	PHE	-	LINKER	UNP P0C872

- Molecule 2 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 3 is CITRATE ANION (three-letter code: FLC) (formula:  $C_6H_5O_7^-$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose-binding periplasmic protein, JmjC domain-containing protein 7 chimera



- Molecule 1: Maltose-binding periplasmic protein, JmjC domain-containing protein 7 chimera



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.09 Å 158.92 Å 100.01 Å 90.00° 94.08° 90.00°	Depositor
Resolution (Å)	62.15 – 2.86 62.15 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.7 (62.15-2.86) 97.8 (62.15-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.66 (at 2.69 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, $R_{free}$	0.191 , 0.264 0.191 , 0.264	Depositor DCC
$R_{free}$ test set	1681 reflections (4.98%)	DCC
Wilson B-factor (Å <sup>2</sup> )	38.0	Xtriage
Anisotropy	0.208	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 29.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	10832	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.99	1/5547 (0.0%)	0.71	3/7567 (0.0%)
1	B	0.50	0/5518	0.65	1/7531 (0.0%)
All	All	0.79	1/11065 (0.0%)	0.68	4/15098 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	316	ASP	CG-OD2	63.77	2.72	1.25

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	316	ASP	CB-CG-OD2	-15.69	104.18	118.30
1	A	139	LEU	CA-CB-CG	9.65	137.50	115.30
1	A	316	ASP	OD1-CG-OD2	7.57	137.68	123.30
1	B	23	LEU	CA-CB-CG	6.57	130.41	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	134	ASN	Peptide
1	B	-199	GLU	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5395	0	5269	77	0
1	B	5365	0	5239	108	0
2	A	23	0	22	5	0
2	B	23	0	22	4	0
3	A	13	0	5	3	0
3	B	13	0	5	1	0
All	All	10832	0	10562	190	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 190 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:-235:ASP:O	1:B:-231:LYS:N	2.00	0.93
1:A:137:THR:H	1:A:138:GLU:HA	1.33	0.93
1:A:130:LYS:HE3	1:A:167:TRP:CD1	2.07	0.89
1:B:207:PRO:HA	1:B:316:ASP:HB3	1.54	0.87
1:B:-125:VAL:HA	1:B:-48:ASN:HD21	1.43	0.83

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	682/687 (99%)	644 (94%)	35 (5%)	3 (0%)	38	68
1	B	680/687 (99%)	630 (93%)	46 (7%)	4 (1%)	28	60
All	All	1362/1374 (99%)	1274 (94%)	81 (6%)	7 (0%)	32	64

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	134	ASN
1	A	135	LEU
1	A	-202	PHE
1	B	294	MET
1	B	-289	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	567/568 (100%)	552 (97%)	15 (3%)	51	80
1	B	564/568 (99%)	552 (98%)	12 (2%)	59	84
All	All	1131/1136 (100%)	1104 (98%)	27 (2%)	54	82

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	190	SER
1	B	-316	ASP
1	B	186	TYR
1	A	201	SER
1	A	-113	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	134	ASN
1	A	249	GLN
1	B	178	HIS
1	B	304	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLC	A	901	-	12,12,12	1.61	2 (16%)	17,17,17	1.50	4 (23%)
2	GLC	A	902	-	11,11,12	1.81	2 (18%)	13,15,17	1.37	1 (7%)
3	FLC	A	903	-	3,12,12	1.38	0	3,17,17	1.29	0
2	GLC	B	901	-	12,12,12	1.40	1 (8%)	17,17,17	1.44	3 (17%)
2	GLC	B	902	-	11,11,12	1.71	2 (18%)	13,15,17	1.48	3 (23%)
3	FLC	B	903	-	3,12,12	1.41	0	3,17,17	1.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	901	-	-	0/2/22/22	0/1/1/1
2	GLC	A	902	-	-	0/2/19/22	0/1/1/1
3	FLC	A	903	-	-	0/6/16/16	0/0/0/0
2	GLC	B	901	-	-	0/2/22/22	0/1/1/1
2	GLC	B	902	-	-	0/2/19/22	0/1/1/1
3	FLC	B	903	-	-	0/6/16/16	0/0/0/0

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	902	GLC	C2-C3	-4.09	1.47	1.52
2	B	902	GLC	C2-C3	-3.14	1.48	1.52
2	A	901	GLC	C3-C2	-2.45	1.46	1.52
2	B	901	GLC	O5-C1	3.37	1.49	1.43
2	A	902	GLC	O5-C1	3.86	1.50	1.43

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	902	GLC	O3-C3-C2	-3.77	103.16	110.02
2	B	901	GLC	O4-C4-C5	-3.55	100.33	109.28
2	B	901	GLC	C6-C5-C4	-2.71	106.66	113.00
2	A	901	GLC	O4-C4-C3	-2.61	104.67	110.36
2	A	901	GLC	C6-C5-C4	-2.60	106.92	113.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	GLC	2	0
2	A	902	GLC	4	0
3	A	903	FLC	3	0
2	B	901	GLC	3	0
2	B	902	GLC	3	0
3	B	903	FLC	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	686/687 (99%)	-0.57	1 (0%) 95 95	12, 29, 55, 85	0
1	B	682/687 (99%)	-0.41	2 (0%) 93 93	12, 35, 67, 99	0
All	All	1368/1374 (99%)	-0.49	3 (0%) 94 95	12, 31, 63, 99	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	129	GLN	2.7
1	B	-232	LEU	2.5
1	B	316	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	FLC	B	903	13/13	0.78	0.30	3.91	43,62,83,83	0
3	FLC	A	903	13/13	0.91	0.19	3.38	26,34,43,49	0
2	GLC	A	902	11/12	0.96	0.16	2.64	14,17,25,28	0
2	GLC	A	901	12/12	0.96	0.14	0.28	15,18,24,26	0
2	GLC	B	902	11/12	0.94	0.15	-0.17	33,39,46,47	0
2	GLC	B	901	12/12	0.95	0.12	-0.98	25,28,33,37	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.