



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2017 – 07:28 AM EST

PDB ID : 4QTJ  
Title : Complex of WOPR domain of Wor1 in *Candida albicans* with the 13bp dsDNA  
Authors : Zhang, S.; Zhang, T.; Ding, J.  
Deposited on : unknown  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	<div> <div>14%</div> <div>46%</div> <div>10%</div> <div>•</div> <div>44%</div> </div>
2	B	13	<div> <div>85%</div> <div>15%</div> </div>
3	C	13	<div> <div>92%</div> <div>8%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1865 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called White-opaque regulator 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	154	Total	C	N	O	S	0	0	0
			1229	777	217	232	3			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	MET	-	EXPRESSION TAG	UNP Q5AP80
A	273	HIS	-	EXPRESSION TAG	UNP Q5AP80
A	274	HIS	-	EXPRESSION TAG	UNP Q5AP80
A	275	HIS	-	EXPRESSION TAG	UNP Q5AP80
A	276	HIS	-	EXPRESSION TAG	UNP Q5AP80
A	277	HIS	-	EXPRESSION TAG	UNP Q5AP80
A	278	HIS	-	EXPRESSION TAG	UNP Q5AP80

- Molecule 2 is a DNA chain called DNA (5'-D(\*AP\*AP\*AP\*AP\*GP\*TP\*TP\*TP\*AP\*AP\*CP\*TP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	13	Total	C	N	O	P	0	0	0
			264	129	48	75	12			

- Molecule 3 is a DNA chain called DNA (5'-D(\*AP\*AP\*GP\*TP\*TP\*AP\*AP\*AP\*CP\*TP\*TP\*TP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	13	Total	C	N	O	P	0	0	0
			263	129	45	77	12			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	46	Total 46	O 46	0	0
4	B	39	Total 39	O 39	0	0
4	C	24	Total 24	O 24	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.63Å 46.63Å 215.50Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.49 – 2.10 24.48 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.1 (24.49-2.10) 95.2 (24.48-2.10)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.23 (at 2.10Å)	Xtriage
Refinement program	PHENIX, REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.206 , 0.248 0.205 , 0.252	Depositor DCC
$R_{free}$ test set	810 reflections (5.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	49.5	Xtriage
Anisotropy	0.277	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 65.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.038 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	1865	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.86	0/1252	0.91	0/1692
2	B	0.86	0/296	1.10	0/455
3	C	0.83	0/294	1.06	1/452 (0.2%)
All	All	0.85	0/1842	0.97	1/2599 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	C	4	DT	O5'-P-OP1	-5.44	100.81	105.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1229	0	1237	27	0
2	B	264	0	150	3	0
3	C	263	0	151	0	0
4	A	46	0	0	0	0
4	B	39	0	0	0	0
4	C	24	0	0	0	0
All	All	1865	0	1538	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 9.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:VAL:HG21	1:A:55:VAL:HG21	1.56	0.86
1:A:214:THR:HB	1:A:217:GLU:OE2	1.85	0.76
1:A:214:THR:HG21	1:A:224:ALA:HB1	1.70	0.71
2:B:1:DA:H2''	2:B:2:DA:H5''	1.73	0.69
1:A:214:THR:HG22	1:A:216:LYS:H	1.59	0.67
1:A:214:THR:CG2	1:A:224:ALA:HB1	2.31	0.60
1:A:248:PRO:O	1:A:250:GLU:O	2.21	0.59
1:A:27:VAL:HG21	1:A:55:VAL:CG2	2.30	0.58
1:A:19:ASP:HB3	1:A:248:PRO:CG	2.34	0.57
1:A:26:GLN:HG3	1:A:31:GLN:HB2	1.86	0.57
1:A:260:PRO:O	1:A:264:THR:HG23	2.06	0.56
1:A:19:ASP:CG	1:A:248:PRO:HD2	2.26	0.56
2:B:1:DA:C2'	2:B:2:DA:H5''	2.35	0.56
1:A:57:ILE:N	1:A:57:ILE:HD12	2.26	0.51
1:A:19:ASP:HA	1:A:248:PRO:HG2	1.95	0.48
1:A:7:VAL:HG22	1:A:8:PRO:HD2	1.95	0.48
1:A:19:ASP:CB	1:A:248:PRO:CG	2.91	0.48
2:B:1:DA:H2''	2:B:2:DA:C5'	2.41	0.48
1:A:239:ASP:HB2	1:A:245:LEU:HD22	1.95	0.48
1:A:231:LEU:HD23	1:A:231:LEU:C	2.35	0.47
1:A:19:ASP:CB	1:A:248:PRO:HG2	2.44	0.47
1:A:19:ASP:HB3	1:A:248:PRO:HG3	1.98	0.45
1:A:19:ASP:OD1	1:A:248:PRO:HG2	2.18	0.43
1:A:10:TYR:HE1	1:A:252:ASP:OD2	2.00	0.43
1:A:5:MET:CG	1:A:37:ARG:HH21	2.32	0.43
1:A:251:SER:OG	1:A:252:ASP:N	2.52	0.43
1:A:57:ILE:N	1:A:57:ILE:CD1	2.83	0.42
1:A:19:ASP:CA	1:A:248:PRO:HG2	2.50	0.42
1:A:5:MET:HG2	1:A:37:ARG:HH21	1.84	0.41
1:A:247:ARG:HB3	1:A:248:PRO:HD2	2.04	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/274 (55%)	147 (98%)	3 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	138/245 (56%)	132 (96%)	6 (4%)	33	32

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	VAL
1	A	223	LYS
1	A	245	LEU
1	A	257	GLN
1	A	259	SER
1	A	270	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	154/274 (56%)	1.13	38 (24%) 1 1	36, 60, 90, 103	0
2	B	13/13 (100%)	-0.56	0 100 100	49, 50, 62, 73	0
3	C	13/13 (100%)	0.00	0 100 100	39, 49, 79, 79	0
All	All	180/300 (60%)	0.93	38 (21%) 1 1	36, 58, 89, 103	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	231	LEU	7.8
1	A	232	ILE	6.0
1	A	209	ILE	5.9
1	A	55	VAL	5.7
1	A	240	ILE	5.7
1	A	223	LYS	4.8
1	A	23	VAL	4.8
1	A	53	VAL	4.3
1	A	241	ASP	4.1
1	A	66	TRP	4.1
1	A	270	SER	4.0
1	A	250	GLU	3.9
1	A	24	ILE	3.9
1	A	56	PHE	3.8
1	A	224	ALA	3.8
1	A	54	PHE	3.7
1	A	20	ALA	3.5
1	A	233	SER	3.5
1	A	229	ILE	3.5
1	A	249	SER	3.2
1	A	230	HIS	3.1
1	A	82	LEU	2.9
1	A	208	THR	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	234	TYR	2.6
1	A	219	HIS	2.6
1	A	222	GLY	2.6
1	A	237	LYS	2.5
1	A	255	HIS	2.5
1	A	251	SER	2.4
1	A	14	ILE	2.4
1	A	254	LYS	2.4
1	A	267	GLN	2.2
1	A	214	THR	2.2
1	A	81	PHE	2.1
1	A	21	LEU	2.1
1	A	242	SER	2.1
1	A	225	GLU	2.1
1	A	210	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.