



Full wwPDB X-ray Structure Validation Report i

Jul 25, 2017 – 08:41 PM EDT

PDB ID : 2QU6
Title : Crystal structure of the VEGFR2 kinase domain in complex with a benzoxazole inhibitor
Authors : Whittington, D.A.; Kim, J.L.; Long, A.M.; Rose, P.; Gu, Y.; Zhao, H.
Deposited on : unknown
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20029824
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4698 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Vascular endothelial growth factor receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	0	0
			2297	1472	404	405	16			
1	B	260	Total	C	N	O	S	0	0	0
			2092	1346	361	369	16			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	817	ALA	CYS	ENGINEERED	UNP P35968
A	916	THR	VAL	ENGINEERED	UNP P35968
A	990	VAL	GLU	ENGINEERED	UNP P35968
A	1054	PTR	TYR	MODIFIED RESIDUE	UNP P35968
A	1059	PTR	TYR	MODIFIED RESIDUE	UNP P35968
A	1172	ARG	-	EXPRESSION TAG	UNP P35968
A	1173	HIS	-	EXPRESSION TAG	UNP P35968
A	1174	HIS	-	EXPRESSION TAG	UNP P35968
A	1175	HIS	-	EXPRESSION TAG	UNP P35968
A	1176	HIS	-	EXPRESSION TAG	UNP P35968
A	1177	HIS	-	EXPRESSION TAG	UNP P35968
A	1178	HIS	-	EXPRESSION TAG	UNP P35968
B	817	ALA	CYS	ENGINEERED	UNP P35968
B	916	THR	VAL	ENGINEERED	UNP P35968
B	990	VAL	GLU	ENGINEERED	UNP P35968
B	1054	PTR	TYR	MODIFIED RESIDUE	UNP P35968
B	1059	PTR	TYR	MODIFIED RESIDUE	UNP P35968
B	1172	ARG	-	EXPRESSION TAG	UNP P35968
B	1173	HIS	-	EXPRESSION TAG	UNP P35968
B	1174	HIS	-	EXPRESSION TAG	UNP P35968
B	1175	HIS	-	EXPRESSION TAG	UNP P35968
B	1176	HIS	-	EXPRESSION TAG	UNP P35968
B	1177	HIS	-	EXPRESSION TAG	UNP P35968
B	1178	HIS	-	EXPRESSION TAG	UNP P35968

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 4-({2-[{4-chloro-3-[(2S)-1-methylpyrrolidin-2-yl]methoxy}phenyl]amino}-1,3-benzoxazol-5-yl)oxy)-N-methylpyridine-2-carboxamide (three-letter code: 857) (formula: C₂₆H₂₆ClN₅O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf				
3	A	1	Total	C	Cl	N	O	0	0
			36	26	1	5	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
3	B	1	36	26	1	5	4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	176	Total	O 176	0	0
4	B	56	Total	O 56	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Vascular endothelial growth factor receptor 2



- Molecule 1: Vascular endothelial growth factor receptor 2



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.37 Å 67.37 Å 89.62 Å 90.00° 92.98° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10 29.83 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.6 (30.00-2.10) 93.4 (29.83-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.93 (at 2.00 Å)	Xtriage
Refinement program	CNS	Depositor
R , R_{free}	0.242 , 0.272 0.219 , 0.246	Depositor DCC
R_{free} test set	1487 reflections (4.03%)	DCC
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.474	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 57.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4698	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 857, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/2355	0.63	0/3176
1	B	0.37	0/2141	0.59	0/2889
All	All	0.38	0/4496	0.61	0/6065

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2297	0	2282	29	0
1	B	2092	0	2106	44	0
2	A	5	0	0	1	0
3	A	36	0	26	0	0
3	B	36	0	26	2	0
4	A	176	0	0	2	0
4	B	56	0	0	2	0
All	All	4698	0	4440	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:838:LYS:HD2	1:B:1039:LYS:HE2	1.64	0.78
1:B:1027:ARG:HH22	3:B:502:857:H22	1.59	0.67
1:B:1159:HIS:NE2	1:B:1163:LEU:HD11	2.10	0.66
1:B:1160:LEU:HD23	1:B:1163:LEU:HD12	1.79	0.64
1:A:839:PRO:HA	1:A:849:ILE:HG22	1.80	0.61
1:B:920:LYS:HD2	1:B:921:PHE:CE1	2.36	0.61
1:A:884:SER:O	1:A:888:ILE:HG13	2.02	0.60
1:A:875:THR:HG22	1:A:877:SER:H	1.66	0.59
1:B:1128:PRO:HG2	1:B:1131:THR:HB	1.84	0.59
1:B:833:ARG:HH21	1:B:855:GLY:HA3	1.69	0.57
1:A:924:LEU:O	1:A:928:LEU:HG	2.05	0.57
1:A:993:GLU:OE1	1:A:1175:HIS:HE1	1.88	0.57
1:B:870:LEU:HD11	1:B:879:HIS:HA	1.87	0.56
1:B:920:LYS:HD2	1:B:921:PHE:CZ	2.40	0.56
1:B:933:ASN:HB2	4:B:263:HOH:O	2.05	0.56
1:A:1026:HIS:O	1:A:1027:ARG:HB2	2.06	0.56
1:B:1152:THR:OG1	1:B:1155:GLU:HG3	2.07	0.54
1:A:838:LYS:HB3	1:B:1038:GLU:HG2	1.89	0.53
1:B:1067:LEU:HB3	1:B:1072:MET:CE	2.39	0.53
1:B:1075:GLU:HG2	1:B:1076:THR:N	2.23	0.53
1:B:899:VAL:HG13	1:B:899:VAL:O	2.08	0.53
1:A:1070:LYS:HD3	1:A:1111:ILE:HD11	1.90	0.53
1:A:1169:GLN:HG3	2:A:503:SO4:O2	2.09	0.51
1:A:900:ASN:HD22	1:A:900:ASN:N	2.06	0.51
1:B:1067:LEU:HB3	1:B:1072:MET:HE1	1.90	0.51
1:B:1081:VAL:O	1:B:1081:VAL:HG23	2.10	0.51
1:B:1085:GLN:NE2	1:B:1147:PRO:O	2.43	0.51
1:A:1022:ARG:HG3	1:A:1022:ARG:HH21	1.76	0.50
1:A:1117:ARG:O	1:A:1121:GLU:HG3	2.11	0.49
1:A:1099:PHE:CE2	1:A:1135:MET:HG2	2.48	0.49
1:B:849:ILE:HG22	1:B:850:GLU:N	2.28	0.49
1:B:894:HIS:CG	1:B:895:HIS:N	2.81	0.49
1:B:892:ILE:HG13	1:B:893:GLY:N	2.27	0.48
1:B:1153:PHE:O	1:B:1157:VAL:HG23	2.13	0.48
1:B:888:ILE:O	1:B:892:ILE:HG23	2.14	0.47
1:A:899:VAL:O	1:A:899:VAL:HG13	2.15	0.46
1:B:1027:ARG:NH2	3:B:502:857:H22	2.27	0.46
1:A:838:LYS:HD3	1:B:1038:GLU:HG2	1.97	0.46
1:A:993:GLU:OE1	1:A:1175:HIS:CE1	2.69	0.45
1:A:1028:ASP:CG	1:A:1032:ARG:HH22	2.19	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:855:GLY:HA2	4:B:241:HOH:O	2.17	0.45
1:B:1159:HIS:CD2	1:B:1163:LEU:HD11	2.52	0.44
1:A:870:LEU:HD11	1:A:879:HIS:HA	2.00	0.44
1:B:1022:ARG:O	1:B:1023:LYS:HB2	2.17	0.44
1:B:1118:ARG:HD3	1:B:1123:THR:OG1	2.17	0.44
1:B:892:ILE:O	1:B:1022:ARG:NE	2.51	0.44
1:A:1020:ALA:HB1	1:A:1084:ILE:HD13	2.00	0.43
1:B:1127:ALA:HB1	1:B:1131:THR:HG21	2.00	0.43
1:B:1134:GLU:OE1	1:B:1134:GLU:N	2.38	0.43
1:B:885:GLU:O	1:B:889:LEU:HG	2.17	0.43
1:B:847:GLN:HG2	1:B:849:ILE:HD11	2.00	0.43
1:B:838:LYS:HA	1:B:839:PRO:HD3	1.83	0.43
1:B:1112:ASP:O	1:B:1115:PHE:HB3	2.19	0.43
1:A:838:LYS:CD	1:B:1039:LYS:HE2	2.42	0.42
1:B:884:SER:O	1:B:888:ILE:HG13	2.19	0.42
1:B:1010:PHE:CZ	1:B:1014:LYS:HD3	2.54	0.42
1:B:1139:MET:HB3	1:B:1143:TRP:CH2	2.54	0.42
1:A:1075:GLU:HB3	4:A:204:HOH:O	2.19	0.42
1:A:817:ALA:O	1:A:820:LEU:HB2	2.19	0.42
1:B:1089:TRP:CE3	1:B:1143:TRP:HA	2.55	0.42
1:A:898:VAL:O	1:A:899:VAL:C	2.56	0.42
1:B:823:ASP:OD1	1:B:825:SER:HB3	2.20	0.42
1:B:1025:ILE:O	1:B:1027:ARG:NH2	2.54	0.41
1:A:817:ALA:HA	1:A:820:LEU:HD12	2.03	0.41
1:A:1027:ARG:HD2	4:A:13:HOH:O	2.21	0.41
1:B:886:LEU:O	1:B:890:ILE:HG13	2.21	0.41
1:A:1106:TYR:CE1	1:A:1125:MET:HG3	2.57	0.40
1:A:899:VAL:CG1	1:A:899:VAL:O	2.68	0.40
1:A:919:CYS:SG	1:A:1043:LYS:HD2	2.61	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/314 (87%)	267 (98%)	6 (2%)	0	100	100
1	B	252/314 (80%)	246 (98%)	6 (2%)	0	100	100
All	All	525/628 (84%)	513 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/270 (93%)	247 (99%)	3 (1%)	75	81
1	B	229/270 (85%)	225 (98%)	4 (2%)	66	72
All	All	479/540 (89%)	472 (98%)	7 (2%)	70	76

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	929	ARG
1	A	995	LEU
1	A	1039	LYS
1	B	838	LYS
1	B	883	MET
1	B	1038	GLU
1	B	1165	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	894	HIS
1	A	900	ASN
1	A	933	ASN
1	A	1040	ASN
1	A	1162	ASN
1	A	1165	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1167	ASN
1	A	1173	HIS
1	A	1175	HIS
1	B	847	GLN
1	B	933	ASN
1	B	1040	ASN
1	B	1165	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	857	A	501	-	36,40,40	2.25	13 (36%)	44,56,56	2.89	17 (38%)
2	SO4	A	503	-	4,4,4	0.27	0	6,6,6	0.30	0
3	857	B	502	-	36,40,40	2.33	12 (33%)	44,56,56	2.77	17 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	857	A	501	-	-	0/17/29/29	0/4/5/5
2	SO4	A	503	-	-	0/0/0/0	0/0/0/0
3	857	B	502	-	-	0/17/29/29	0/4/5/5

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	857	C25-N2	-2.38	1.35	1.40
3	B	502	857	C11-C20	-2.31	1.45	1.50
3	B	502	857	C6-C25	2.08	1.43	1.39
3	A	501	857	C11-N3	2.17	1.38	1.34
3	A	501	857	C21-C24	2.19	1.42	1.38
3	A	501	857	C21-C10	2.20	1.43	1.38
3	B	502	857	C5-N1	2.24	1.50	1.48
3	A	501	857	C5-N1	2.41	1.51	1.48
3	B	502	857	C7-C10	2.46	1.44	1.39
3	B	502	857	C11-N3	2.48	1.39	1.34
3	B	502	857	C6-C7	2.58	1.43	1.38
3	A	501	857	C6-C7	2.75	1.43	1.38
3	A	501	857	C7-C10	3.40	1.45	1.39
3	B	502	857	C8-C13	3.47	1.45	1.38
3	A	501	857	C8-C13	3.48	1.45	1.38
3	A	501	857	C1-C18	3.77	1.46	1.38
3	B	502	857	C1-C18	3.83	1.46	1.38
3	A	501	857	C17-C1	3.91	1.44	1.36
3	A	501	857	C24-C25	4.09	1.46	1.39
3	B	502	857	C17-C1	4.16	1.45	1.36
3	A	501	857	C19-N2	4.75	1.44	1.36
3	B	502	857	C24-C25	4.82	1.47	1.39
3	B	502	857	C14-C18	5.23	1.46	1.37
3	B	502	857	C19-N2	5.27	1.45	1.36
3	A	501	857	C14-C18	5.35	1.46	1.37

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	857	C22-N5-C20	-8.64	111.83	121.84
3	A	501	857	C8-C9-N3	-7.62	115.14	123.92
3	B	502	857	C8-C9-N3	-7.11	115.72	123.92
3	A	501	857	C22-N5-C20	-6.80	113.96	121.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	857	C18-C14-C15	-5.94	115.40	119.13
3	B	502	857	C18-C14-C15	-5.01	115.98	119.13
3	B	502	857	C24-C21-C10	-3.59	114.48	119.98
3	A	501	857	C24-C21-C10	-3.25	115.00	119.98
3	A	501	857	O1-C13-C12	-3.04	109.26	119.08
3	B	502	857	O1-C13-C12	-2.87	109.82	119.08
3	A	501	857	C6-C25-N2	-2.84	111.28	120.36
3	B	502	857	C12-C11-N3	-2.71	119.86	123.17
3	A	501	857	C17-C1-C18	-2.51	116.79	120.15
3	B	502	857	C6-C25-N2	-2.43	112.57	120.36
3	A	501	857	C25-C6-C7	-2.34	116.24	119.58
3	A	501	857	C12-C11-N3	-2.33	120.32	123.17
3	B	502	857	C7-C10-CL	-2.24	116.83	119.42
3	B	502	857	C25-C6-C7	-2.16	116.49	119.58
3	B	502	857	C18-O1-C13	2.31	124.36	118.78
3	B	502	857	C24-C25-C6	2.37	122.44	119.66
3	B	502	857	C1-C18-C14	2.43	124.10	120.82
3	B	502	857	C8-C13-C12	2.53	123.96	120.53
3	A	501	857	C3-C4-C5	2.70	111.08	104.72
3	A	501	857	C21-C10-C7	2.90	125.12	120.88
3	B	502	857	C3-C4-C5	2.92	111.58	104.72
3	A	501	857	C8-C13-C12	3.05	124.67	120.53
3	B	502	857	C21-C10-C7	3.25	125.62	120.88
3	A	501	857	C1-C18-C14	3.28	125.25	120.82
3	A	501	857	C20-C11-N3	3.69	121.90	117.61
3	A	501	857	C24-C25-C6	3.70	124.01	119.66
3	A	501	857	O3-C7-C10	4.44	121.85	116.38
3	B	502	857	C20-C11-N3	4.87	123.27	117.61
3	B	502	857	C9-N3-C11	7.56	126.95	116.94
3	A	501	857	C9-N3-C11	8.40	128.06	116.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	503	SO4	1	0
3	B	502	857	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/314 (90%)	0.07	11 (3%) 40 47	18, 29, 51, 68	0
1	B	260/314 (82%)	0.58	30 (11%) 5 7	26, 44, 64, 73	0
All	All	543/628 (86%)	0.31	41 (7%) 15 19	18, 35, 61, 73	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	842	ARG	5.2
1	A	858	LYS	4.7
1	B	1078	PHE	4.6
1	A	846	GLY	4.5
1	A	819	ARG	4.4
1	B	872	GLU	4.2
1	B	871	LYS	4.1
1	B	841	GLY	4.1
1	B	908	PRO	3.8
1	A	871	LYS	3.7
1	A	870	LEU	3.7
1	B	1094	LEU	3.7
1	A	876	HIS	3.7
1	B	1067	LEU	3.6
1	B	1095	LEU	3.5
1	B	1080	ARG	3.5
1	B	1110	LYS	3.3
1	B	1126	ARG	3.1
1	A	878	GLU	2.9
1	B	1091	PHE	2.9
1	B	1118	ARG	2.9
1	A	880	ARG	2.8
1	B	1079	ASP	2.8
1	B	1165	GLN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1111	ILE	2.7
1	B	1012	VAL	2.6
1	B	1044	ILE	2.6
1	B	1117	ARG	2.6
1	B	909	GLY	2.5
1	B	1098	ILE	2.4
1	B	825	SER	2.3
1	B	999	PHE	2.3
1	A	817	ALA	2.3
1	B	1085	GLN	2.2
1	B	1114	GLU	2.2
1	B	1158	GLU	2.2
1	B	1029	LEU	2.1
1	B	899	VAL	2.1
1	A	877	SER	2.1
1	B	1034	ILE	2.1
1	A	1094	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	857	B	502	36/36	0.91	0.14	-0.19	32,37,51,52	0
3	857	A	501	36/36	0.95	0.11	-0.80	27,33,44,46	0
2	SO4	A	503	5/5	0.76	0.33	-	69,70,70,72	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.