



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 09:01 pm GMT

PDB ID : 4QWQ
Title : Crystal structure of the DNA-binding domain of the response regulator SaeR from *Staphylococcus aureus*
Authors : Fan, X.; Zhu, Y.; Zhang, X.; Teng, M.; Li, X.
Deposited on : 2014-07-17
Resolution : 2.50 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	110	
1	B	110	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1585 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Response regulator SaeR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	100	800	520	129	150	1	0	0	0
1	A	96	774	504	129	140	1	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	119	HIS	-	EXPRESSION TAG	UNP H9BRQ0
B	120	HIS	-	EXPRESSION TAG	UNP H9BRQ0
B	121	HIS	-	EXPRESSION TAG	UNP H9BRQ0
B	122	HIS	-	EXPRESSION TAG	UNP H9BRQ0
A	119	HIS	-	EXPRESSION TAG	UNP H9BRQ0
A	120	HIS	-	EXPRESSION TAG	UNP H9BRQ0
A	121	HIS	-	EXPRESSION TAG	UNP H9BRQ0
A	122	HIS	-	EXPRESSION TAG	UNP H9BRQ0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	6	Total	O	0	0
			6	6		
2	A	5	Total	O	0	0
			5	5		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	31.97Å 62.65Å 116.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.32 – 2.50 31.32 – 2.50	Depositor EDS
% Data completeness (in resolution range)	95.5 (31.32-2.50) 99.2 (31.32-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.24 (at 2.51Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6_289)	Depositor
R, R_{free}	0.205 , 0.281 0.234 , 0.266	Depositor DCC
R_{free} test set	399 reflections (4.91%)	DCC
Wilson B-factor (Å ²)	47.0	Xtrriage
Anisotropy	0.911	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 59.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1585	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.22	1/790 (0.1%)	0.76	1/1071 (0.1%)
1	B	1.56	2/818 (0.2%)	0.85	0/1113
All	All	1.40	3/1608 (0.2%)	0.80	1/2184 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	222	TYR	CD1-CE1	-5.64	1.30	1.39
1	B	184	TYR	CD2-CE2	-5.33	1.31	1.39
1	A	222	TYR	CD2-CE2	-5.24	1.31	1.39

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	134	GLU	N-CA-C	5.11	124.80	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	774	0	756	31	0
1	B	800	0	750	28	0
2	A	5	0	0	0	0
2	B	6	0	0	0	0
All	All	1585	0	1506	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:201:ARG:O	1:B:205:GLU:HG3	1.61	1.00
1:A:209:PHE:CE1	1:A:211:THR:HG22	1.97	0.98
1:B:216:THR:HG22	1:B:222:TYR:CE1	2.10	0.86
1:B:216:THR:HG22	1:B:222:TYR:HE1	1.38	0.85
1:A:131:SER:OG	1:A:136:THR:HG22	1.77	0.83
1:A:133:ASP:CB	1:A:211:THR:HG21	2.10	0.81
1:B:130:LEU:HD23	1:B:162:TRP:CZ3	2.18	0.77
1:B:172:ILE:O	1:B:221:GLY:HA3	1.89	0.73
1:B:134:GLU:N	1:B:134:GLU:OE2	2.22	0.71
1:A:209:PHE:HE1	1:A:211:THR:HG22	1.53	0.70
1:A:139:ASN:O	1:A:142:LYS:HD2	1.91	0.69
1:A:131:SER:OG	1:A:136:THR:CG2	2.41	0.68
1:A:201:ARG:NH1	1:A:214:ILE:O	2.27	0.67
1:B:128:GLU:HG3	1:B:129:GLN:H	1.62	0.64
1:B:216:THR:CG2	1:B:222:TYR:HE1	2.12	0.62
1:A:209:PHE:CE1	1:A:211:THR:CG2	2.81	0.61
1:A:144:VAL:HB	1:A:151:VAL:CG2	2.31	0.61
1:B:168:GLU:O	1:B:169:ASN:HB2	2.01	0.60
1:A:129:GLN:HB3	1:A:162:TRP:HH2	1.69	0.58
1:B:128:GLU:HG3	1:B:129:GLN:N	2.19	0.57
1:A:151:VAL:HG23	1:A:151:VAL:O	2.04	0.57
1:B:130:LEU:CD2	1:B:162:TRP:CZ3	2.88	0.57
1:A:211:THR:HG23	1:A:212:TYR:HD1	1.69	0.57
1:A:168:GLU:O	1:A:169:ASN:HB2	2.04	0.56
1:B:201:ARG:O	1:B:205:GLU:CG	2.48	0.54
1:B:153:MET:HG2	1:B:203:LYS:HD3	1.89	0.54
1:B:130:LEU:CD2	1:B:162:TRP:CH2	2.91	0.53
1:A:171:VAL:HG22	1:A:217:VAL:HG21	1.91	0.53
1:A:155:ILE:O	1:A:159:GLU:HG3	2.08	0.53
1:B:172:ILE:O	1:B:222:TYR:N	2.39	0.53
1:A:197:ILE:O	1:A:201:ARG:HG3	2.09	0.52
1:B:176:GLU:O	1:B:180:LYS:HG2	2.10	0.51
1:A:209:PHE:HE1	1:A:211:THR:CG2	2.19	0.50
1:A:191:ASN:OD1	1:A:194:ASN:HB2	2.12	0.50
1:B:158:PHE:CD1	1:B:158:PHE:C	2.85	0.50
1:B:130:LEU:HD22	1:B:162:TRP:CH2	2.48	0.48
1:A:144:VAL:HB	1:A:151:VAL:HG22	1.94	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:143:VAL:HG13	1:B:150:GLU:HG3	1.94	0.48
1:B:163:TYR:CZ	1:B:180:LYS:HG3	2.48	0.48
1:B:139:ASN:HD21	1:B:142:LYS:NZ	2.11	0.48
1:A:131:SER:HA	1:A:135:LEU:O	2.13	0.48
1:A:130:LEU:HD12	1:A:162:TRP:CZ3	2.49	0.48
1:A:174:LYS:NZ	1:A:194:ASN:HD21	2.12	0.47
1:A:174:LYS:HE2	1:A:194:ASN:OD1	2.14	0.46
1:B:160:LEU:HD21	1:B:197:ILE:HD13	1.96	0.46
1:A:160:LEU:CD1	1:A:177:LEU:HD22	2.47	0.45
1:A:144:VAL:HB	1:A:151:VAL:HG23	1.99	0.45
1:B:172:ILE:O	1:B:221:GLY:CA	2.62	0.45
1:A:171:VAL:HG23	1:A:223:LYS:HE2	1.98	0.45
1:B:139:ASN:HD22	1:B:139:ASN:C	2.21	0.44
1:B:139:ASN:ND2	1:B:142:LYS:NZ	2.66	0.43
1:B:211:THR:OG1	1:B:212:TYR:CD1	2.72	0.43
1:B:172:ILE:HG22	1:B:173:SER:N	2.34	0.42
1:A:196:HIS:O	1:A:200:ILE:HG13	2.19	0.42
1:A:151:VAL:HA	1:A:152:PRO:HD3	1.87	0.42
1:A:210:THR:O	1:A:211:THR:C	2.57	0.42
1:A:203:LYS:O	1:A:207:GLU:HG2	2.20	0.41
1:B:190:ALA:C	1:B:192:THR:N	2.73	0.41
1:A:137:LEU:HD23	1:A:144:VAL:HG13	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	92/110 (84%)	84 (91%)	8 (9%)	0	100	100
1	B	98/110 (89%)	93 (95%)	5 (5%)	0	100	100
All	All	190/220 (86%)	177 (93%)	13 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	82/104 (79%)	79 (96%)	3 (4%)	39	66
1	B	81/104 (78%)	62 (76%)	19 (24%)	1	1
All	All	163/208 (78%)	141 (86%)	22 (14%)	4	8

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	129	GLN
1	B	130	LEU
1	B	131	SER
1	B	134	GLU
1	B	136	THR
1	B	137	LEU
1	B	139	ASN
1	B	151	VAL
1	B	164	LEU
1	B	178	LEU
1	B	179	GLU
1	B	185	ASP
1	B	192	THR
1	B	204	LEU
1	B	208	SER
1	B	211	THR
1	B	216	THR
1	B	223	LYS
1	B	225	GLU
1	A	175	SER
1	A	189	ASP
1	A	220	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	129	GLN
1	B	139	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	96/110 (87%)	-0.22	1 (1%) 82 83	28, 52, 77, 80	0
1	B	100/110 (90%)	-0.22	0 100 100	28, 59, 86, 98	0
All	All	196/220 (89%)	-0.22	1 (0%) 90 91	28, 55, 80, 98	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	128	GLU	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.