



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 11:48 pm GMT

PDB ID : 3R04  
Title : The discovery of novel benzofuran-2-carboxylic acids as potent Pim-1 inhibitors  
Authors : Xiang, Y.; Hirth, B.; Asmussen, G.; Biemann, H.-P.; Good, A.; Fitzgerald, M.; Gladysheva, T.; Jancsics, K.; Liu, J.; Metz, M.; Papoulis, A.; Skerlj, R.; Stepp, D.J.; Wei, R.R.  
Deposited on : 2011-03-07  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3453 (1.70-1.70)
Clashscore	112137	3876 (1.70-1.70)
Ramachandran outliers	110173	3815 (1.70-1.70)
Sidechain outliers	110143	3815 (1.70-1.70)
RSRZ outliers	101464	3491 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IMD	A	4497	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2497 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proto-oncogene serine/threonine-protein kinase pim-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	271	2225	1422	389	406	8	0	2	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	MET	-	INITIATING METHIONINE	UNP P11309
A	314	ALA	-	EXPRESSION TAG	UNP P11309
A	315	ALA	-	EXPRESSION TAG	UNP P11309
A	316	ALA	-	EXPRESSION TAG	UNP P11309
A	317	LEU	-	EXPRESSION TAG	UNP P11309
A	318	GLU	-	EXPRESSION TAG	UNP P11309
A	319	HIS	-	EXPRESSION TAG	UNP P11309
A	320	HIS	-	EXPRESSION TAG	UNP P11309
A	321	HIS	-	EXPRESSION TAG	UNP P11309
A	322	HIS	-	EXPRESSION TAG	UNP P11309
A	323	HIS	-	EXPRESSION TAG	UNP P11309
A	324	HIS	-	EXPRESSION TAG	UNP P11309
A	325	HIS	-	EXPRESSION TAG	UNP P11309
A	326	HIS	-	EXPRESSION TAG	UNP P11309

- Molecule 2 is 5-{6-[(TRANS-4-AMINOCYCLOHEXYL)AMINO]PYRAZIN-2-YL}-1-BENZOFURAN-2-CARBOXYLIC ACID (three-letter code: UNQ) (formula: C<sub>19</sub>H<sub>20</sub>N<sub>4</sub>O<sub>3</sub>).




- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	231	Total 231	O 231	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene serine/threonine-protein kinase pim-1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.94Å 97.94Å 80.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	84.82 – 1.70 42.41 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.3 (84.82-1.70) 99.3 (42.41-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.90 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.177 , 0.206 0.184 , 0.214	Depositor DCC
$R_{free}$ test set	2435 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.6	Xtrriage
Anisotropy	0.345	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 42.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.037 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2497	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNQ, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.56	20/2284 (0.9%)	1.27	21/3101 (0.7%)

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	258	ARG	CZ-NH1	13.33	1.50	1.33
1	A	258	ARG	CZ-NH2	8.87	1.44	1.33
1	A	100	PHE	CE1-CZ	8.86	1.54	1.37
1	A	246	GLU	CG-CD	8.31	1.64	1.51
1	A	235	MET	SD-CE	-7.96	1.33	1.77
1	A	160	ASN	CB-CG	-6.42	1.36	1.51
1	A	269	TRP	CE3-CZ3	6.22	1.49	1.38
1	A	260	SER	CA-CB	6.03	1.61	1.52
1	A	226	TRP	CZ3-CH2	5.96	1.49	1.40
1	A	146	SER	CB-OG	5.89	1.50	1.42
1	A	211	GLU	CB-CG	5.75	1.63	1.52
1	A	246	GLU	CB-CG	5.72	1.63	1.52
1	A	281	PHE	CD1-CE1	5.68	1.50	1.39
1	A	289	TRP	CB-CG	5.52	1.60	1.50
1	A	254	PHE	CE2-CZ	5.51	1.47	1.37
1	A	262	GLU	CD-OE1	5.42	1.31	1.25
1	A	282	GLU	CG-CD	5.33	1.59	1.51
1	A	69	VAL	CB-CG2	5.30	1.64	1.52
1	A	291	GLN	CG-CD	5.14	1.62	1.51
1	A	176	ASP	CB-CG	5.12	1.62	1.51

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	258	ARG	NE-CZ-NH2	-12.42	114.09	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	136	ARG	NE-CZ-NH2	-12.00	114.30	120.30
1	A	145	ARG	NE-CZ-NH2	-8.51	116.04	120.30
1	A	258	ARG	NH1-CZ-NH2	7.77	127.95	119.40
1	A	57	ARG	NE-CZ-NH2	-7.47	116.56	120.30
1	A	57	ARG	NE-CZ-NH1	6.87	123.73	120.30
1	A	128	ASP	CB-CG-OD1	6.84	124.45	118.30
1	A	136	ARG	NE-CZ-NH1	6.73	123.66	120.30
1	A	179	ARG	NE-CZ-NH1	6.60	123.60	120.30
1	A	170	ASP	CB-CG-OD1	6.35	124.02	118.30
1	A	174	LEU	CB-CG-CD1	-6.31	100.27	111.00
1	A	147	PHE	CB-CG-CD2	-6.26	116.42	120.80
1	A	239	ASP	CB-CG-OD2	-6.02	112.88	118.30
1	A	166	ARG	NE-CZ-NH2	-6.01	117.30	120.30
1	A	258	ARG	CD-NE-CZ	-5.66	115.68	123.60
1	A	232	LEU	CB-CG-CD2	-5.43	101.77	111.00
1	A	152	LEU	CB-CG-CD2	5.39	120.17	111.00
1	A	249	ILE	CG1-CB-CG2	5.25	122.96	111.40
1	A	237	CYS	CA-CB-SG	-5.24	104.57	114.00
1	A	235	MET	CA-CB-CG	-5.23	104.41	113.30
1	A	136	ARG	CG-CD-NE	-5.06	101.17	111.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2225	0	2165	14	0
2	A	26	0	19	0	0
3	A	15	0	15	0	0
4	A	231	0	0	4	1
All	All	2497	0	2199	14	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124[A]:GLU:O	1:A:124[A]:GLU:HG3	1.71	0.86
1:A:287:HIS:HD2	1:A:289:TRP:H	1.25	0.83
1:A:287:HIS:CD2	1:A:289:TRP:H	2.05	0.75
1:A:74:ILE:HD13	1:A:87:PRO:HG3	1.75	0.69
1:A:262:GLU:HG3	4:A:1:HOH:O	1.95	0.65
1:A:265:HIS:ND1	1:A:287:HIS:HE1	1.94	0.64
1:A:97:SER:O	1:A:98:SER:HB3	2.02	0.58
1:A:292:ASP:HB3	4:A:331:HOH:O	2.04	0.56
1:A:80:LEU:O	1:A:82:ASN:N	2.41	0.52
1:A:105:ARG:NH1	4:A:444:HOH:O	2.35	0.51
1:A:81:PRO:O	1:A:82:ASN:HB3	2.10	0.49
1:A:213:ILE:HG22	1:A:249:ILE:HD11	1.94	0.48
1:A:292:ASP:CB	4:A:331:HOH:O	2.65	0.44
1:A:101:SER:O	1:A:183:LYS:NZ	2.40	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:376:HOH:O	4:A:396:HOH:O[2_544]	2.07	0.13

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	271/299 (91%)	263 (97%)	5 (2%)	3 (1%)	17   4

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	82	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	81	PRO
1	A	98	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	244/266 (92%)	236 (97%)	8 (3%)	43 21

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	ASP
1	A	79	GLU
1	A	84	THR
1	A	98	SER
1	A	101	SER
1	A	126[A]	VAL
1	A	126[B]	VAL
1	A	290	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	287	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	IMD	A	4496	-	3,5,5	0.69	0	4,5,5	0.59	0
3	IMD	A	4497	-	3,5,5	0.85	0	4,5,5	0.63	0
3	IMD	A	4498	-	3,5,5	0.64	0	4,5,5	0.93	0
2	UNQ	A	555	-	21,29,29	1.63	5 (23%)	28,41,41	2.10	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IMD	A	4496	-	-	0/0/0/0	0/1/1/1
3	IMD	A	4497	-	-	0/0/0/0	0/1/1/1
3	IMD	A	4498	-	-	0/0/0/0	0/1/1/1
2	UNQ	A	555	-	-	0/8/22/22	0/3/4/4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	555	UNQ	C03-C02	-2.13	1.37	1.42
2	A	555	UNQ	C21-C20	2.04	1.57	1.51
2	A	555	UNQ	C13-N15	2.23	1.38	1.34
2	A	555	UNQ	C25-C20	2.28	1.57	1.51
2	A	555	UNQ	C14-N15	3.06	1.40	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	555	UNQ	C14-N19-C20	-5.59	117.52	124.13
2	A	555	UNQ	C17-C13-C01	-3.93	116.57	122.03
2	A	555	UNQ	C01-C03-C02	-3.11	116.43	121.52
2	A	555	UNQ	C01-C13-N15	5.95	124.27	116.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	271/299 (90%)	0.16	14 (5%) <span style="border: 1px solid red; padding: 2px;">28</span> <span style="border: 1px solid red; padding: 2px;">31</span>	17, 29, 57, 79	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	80	LEU	7.6
1	A	83	GLY	5.9
1	A	81	PRO	5.9
1	A	84	THR	4.7
1	A	82	ASN	4.7
1	A	99	GLY	3.5
1	A	85	ARG	3.2
1	A	48	GLY	2.9
1	A	79	GLU	2.9
1	A	301	GLU	2.8
1	A	49	PHE	2.8
1	A	101	SER	2.5
1	A	305	HIS	2.3
1	A	100	PHE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	IMD	A	4497	5/5	0.92	0.26	14.61	20,20,20,20	0
2	UNQ	A	555	26/26	0.95	0.10	0.44	16,22,31,33	0
3	IMD	A	4496	5/5	0.89	0.26	-	20,20,20,20	0
3	IMD	A	4498	5/5	0.79	0.29	-	20,20,20,20	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.