



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 06:50 am GMT

PDB ID : 1R1L  
Title : Structure of dimeric antithrombin complexed with a P14-P9 reactive loop peptide and an exogenous tripeptide (formyl-norleucine-LF)  
Authors : Zhou, A.; Huntington, J.A.; Lomas, D.A.; Stein, P.E.; Carrell, R.W.  
Deposited on : 2003-09-24  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2259 (2.70-2.70)
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	I	432	<div> <div>3%</div> <div> <div></div> <div>63%</div> <div>29%</div> <div>• •</div> </div> </div>
1	L	432	<div> <div>4%</div> <div> <div></div> <div>50%</div> <div>41%</div> <div>• 5%</div> </div> </div>
2	C	7	<div> <div></div> <div> <div>86%</div> <div>14%</div> </div> </div>
3	D	3	<div> <div></div> <div> <div>33%</div> <div>67%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	I	901	X	-	-	-
4	NAG	I	941	-	-	X	-
4	NAG	I	942	X	-	-	-
4	NAG	I	962	X	-	-	-
4	NAG	L	841	X	-	X	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6898 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antithrombin-III.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	409	Total	C	N	O	S	0	0	0
			3270	2085	550	617	18			
1	I	416	Total	C	N	O	S	0	0	0
			3328	2121	562	627	18			

- Molecule 2 is a protein called Antithrombin P14-P9 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	7	Total	C	N	O	0	0	0
			40	22	6	12			

- Molecule 3 is a protein called EXOGENOUS TRIPEPTIDE formyl-(NLE)LF.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	3	Total	C	N	O	0	0	0
			28	21	3	4			

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	L	1	Total	C	N	O	0	0
			15	8	1	6		
4	L	1	Total	C	N	O	0	0
			15	8	1	6		
4	I	1	Total	C	N	O	0	0
			15	8	1	6		
4	I	1	Total	C	N	O	0	0
			15	8	1	6		
4	I	1	Total	C	N	O	0	0
			15	8	1	6		
4	I	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 5 is FORMYL GROUP (three-letter code: FOR) (formula: CH<sub>2</sub>O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			2	1	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	1	Total 1	O 1	0	0
7	I	76	Total 76	O 76	0	0
7	L	42	Total 42	O 42	0	0







- Molecule 2: Antithrombin P14-P9 peptide

Chain C:  86% 14%



- Molecule 3: EXOGENOUS TRIPEPTIDE formyl-(NLE)LF

Chain D:  33% 67%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.17Å 100.42Å 87.20Å 90.00° 104.29° 90.00°	Depositor
Resolution (Å)	24.99 – 2.70 24.99 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.5 (24.99-2.70) 99.6 (24.99-2.70)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	0.14	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.71 (at 2.72Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.204 , 0.249 0.203 , 0.251	Depositor DCC
$R_{free}$ test set	1602 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.7	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 63.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6898	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NLE, FOR, ACE, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	I	0.37	0/3394	0.61	0/4580
1	L	0.37	0/3334	0.61	1/4501 (0.0%)
2	C	0.46	0/37	0.46	0/48
3	D	0.75	0/20	0.84	0/24
All	All	0.38	0/6785	0.61	1/9153 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	316	LEU	CA-CB-CG	5.94	128.97	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	I	3328	0	3335	145	0
1	L	3270	0	3270	176	0
2	C	40	0	34	1	0
3	D	28	0	30	2	0
4	I	75	0	75	18	0
4	L	30	0	30	9	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	2	0	0	0	0
6	C	6	0	8	1	0
7	D	1	0	0	0	0
7	I	76	0	0	3	0
7	L	42	0	0	1	0
All	All	6898	0	6782	325	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 325 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:155:ASN:HD21	4:L:841:NAG:H1	1.14	1.13
1:L:155:ASN:HD21	4:L:841:NAG:C1	1.60	1.12
1:I:96:ASN:HD21	4:I:901:NAG:C1	1.62	1.11
1:L:155:ASN:ND2	4:L:841:NAG:H1	1.69	1.05
1:I:57:ARG:HB3	1:I:57:ARG:HH11	1.21	1.03

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	412/432 (95%)	381 (92%)	22 (5%)	9 (2%)	8	20
1	L	405/432 (94%)	364 (90%)	36 (9%)	5 (1%)	15	37
2	C	5/7 (71%)	5 (100%)	0	0	100	100
3	D	1/3 (33%)	1 (100%)	0	0	100	100
All	All	823/874 (94%)	751 (91%)	58 (7%)	14 (2%)	11	27

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	28	LYS
1	I	42	GLU
1	I	43	ALA
1	L	112	SER
1	L	242	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	I	370/383 (97%)	353 (95%)	17 (5%)	31	61
1	L	363/383 (95%)	341 (94%)	22 (6%)	22	47
2	C	3/3 (100%)	3 (100%)	0	100	100
3	D	2/2 (100%)	0	2 (100%)	0	0
All	All	738/771 (96%)	697 (94%)	41 (6%)	25	51

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	366	ASP
1	I	13	ARG
1	I	361	ASP
1	L	398	ASN
1	L	405	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	L	428	ASN
1	I	118	GLN
1	I	217	ASN
1	L	405	ASN
1	I	254	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NLE	D	10	3,5	7,7,8	0.91	0	4,7,9	1.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NLE	D	10	3,5	-	0/4/6/8	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	GOL	C	200	-	5,5,5	0.89	0	5,5,5	0.49	0
5	FOR	D	9	3	0,1,1	0.00	-	0,0,0	0.00	-
4	NAG	I	901	-	15,15,15	0.41	0	21,21,21	0.68	0
4	NAG	I	941	-	15,15,15	0.38	0	21,21,21	0.56	0
4	NAG	I	942	-	15,15,15	0.49	0	21,21,21	0.55	0
4	NAG	I	961	-	15,15,15	0.50	0	21,21,21	0.66	0
4	NAG	I	962	-	15,15,15	0.38	0	21,21,21	0.55	0
4	NAG	L	801	-	15,15,15	0.50	0	21,21,21	0.53	0
4	NAG	L	841	-	15,15,15	0.42	0	21,21,21	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	C	200	-	-	0/4/4/4	0/0/0/0
5	FOR	D	9	3	-	0/0/0/0	0/0/0/0
4	NAG	I	901	-	1/1/6/7	1/6/26/26	0/1/1/1
4	NAG	I	941	-	-	0/6/26/26	0/1/1/1
4	NAG	I	942	-	1/1/6/7	0/6/26/26	0/1/1/1
4	NAG	I	961	-	-	0/6/26/26	0/1/1/1
4	NAG	I	962	-	1/1/6/7	0/6/26/26	0/1/1/1
4	NAG	L	801	-	-	0/6/26/26	0/1/1/1
4	NAG	L	841	-	1/1/6/7	0/6/26/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	L	841	NAG	C1
4	I	962	NAG	C1
4	I	901	NAG	C1
4	I	942	NAG	C1

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	I	901	NAG	O7-C7-N2-C2

There are no ring outliers.

8 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	200	GOL	1	0
4	I	901	NAG	4	0
4	I	941	NAG	7	0
4	I	942	NAG	4	0
4	I	961	NAG	6	0
4	I	962	NAG	1	0
4	L	801	NAG	1	0
4	L	841	NAG	8	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	I	416/432 (96%)	-0.17	12 (2%) 52 52	14, 36, 88, 196	0
1	L	409/432 (94%)	0.09	19 (4%) 33 31	18, 43, 114, 191	0
2	C	6/7 (85%)	-0.41	0 100 100	19, 28, 32, 36	0
3	D	2/3 (66%)	0.33	0 100 100	35, 35, 35, 43	0
All	All	833/874 (95%)	-0.04	31 (3%) 42 41	14, 39, 104, 196	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	134	ALA	6.2
1	L	431	VAL	6.0
1	I	358	GLY	5.6
1	L	135	ASN	5.4
1	I	357	GLU	5.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NLE	D	10	8/9	0.94	0.24	-	23,25,29,30	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	L	841	15/15	0.76	0.28	2.42	44,75,85,94	0
6	GOL	C	200	6/6	0.91	0.27	1.33	45,45,45,45	0
4	NAG	I	941	15/15	0.88	0.20	-0.01	54,54,54,54	0
4	NAG	L	801	15/15	0.70	0.39	-	48,103,112,121	0
5	FOR	D	9	2/2	0.97	0.25	-	33,33,33,33	0
4	NAG	I	961	15/15	0.64	0.32	-	88,88,88,88	0
4	NAG	I	962	15/15	0.64	0.44	-	109,109,109,109	0
4	NAG	I	901	15/15	0.83	0.25	-	66,66,66,66	0
4	NAG	I	942	15/15	0.54	0.52	-	143,143,143,143	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.