



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 15, 2017 – 06:53 AM EDT

PDB ID : 2R17
Title : Functional architecture of the retromer cargo-recognition complex
Authors : Hierro, A.; Rojas, A.L.; Rojas, R.; Murthy, N.; Effantin, G.; Kajava, A.V.;
Steven, A.C.; Bonifacino, J.S.; Hurley, J.H.
Deposited on : unknown
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	183	<div> <div>3%</div> <div>52%</div> <div>41%</div> <div>7%</div> </div>
1	B	183	<div> <div>56%</div> <div>38%</div> <div>5%</div> </div>
2	C	298	<div> <div>%</div> <div>52%</div> <div>43%</div> <div>5%</div> </div>
2	D	298	<div> <div>%</div> <div>53%</div> <div>37%</div> <div>• 7%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7640 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vacuolar protein sorting-associated protein 29.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	183	Total	C	N	O	S	Se	0	0	0
			1455	941	243	264	3	4			
1	B	183	Total	C	N	O	S	Se	0	0	0
			1455	941	243	264	3	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MSE	-	INITIATING METHIONINE	UNP Q9UBQ0
B	0	MSE	-	INITIATING METHIONINE	UNP Q9UBQ0

- Molecule 2 is a protein called Vacuolar protein sorting-associated protein 35.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	298	Total	C	N	O	S	Se	0	0	0
			2416	1534	422	450	6	4			
2	D	276	Total	C	N	O	S	Se	0	0	0
			2253	1442	390	411	6	4			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	8	Total	O	0	0
			8	8		
4	B	10	Total	O	0	0
			10	10		
4	C	17	Total	O	0	0
			17	17		
4	D	14	Total	O	0	0
			14	14		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vacuolar protein sorting-associated protein 29



- Molecule 1: Vacuolar protein sorting-associated protein 29



- Molecule 2: Vacuolar protein sorting-associated protein 35



N744
Q745
L746
L747
Q748
K749
L750
L754
P755
S759
S760
E761
E762
E764
Q765
H769
T773
L777
R778
L779
R780

• Molecule 2: Vacuolar protein sorting-associated protein 35

Chain D: 53% 37% 7%

ASP
F484
A485
D486
E487
L490
R493
F494
L497
L498
R499
S500
E501
D502
P503
D504
Q505
L508
T512
A513
R514
P515
H516
F517
G520
Q523
R524
I525
T528
L529
P530
P531
R542
Y543
K544
E545
H546
S547
K556
I560
F561
S562
F563
A564
H565
Q566
T567
I568
K573
A574
E575
L576
A577
E578
L579
P580
L581
R582
L583
F584
L585
L589
E593
I594
E597
N598
E605
F606
M607
S608
Q609
A610
F611
L625
T629
R630
T634
F635
E636
R637
M638
K639
M645
H646
E647
P648
L649
R650
T651
L660
P664
R668
T672
M678
SER
GLY
ARG
ASN
THR
ASP
LYS
ASN
GLY
GLU
GLU
LEU
HIS
GLY
GLY
LYS
R695
V696
M697
L700
K701
K702
A703
L704
K705
I706
A707
P713
S714
L715
Q716
V717
Q718
L719
F720
I721
E722
I723
L724
N725
R726
Y727
I728
E732
K733
GLU
ASN
ASP
ALA
VAL
I739
I740
I746
I750
L754
P755
E758
S759
S760
E761
E762
T763
E764
Q765
K768
H769
F770
H771
N772
T773
L774
E775
R778
L779
R780

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.25Å 128.47Å 140.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 48.19 – 2.80	Depositor EDS
% Data completeness (in resolution range)	92.1 (50.00-2.80) 98.8 (48.19-2.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.09 (at 2.81Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.216 , 0.268 0.226 , 0.237	Depositor DCC
R_{free} test set	1509 reflections (5.05%)	DCC
Wilson B-factor (Å ²)	53.5	Xtriage
Anisotropy	0.702	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 61.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7640	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/1486	0.68	0/2009
1	B	0.41	0/1486	0.67	1/2009 (0.0%)
2	C	0.41	0/2459	0.60	1/3303 (0.0%)
2	D	0.41	0/2293	0.61	0/3078
All	All	0.40	0/7724	0.63	2/10399 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	547	SER	N-CA-C	5.58	126.07	111.00
1	B	40	LEU	N-CA-C	-5.35	96.55	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1455	0	1468	89	0
1	B	1455	0	1468	63	0
2	C	2416	0	2401	127	0
2	D	2253	0	2256	126	0
3	D	12	0	16	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	8	0	0	1	0
4	B	10	0	0	1	0
4	C	17	0	0	2	0
4	D	14	0	0	6	0
All	All	7640	0	7609	393	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 393 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:525:ILE:HA	2:D:528:THR:HG22	1.28	1.14
2:D:520:GLY:HA3	2:D:524:ARG:HD3	1.15	1.14
1:A:8:ASP:H	1:A:38:GLY:HA3	1.15	1.10
2:D:635:PHE:HA	2:D:638:MSE:HE2	1.37	1.02
2:C:712:ASP:O	2:C:716:GLN:HB2	1.63	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/183 (99%)	158 (87%)	22 (12%)	1 (1%)	28	62
1	B	181/183 (99%)	172 (95%)	7 (4%)	2 (1%)	17	47
2	C	296/298 (99%)	277 (94%)	18 (6%)	1 (0%)	44	77
2	D	270/298 (91%)	248 (92%)	21 (8%)	1 (0%)	38	72
All	All	928/962 (96%)	855 (92%)	68 (7%)	5 (0%)	32	67

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	710	CYS
2	D	520	GLY
1	A	38	GLY
1	B	41	CYS
1	B	157	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	161/157 (102%)	150 (93%)	11 (7%)	18	47
1	B	161/157 (102%)	148 (92%)	13 (8%)	14	37
2	C	259/255 (102%)	240 (93%)	19 (7%)	16	42
2	D	242/255 (95%)	229 (95%)	13 (5%)	26	58
All	All	823/824 (100%)	767 (93%)	56 (7%)	18	47

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	498	LEU
2	C	555	LYS
2	D	630	LEU
2	C	505	GLN
2	C	528	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	D	1	-	5,5,5	0.33	0	5,5,5	0.28	0
3	GOL	D	2	-	5,5,5	0.32	0	5,5,5	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	D	1	-	-	0/4/4/4	0/0/0/0
3	GOL	D	2	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	179/183 (97%)	0.07	6 (3%) 46 34	24, 62, 91, 105	0
1	B	179/183 (97%)	-0.28	0 100 100	24, 46, 65, 75	0
2	C	294/298 (98%)	-0.19	3 (1%) 82 77	26, 55, 89, 104	0
2	D	272/298 (91%)	-0.17	2 (0%) 87 83	21, 48, 86, 100	0
All	All	924/962 (96%)	-0.15	11 (1%) 79 72	21, 52, 86, 105	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	155	ILE	2.9
1	A	182	PRO	2.8
1	A	31	ILE	2.4
2	D	780	ARG	2.2
2	C	547	SER	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	D	2	6/6	0.84	0.27	-	68,72,73,73	0
3	GOL	D	1	6/6	0.91	0.18	-	78,79,80,81	0

6.5 Other polymers [i](#)

There are no such residues in this entry.