



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 31, 2017 – 05:53 PM EDT

PDB ID : 3R1C
Title : Crystal structure of GCGGCGGC duplex
Authors : Kiliszek, A.; Kierzek, R.; Krzyzosiak, W.J.; Rypniewski, W.
Deposited on : unknown
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

| | | |
|--------------------------------|---|--|
| MolProbity | : | 4.02b-467 |
| Mogul | : | 1.7.2 (RC1), CSD as538be (2017) |
| Xtriage (Phenix) | : | 1.9-1692 |
| EDS | : | rb-20030345 |
| Percentile statistics | : | 20161228.v01 (using entries in the PDB archive December 28th 2016) |
| Refmac | : | 5.8.0135 |
| CCP4 | : | 6.5.0 |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | rb-20030345 |


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|---------------|-----------------------------|---|
| R_{free} | 100719 | 2028 (2.08-2.04) |
| Clashscore | 112137 | 2143 (2.08-2.04) |
| RSRZ outliers | 101464 | 2035 (2.08-2.04) |
| RNA backbone | 2435 | 1078 (2.70-1.42) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 1 | A | 8 | <div> <div></div> <div>75%25%</div> </div> |
| 1 | B | 8 | <div> <div></div> <div>100%</div> </div> |
| 1 | C | 8 | <div> <div></div> <div>88%13%</div> </div> |
| 1 | D | 8 | <div> <div></div> <div>75%25%</div> </div> |
| 1 | E | 8 | <div> <div></div> <div>75%25%</div> </div> |
| 1 | F | 8 | <div> <div></div> <div>75%25%</div> </div> |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|---|
| 1 | G | 8 |  75%25% |
| 1 | H | 8 |  88%13% |
| 1 | I | 8 |  88%13% |
| 1 | J | 8 |  100% |
| 1 | K | 8 |  88%13% |
| 1 | L | 8 |  100% |
| 1 | M | 8 |  100% |
| 1 | N | 8 |  100% |
| 1 | O | 8 |  88%13% |
| 1 | P | 8 |  88%13% |
| 1 | Q | 8 |  88%13% |
| 1 | R | 8 |  100% |
| 1 | S | 8 |  88%13% |
| 1 | T | 8 |  100% |
| 1 | U | 8 |  100% |
| 1 | V | 8 |  88%13% |
| 1 | W | 8 |  88%13% |
| 1 | X | 8 |  88%13% |
| 1 | Y | 8 |  63%38% |
| 1 | Z | 8 |  88%13% |
| 1 | a | 8 |  100% |
| 1 | b | 8 |  100% |
| 1 | c | 8 |  100% |
| 1 | d | 8 |  100% |
| 1 | e | 8 |  100% |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|---|
| 1 | f | 8 |  100% |
| 1 | g | 8 |  100% |
| 1 | h | 8 |  88%13% |
| 1 | i | 8 |  100% |
| 1 | j | 8 |  100% |

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6865 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3').

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|----|----|----|---|---------|---------|-------|
| 1 | A | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | B | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | C | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | D | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | E | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | F | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | G | 8 | Total | C | N | O | P | 0 | 1 | 0 |
| | | | 195 | 87 | 39 | 61 | 8 | | | |
| 1 | H | 8 | Total | C | N | O | P | 0 | 1 | 0 |
| | | | 195 | 87 | 39 | 61 | 8 | | | |
| 1 | I | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | J | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | K | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | L | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | M | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | N | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | O | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |
| 1 | P | 8 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 172 | 77 | 34 | 54 | 7 | | | |

Continued on next page...

Continued from previous page...

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|--------------|---------|---------|---------|--------|---------|---------|-------|
| 1 | Q | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | R | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | S | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | Y | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | T | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | U | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | W | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | X | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | V | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | Z | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | a | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | b | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | c | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | d | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | e | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | f | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | g | 8 | Total 215 | C 96 | N 42 | O 68 | P 9 | 0 | 2 | 0 |
| 1 | h | 8 | Total 195 | C 87 | N 39 | O 61 | P 8 | 0 | 1 | 0 |
| 1 | i | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |
| 1 | j | 8 | Total 172 | C 77 | N 34 | O 54 | P 7 | 0 | 0 | 0 |

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---------|---------|
| 2 | S | 1 | Total | O | S | 0 | 0 |
| | | | 5 | 4 | 1 | | |
| 2 | W | 1 | Total | O | S | 0 | 1 |
| | | | 5 | 4 | 1 | | |
| 2 | X | 1 | Total | O | S | 0 | 0 |
| | | | 5 | 4 | 1 | | |
| 2 | b | 1 | Total | O | S | 0 | 1 |
| | | | 5 | 4 | 1 | | |
| 2 | c | 1 | Total | O | S | 0 | 1 |
| | | | 5 | 4 | 1 | | |
| 2 | d | 1 | Total | O | S | 0 | 0 |
| | | | 5 | 4 | 1 | | |
| 2 | e | 1 | Total | O | S | 0 | 0 |
| | | | 5 | 4 | 1 | | |

- Molecule 3 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 3 | A | 21 | Total | O | 0 | 0 |
| | | | 21 | 21 | | |
| 3 | B | 19 | Total | O | 0 | 0 |
| | | | 19 | 19 | | |
| 3 | C | 19 | Total | O | 0 | 0 |
| | | | 19 | 19 | | |
| 3 | D | 13 | Total | O | 0 | 0 |
| | | | 13 | 13 | | |
| 3 | E | 20 | Total | O | 0 | 0 |
| | | | 20 | 20 | | |

Continued on next page...

Continued from previous page...

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------------|---------|---------|---------|
| 3 | F | 13 | Total 13 | O 13 | 0 | 0 |
| 3 | G | 18 | Total 18 | O 18 | 0 | 0 |
| 3 | H | 10 | Total 10 | O 10 | 0 | 1 |
| 3 | I | 29 | Total 29 | O 29 | 0 | 0 |
| 3 | J | 28 | Total 28 | O 28 | 0 | 0 |
| 3 | K | 29 | Total 29 | O 29 | 0 | 0 |
| 3 | L | 26 | Total 26 | O 26 | 0 | 0 |
| 3 | M | 17 | Total 17 | O 17 | 0 | 0 |
| 3 | N | 6 | Total 6 | O 6 | 0 | 0 |
| 3 | O | 12 | Total 12 | O 12 | 0 | 0 |
| 3 | P | 12 | Total 12 | O 12 | 0 | 0 |
| 3 | Q | 23 | Total 23 | O 23 | 0 | 0 |
| 3 | R | 15 | Total 15 | O 15 | 0 | 0 |
| 3 | S | 2 | Total 2 | O 2 | 0 | 0 |
| 3 | Y | 3 | Total 3 | O 3 | 0 | 0 |
| 3 | T | 10 | Total 10 | O 10 | 0 | 0 |
| 3 | U | 19 | Total 19 | O 19 | 0 | 0 |
| 3 | W | 11 | Total 11 | O 11 | 0 | 0 |
| 3 | X | 18 | Total 19 | O 19 | 0 | 1 |
| 3 | V | 4 | Total 4 | O 4 | 0 | 0 |
| 3 | Z | 8 | Total 8 | O 8 | 0 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------------|---------|---------|---------|
| 3 | a | 3 | Total 3 | O 3 | 0 | 0 |
| 3 | b | 4 | Total 4 | O 4 | 0 | 0 |
| 3 | c | 13 | Total 13 | O 13 | 0 | 0 |
| 3 | d | 14 | Total 14 | O 14 | 0 | 0 |
| 3 | e | 22 | Total 22 | O 22 | 0 | 0 |
| 3 | f | 17 | Total 17 | O 17 | 0 | 0 |
| 3 | g | 9 | Total 9 | O 9 | 0 | 0 |
| 3 | h | 10 | Total 10 | O 10 | 0 | 0 |
| 3 | i | 17 | Total 18 | O 18 | 0 | 1 |
| 3 | j | 10 | Total 10 | O 10 | 0 | 0 |

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain A: 




- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain B: 

There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain C: 



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain D: 



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain E: 




- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')


Chain F: 




- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')



There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')



There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')




There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')




There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain O:  88% 13%




- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain P:  88% 13%



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain Q:  88% 13%





- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain R:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain S:  88% 13%



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain Y:  63% 38%



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain T:  100%


There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain U:  100%


There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain W:  88% 13%




- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain X:  88% 13%




- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain V:  88% 13%



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain Z:  88% 13%



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain a:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain b:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain c:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain d:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain e:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain f:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain g:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain h:  88% 13%



- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain i:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: RNA (5'-R(*GP*CP*GP*GP*CP*GP*GP*C)-3')

Chain j:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 1 | Depositor |
| Cell constants a, b, c, α , β , γ | 39.70Å 76.89Å 85.40Å 89.98° 88.61° 77.29° | Depositor |
| Resolution (Å) | 19.36 – 2.05 19.83 – 2.05 | Depositor EDS |
| % Data completeness (in resolution range) | 97.7 (19.36-2.05) 98.0 (19.83-2.05) | Depositor EDS |
| R_{merge} | (Not available) | Depositor |
| R_{sym} | 0.09 | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 3.12 (at 2.06Å) | Xtriage |
| Refinement program | PHENIX | Depositor |
| R, R_{free} | 0.216 , 0.257 0.203 , 0.247 | Depositor DCC |
| R_{free} test set | 3037 reflections (5.06%) | DCC |
| Wilson B-factor (Å ²) | 28.5 | Xtriage |
| Anisotropy | 0.078 | Xtriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.32 , 48.2 | EDS |
| L-test for twinning ² | $\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$ | Xtriage |
| Estimated twinning fraction | 0.000 for h,h-k,-l 0.000 for -h,-k,l 0.000 for -h,-h+k,-l | Xtriage |
| F_o, F_c correlation | 0.96 | EDS |
| Total number of atoms | 6865 | wwPDB-VP |
| Average B, all atoms (Å ²) | 26.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 69.39 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.7299e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------|-------------|--------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | A | 0.60 | 0/192 | 1.14 | 0/299 |
| 1 | B | 0.61 | 0/192 | 1.17 | 0/299 |
| 1 | C | 0.51 | 0/192 | 1.07 | 0/299 |
| 1 | D | 0.60 | 0/192 | 1.14 | 1/299 (0.3%) |
| 1 | E | 0.61 | 0/192 | 1.13 | 0/299 |
| 1 | F | 0.62 | 0/192 | 1.25 | 0/299 |
| 1 | G | 0.58 | 0/218 | 1.18 | 0/340 |
| 1 | H | 0.46 | 0/218 | 1.04 | 0/340 |
| 1 | I | 0.66 | 0/192 | 1.18 | 0/299 |
| 1 | J | 0.64 | 0/192 | 1.08 | 0/299 |
| 1 | K | 0.70 | 0/192 | 1.26 | 1/299 (0.3%) |
| 1 | L | 0.63 | 0/192 | 1.06 | 0/299 |
| 1 | M | 0.57 | 0/192 | 1.22 | 0/299 |
| 1 | N | 0.55 | 0/192 | 1.13 | 0/299 |
| 1 | O | 0.57 | 0/192 | 1.07 | 0/299 |
| 1 | P | 0.60 | 0/192 | 1.19 | 0/299 |
| 1 | Q | 0.61 | 0/192 | 1.18 | 1/299 (0.3%) |
| 1 | R | 0.62 | 0/192 | 1.06 | 0/299 |
| 1 | S | 0.49 | 0/192 | 0.96 | 0/299 |
| 1 | T | 0.51 | 0/192 | 1.12 | 0/299 |
| 1 | U | 0.57 | 0/192 | 1.20 | 0/299 |
| 1 | V | 0.49 | 0/192 | 1.04 | 0/299 |
| 1 | W | 0.55 | 0/192 | 1.15 | 0/299 |
| 1 | X | 0.57 | 0/192 | 1.12 | 0/299 |
| 1 | Y | 0.49 | 0/192 | 1.05 | 0/299 |
| 1 | Z | 0.49 | 0/192 | 1.07 | 0/299 |
| 1 | a | 0.45 | 0/192 | 1.09 | 0/299 |
| 1 | b | 0.47 | 0/192 | 0.95 | 0/299 |
| 1 | c | 0.50 | 0/192 | 1.12 | 0/299 |
| 1 | d | 0.60 | 0/192 | 1.14 | 0/299 |
| 1 | e | 0.59 | 0/192 | 1.17 | 0/299 |
| 1 | f | 0.56 | 0/192 | 1.07 | 0/299 |

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------|-------------|----------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | g | 0.56 | 0/240 | 1.04 | 0/374 |
| 1 | h | 0.60 | 0/218 | 1.12 | 1/340 (0.3%) |
| 1 | i | 0.64 | 0/192 | 1.08 | 0/299 |
| 1 | j | 0.56 | 0/192 | 1.02 | 0/299 |
| All | All | 0.57 | 0/7038 | 1.11 | 4/10962 (0.0%) |

There are no bond length outliers.

All (4) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|------------|------|-------------|----------|
| 1 | K | 3 | G | O4'-C1'-N9 | 6.59 | 113.47 | 108.20 |
| 1 | Q | 6 | G | O4'-C1'-N9 | 5.62 | 112.70 | 108.20 |
| 1 | D | 6 | G | O4'-C1'-N9 | 5.32 | 112.45 | 108.20 |
| 1 | h | 6 | G | O4'-C1'-N9 | 5.04 | 112.23 | 108.20 |

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 172 | 0 | 90 | 1 | 0 |
| 1 | B | 172 | 0 | 90 | 0 | 0 |
| 1 | C | 172 | 0 | 90 | 1 | 0 |
| 1 | D | 172 | 0 | 90 | 1 | 0 |
| 1 | E | 172 | 0 | 90 | 2 | 0 |
| 1 | F | 172 | 0 | 90 | 2 | 0 |
| 1 | G | 195 | 0 | 101 | 4 | 0 |
| 1 | H | 195 | 0 | 101 | 2 | 0 |
| 1 | I | 172 | 0 | 90 | 2 | 0 |
| 1 | J | 172 | 0 | 90 | 0 | 0 |
| 1 | K | 172 | 0 | 90 | 0 | 0 |
| 1 | L | 172 | 0 | 90 | 0 | 0 |
| 1 | M | 172 | 0 | 90 | 0 | 0 |
| 1 | N | 172 | 0 | 90 | 0 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | O | 172 | 0 | 90 | 1 | 0 |
| 1 | P | 172 | 0 | 90 | 1 | 0 |
| 1 | Q | 172 | 0 | 90 | 0 | 0 |
| 1 | R | 172 | 0 | 90 | 0 | 0 |
| 1 | S | 172 | 0 | 90 | 1 | 0 |
| 1 | T | 172 | 0 | 90 | 0 | 0 |
| 1 | U | 172 | 0 | 90 | 0 | 0 |
| 1 | V | 172 | 0 | 90 | 2 | 0 |
| 1 | W | 172 | 0 | 90 | 1 | 0 |
| 1 | X | 172 | 0 | 90 | 1 | 0 |
| 1 | Y | 172 | 0 | 90 | 2 | 0 |
| 1 | Z | 172 | 0 | 90 | 1 | 0 |
| 1 | a | 172 | 0 | 90 | 0 | 0 |
| 1 | b | 172 | 0 | 90 | 0 | 0 |
| 1 | c | 172 | 0 | 90 | 0 | 0 |
| 1 | d | 172 | 0 | 90 | 0 | 0 |
| 1 | e | 172 | 0 | 90 | 0 | 0 |
| 1 | f | 172 | 0 | 90 | 0 | 0 |
| 1 | g | 215 | 0 | 110 | 0 | 0 |
| 1 | h | 195 | 0 | 99 | 0 | 0 |
| 1 | i | 172 | 0 | 90 | 0 | 0 |
| 1 | j | 172 | 0 | 90 | 0 | 0 |
| 2 | S | 5 | 0 | 0 | 0 | 0 |
| 2 | W | 5 | 0 | 0 | 0 | 0 |
| 2 | X | 5 | 0 | 0 | 0 | 0 |
| 2 | b | 5 | 0 | 0 | 0 | 0 |
| 2 | c | 5 | 0 | 0 | 0 | 0 |
| 2 | d | 5 | 0 | 0 | 0 | 0 |
| 2 | e | 5 | 0 | 0 | 0 | 0 |
| 3 | A | 21 | 0 | 0 | 0 | 0 |
| 3 | B | 19 | 0 | 0 | 0 | 0 |
| 3 | C | 19 | 0 | 0 | 0 | 0 |
| 3 | D | 13 | 0 | 0 | 0 | 0 |
| 3 | E | 20 | 0 | 0 | 0 | 0 |
| 3 | F | 13 | 0 | 0 | 0 | 0 |
| 3 | G | 18 | 0 | 0 | 1 | 0 |
| 3 | H | 10 | 0 | 0 | 0 | 0 |
| 3 | I | 29 | 0 | 0 | 0 | 0 |
| 3 | J | 28 | 0 | 0 | 0 | 0 |
| 3 | K | 29 | 0 | 0 | 0 | 0 |
| 3 | L | 26 | 0 | 0 | 0 | 0 |
| 3 | M | 17 | 0 | 0 | 0 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 3 | N | 6 | 0 | 0 | 0 | 0 |
| 3 | O | 12 | 0 | 0 | 0 | 0 |
| 3 | P | 12 | 0 | 0 | 0 | 0 |
| 3 | Q | 23 | 0 | 0 | 0 | 0 |
| 3 | R | 15 | 0 | 0 | 0 | 0 |
| 3 | S | 2 | 0 | 0 | 0 | 0 |
| 3 | T | 10 | 0 | 0 | 0 | 0 |
| 3 | U | 19 | 0 | 0 | 0 | 0 |
| 3 | V | 4 | 0 | 0 | 0 | 0 |
| 3 | W | 11 | 0 | 0 | 0 | 0 |
| 3 | X | 19 | 0 | 0 | 0 | 0 |
| 3 | Y | 3 | 0 | 0 | 0 | 0 |
| 3 | Z | 8 | 0 | 0 | 1 | 0 |
| 3 | a | 3 | 0 | 0 | 0 | 0 |
| 3 | b | 4 | 0 | 0 | 0 | 0 |
| 3 | c | 13 | 0 | 0 | 0 | 0 |
| 3 | d | 14 | 0 | 0 | 0 | 0 |
| 3 | e | 22 | 0 | 0 | 0 | 0 |
| 3 | f | 17 | 0 | 0 | 0 | 0 |
| 3 | g | 9 | 0 | 0 | 0 | 0 |
| 3 | h | 10 | 0 | 0 | 0 | 0 |
| 3 | i | 18 | 0 | 0 | 0 | 0 |
| 3 | j | 10 | 0 | 0 | 0 | 0 |
| All | All | 6865 | 0 | 3291 | 21 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 21 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|---------------|---------------|--------------------------|-------------------|
| 1:G:3[A]:G:H8 | 3:G:102:HOH:O | 1.71 | 0.73 |
| 1:F:1:G:HO5' | 1:F:1:G:H8 | 3.06 | 0.60 |
| 1:V:1:G:H8 | 1:V:1:G:HO5' | 1.51 | 0.56 |
| 1:W:3:G:H5'' | 1:W:3:G:N3 | 2.22 | 0.55 |
| 1:P:1:G:H8 | 1:P:1:G:HO5' | 1.55 | 0.54 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

| Mol | Chain | Analysed | Backbone Outliers | Pucker Outliers |
|-----|-------|-----------|-------------------|-----------------|
| 1 | A | 7/8 (87%) | 0 | 0 |
| 1 | B | 7/8 (87%) | 0 | 0 |
| 1 | C | 7/8 (87%) | 0 | 0 |
| 1 | D | 7/8 (87%) | 0 | 0 |
| 1 | E | 7/8 (87%) | 0 | 0 |
| 1 | F | 7/8 (87%) | 0 | 0 |
| 1 | G | 6/8 (75%) | 0 | 0 |
| 1 | H | 6/8 (75%) | 0 | 0 |
| 1 | I | 7/8 (87%) | 0 | 0 |
| 1 | J | 7/8 (87%) | 0 | 0 |
| 1 | K | 7/8 (87%) | 0 | 0 |
| 1 | L | 7/8 (87%) | 0 | 0 |
| 1 | M | 7/8 (87%) | 0 | 0 |
| 1 | N | 7/8 (87%) | 0 | 0 |
| 1 | O | 7/8 (87%) | 0 | 0 |
| 1 | P | 7/8 (87%) | 0 | 0 |
| 1 | Q | 7/8 (87%) | 0 | 0 |
| 1 | R | 7/8 (87%) | 0 | 0 |
| 1 | S | 7/8 (87%) | 0 | 0 |
| 1 | T | 7/8 (87%) | 0 | 0 |
| 1 | U | 7/8 (87%) | 0 | 0 |
| 1 | V | 7/8 (87%) | 0 | 0 |
| 1 | W | 7/8 (87%) | 0 | 0 |
| 1 | X | 7/8 (87%) | 0 | 0 |
| 1 | Y | 7/8 (87%) | 0 | 0 |
| 1 | Z | 7/8 (87%) | 0 | 0 |
| 1 | a | 7/8 (87%) | 0 | 0 |
| 1 | b | 7/8 (87%) | 0 | 0 |
| 1 | c | 7/8 (87%) | 0 | 0 |
| 1 | d | 7/8 (87%) | 0 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Backbone Outliers | Pucker Outliers |
|-----|-------|---------------|-------------------|-----------------|
| 1 | e | 7/8 (87%) | 0 | 0 |
| 1 | f | 7/8 (87%) | 0 | 0 |
| 1 | g | 5/8 (62%) | 0 | 0 |
| 1 | h | 6/8 (75%) | 0 | 0 |
| 1 | i | 7/8 (87%) | 0 | 0 |
| 1 | j | 7/8 (87%) | 0 | 0 |
| All | All | 247/288 (85%) | 0 | 0 |

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|------|------|--------------|------|-------------|-------------|------|-------------|
| | | | | | Counts | RMSZ | $\# Z > 2$ | Counts | RMSZ | $\# Z > 2$ |
| 2 | SO4 | S | 9 | - | 4,4,4 | 0.14 | 0 | 6,6,6 | 0.07 | 0 |
| 2 | SO4 | W | 9[A] | - | 4,4,4 | 0.20 | 0 | 6,6,6 | 0.12 | 0 |
| 2 | SO4 | X | 9 | - | 4,4,4 | 0.13 | 0 | 6,6,6 | 0.08 | 0 |
| 2 | SO4 | b | 9[A] | - | 4,4,4 | 0.16 | 0 | 6,6,6 | 0.08 | 0 |
| 2 | SO4 | c | 9[A] | - | 4,4,4 | 0.17 | 0 | 6,6,6 | 0.12 | 0 |
| 2 | SO4 | d | 9 | - | 4,4,4 | 0.15 | 0 | 6,6,6 | 0.12 | 0 |
| 2 | SO4 | e | 9 | - | 4,4,4 | 0.19 | 0 | 6,6,6 | 0.12 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|------|------|---------|----------|---------|
| 2 | SO4 | S | 9 | - | - | 0/0/0/0 | 0/0/0/0 |
| 2 | SO4 | W | 9[A] | - | - | 0/0/0/0 | 0/0/0/0 |
| 2 | SO4 | X | 9 | - | - | 0/0/0/0 | 0/0/0/0 |
| 2 | SO4 | b | 9[A] | - | - | 0/0/0/0 | 0/0/0/0 |
| 2 | SO4 | c | 9[A] | - | - | 0/0/0/0 | 0/0/0/0 |
| 2 | SO4 | d | 9 | - | - | 0/0/0/0 | 0/0/0/0 |
| 2 | SO4 | e | 9 | - | - | 0/0/0/0 | 0/0/0/0 |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|------------|--------|-----------|-----------------------|-------|
| 1 | A | 8/8 (100%) | -0.62 | 0 100 100 | 17, 19, 22, 24 | 0 |
| 1 | B | 8/8 (100%) | -0.71 | 0 100 100 | 17, 21, 23, 27 | 0 |
| 1 | C | 8/8 (100%) | -0.38 | 0 100 100 | 20, 22, 41, 43 | 0 |
| 1 | D | 8/8 (100%) | -0.40 | 0 100 100 | 26, 27, 27, 30 | 0 |
| 1 | E | 8/8 (100%) | -0.67 | 0 100 100 | 17, 19, 24, 25 | 0 |
| 1 | F | 8/8 (100%) | -0.62 | 0 100 100 | 17, 21, 25, 27 | 0 |
| 1 | G | 8/8 (100%) | -0.38 | 0 100 100 | 23, 26, 27, 29 | 0 |
| 1 | H | 8/8 (100%) | -0.44 | 0 100 100 | 26, 27, 33, 34 | 0 |
| 1 | I | 8/8 (100%) | -0.77 | 0 100 100 | 13, 14, 19, 19 | 0 |
| 1 | J | 8/8 (100%) | -0.80 | 0 100 100 | 13, 15, 18, 18 | 0 |
| 1 | K | 8/8 (100%) | -0.83 | 0 100 100 | 13, 14, 17, 19 | 0 |
| 1 | L | 8/8 (100%) | -0.79 | 0 100 100 | 13, 15, 17, 17 | 0 |
| 1 | M | 8/8 (100%) | -0.77 | 0 100 100 | 17, 18, 32, 34 | 0 |
| 1 | N | 8/8 (100%) | -0.70 | 0 100 100 | 18, 31, 35, 41 | 0 |
| 1 | O | 8/8 (100%) | -0.65 | 0 100 100 | 22, 25, 30, 35 | 0 |
| 1 | P | 8/8 (100%) | -0.60 | 0 100 100 | 21, 24, 38, 40 | 0 |
| 1 | Q | 8/8 (100%) | -0.84 | 0 100 100 | 17, 18, 25, 26 | 0 |
| 1 | R | 8/8 (100%) | -0.84 | 0 100 100 | 18, 24, 26, 31 | 0 |
| 1 | S | 8/8 (100%) | -0.20 | 0 100 100 | 29, 43, 51, 52 | 0 |
| 1 | T | 8/8 (100%) | -0.73 | 0 100 100 | 19, 26, 32, 33 | 0 |
| 1 | U | 8/8 (100%) | -0.77 | 0 100 100 | 20, 22, 27, 30 | 0 |
| 1 | V | 8/8 (100%) | -0.24 | 0 100 100 | 32, 36, 38, 40 | 0 |
| 1 | W | 8/8 (100%) | -0.76 | 0 100 100 | 22, 25, 25, 27 | 0 |
| 1 | X | 8/8 (100%) | -0.89 | 0 100 100 | 22, 25, 30, 30 | 0 |

Continued on next page...

Continued from previous page...

Continued from previous page.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 | | |
|-----|-------|----------------|--------|---------|-----------------------|-------|----------------|---|
| 1 | Y | 8/8 (100%) | 0.01 | 0 | 100 | 100 | 27, 46, 53, 53 | 0 |
| 1 | Z | 8/8 (100%) | -0.63 | 0 | 100 | 100 | 29, 30, 32, 35 | 0 |
| 1 | a | 8/8 (100%) | -0.16 | 0 | 100 | 100 | 25, 44, 47, 49 | 0 |
| 1 | b | 8/8 (100%) | -0.48 | 0 | 100 | 100 | 26, 37, 44, 45 | 0 |
| 1 | c | 8/8 (100%) | -0.75 | 0 | 100 | 100 | 21, 23, 26, 27 | 0 |
| 1 | d | 8/8 (100%) | -0.82 | 0 | 100 | 100 | 20, 22, 28, 29 | 0 |
| 1 | e | 8/8 (100%) | -0.70 | 0 | 100 | 100 | 24, 27, 28, 28 | 0 |
| 1 | f | 8/8 (100%) | -0.80 | 0 | 100 | 100 | 24, 26, 32, 32 | 0 |
| 1 | g | 8/8 (100%) | -0.48 | 0 | 100 | 100 | 22, 24, 30, 36 | 0 |
| 1 | h | 8/8 (100%) | -0.50 | 0 | 100 | 100 | 21, 25, 37, 40 | 0 |
| 1 | i | 8/8 (100%) | -0.58 | 0 | 100 | 100 | 22, 23, 27, 32 | 0 |
| 1 | j | 8/8 (100%) | -0.60 | 0 | 100 | 100 | 23, 28, 30, 34 | 0 |
| All | All | 288/288 (100%) | -0.61 | 0 | 100 | 100 | 13, 25, 44, 53 | 0 |

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | LLDF | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|------|----------------------------|-------|
| 2 | SO4 | d | 9 | 5/5 | 0.85 | 0.33 | - | 42,46,59,65 | 5 |

Continued on next page...

Continued from previous page...

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | LLDF | B-factors(\AA^2) | Q<0.9 |
|-----|------|-------|------|-------|------|------|------|-----------------------------|-------|
| 2 | SO4 | e | 9 | 5/5 | 0.83 | 0.26 | - | 31,46,57,62 | 5 |
| 2 | SO4 | b | 9[A] | 5/5 | 0.90 | 0.50 | - | 39,42,48,51 | 5 |
| 2 | SO4 | X | 9 | 5/5 | 0.81 | 0.27 | - | 33,38,45,47 | 5 |
| 2 | SO4 | W | 9[A] | 5/5 | 0.75 | 0.29 | - | 37,40,60,61 | 5 |
| 2 | SO4 | S | 9 | 5/5 | 0.86 | 0.31 | - | 39,41,46,48 | 5 |
| 2 | SO4 | c | 9[A] | 5/5 | 0.87 | 0.26 | - | 32,38,46,52 | 5 |

6.5 Other polymers [i](#)

There are no such residues in this entry.