



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2017 – 04:29 PM EST

PDB ID : 4R1G
Title : Structure of a putative peptidoglycan glycosyltransferase from *Atopobium parvulum* in complex with cloxacillin
Authors : Filippova, E.V.; Minasov, G.; Kiryukhina, O.; Clancy, S.; Joachimiak, A.; Anderson, W.F.; Midwest Center for Structural Genomics (MCSG)
Deposited on : unknown
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6276 (1.94-1.90)
Clashscore	112137	7025 (1.94-1.90)
Ramachandran outliers	110173	6947 (1.94-1.90)
Sidechain outliers	110143	6948 (1.94-1.90)
RSRZ outliers	101464	6332 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	<div> <div>7%</div> <div> <div></div> <div>80%</div> <div>6%</div> <div>13%</div> </div> </div>
1	B	482	<div> <div>9%</div> <div> <div></div> <div>78%</div> <div>8%</div> <div>13%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6541 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptidoglycan glycosyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	417	Total	C	N	O	S	Se	0	0	0
			3010	1874	509	612	2	13			
1	B	419	Total	C	N	O	S	Se	0	1	0
			3035	1886	515	619	2	13			

There are 64 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	473	MSE	-	EXPRESSION TAG	UNP C8W8H7
A	474	HIS	-	EXPRESSION TAG	UNP C8W8H7
A	475	HIS	-	EXPRESSION TAG	UNP C8W8H7
A	476	HIS	-	EXPRESSION TAG	UNP C8W8H7
A	477	HIS	-	EXPRESSION TAG	UNP C8W8H7
A	478	HIS	-	EXPRESSION TAG	UNP C8W8H7
A	479	HIS	-	EXPRESSION TAG	UNP C8W8H7
A	480	SER	-	EXPRESSION TAG	UNP C8W8H7
A	481	SER	-	EXPRESSION TAG	UNP C8W8H7
A	482	GLY	-	EXPRESSION TAG	UNP C8W8H7
A	483	VAL	-	EXPRESSION TAG	UNP C8W8H7
A	484	ASP	-	EXPRESSION TAG	UNP C8W8H7
A	485	LEU	-	EXPRESSION TAG	UNP C8W8H7
A	486	TRP	-	EXPRESSION TAG	UNP C8W8H7
A	487	SER	-	EXPRESSION TAG	UNP C8W8H7
A	488	HIS	-	EXPRESSION TAG	UNP C8W8H7
A	489	PRO	-	EXPRESSION TAG	UNP C8W8H7
A	490	GLN	-	EXPRESSION TAG	UNP C8W8H7
A	491	PHE	-	EXPRESSION TAG	UNP C8W8H7
A	492	GLU	-	EXPRESSION TAG	UNP C8W8H7
A	493	LYS	-	EXPRESSION TAG	UNP C8W8H7
A	494	GLY	-	EXPRESSION TAG	UNP C8W8H7
A	495	THR	-	EXPRESSION TAG	UNP C8W8H7
A	496	GLU	-	EXPRESSION TAG	UNP C8W8H7
A	497	ASN	-	EXPRESSION TAG	UNP C8W8H7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	498	LEU	-	EXPRESSION TAG	UNP C8W8H7
A	499	TYR	-	EXPRESSION TAG	UNP C8W8H7
A	500	PHE	-	EXPRESSION TAG	UNP C8W8H7
A	501	GLN	-	EXPRESSION TAG	UNP C8W8H7
A	502	SER	-	EXPRESSION TAG	UNP C8W8H7
A	503	ASN	-	EXPRESSION TAG	UNP C8W8H7
A	504	ALA	-	EXPRESSION TAG	UNP C8W8H7
B	473	MSE	-	EXPRESSION TAG	UNP C8W8H7
B	474	HIS	-	EXPRESSION TAG	UNP C8W8H7
B	475	HIS	-	EXPRESSION TAG	UNP C8W8H7
B	476	HIS	-	EXPRESSION TAG	UNP C8W8H7
B	477	HIS	-	EXPRESSION TAG	UNP C8W8H7
B	478	HIS	-	EXPRESSION TAG	UNP C8W8H7
B	479	HIS	-	EXPRESSION TAG	UNP C8W8H7
B	480	SER	-	EXPRESSION TAG	UNP C8W8H7
B	481	SER	-	EXPRESSION TAG	UNP C8W8H7
B	482	GLY	-	EXPRESSION TAG	UNP C8W8H7
B	483	VAL	-	EXPRESSION TAG	UNP C8W8H7
B	484	ASP	-	EXPRESSION TAG	UNP C8W8H7
B	485	LEU	-	EXPRESSION TAG	UNP C8W8H7
B	486	TRP	-	EXPRESSION TAG	UNP C8W8H7
B	487	SER	-	EXPRESSION TAG	UNP C8W8H7
B	488	HIS	-	EXPRESSION TAG	UNP C8W8H7
B	489	PRO	-	EXPRESSION TAG	UNP C8W8H7
B	490	GLN	-	EXPRESSION TAG	UNP C8W8H7
B	491	PHE	-	EXPRESSION TAG	UNP C8W8H7
B	492	GLU	-	EXPRESSION TAG	UNP C8W8H7
B	493	LYS	-	EXPRESSION TAG	UNP C8W8H7
B	494	GLY	-	EXPRESSION TAG	UNP C8W8H7
B	495	THR	-	EXPRESSION TAG	UNP C8W8H7
B	496	GLU	-	EXPRESSION TAG	UNP C8W8H7
B	497	ASN	-	EXPRESSION TAG	UNP C8W8H7
B	498	LEU	-	EXPRESSION TAG	UNP C8W8H7
B	499	TYR	-	EXPRESSION TAG	UNP C8W8H7
B	500	PHE	-	EXPRESSION TAG	UNP C8W8H7
B	501	GLN	-	EXPRESSION TAG	UNP C8W8H7
B	502	SER	-	EXPRESSION TAG	UNP C8W8H7
B	503	ASN	-	EXPRESSION TAG	UNP C8W8H7
B	504	ALA	-	EXPRESSION TAG	UNP C8W8H7

- Molecule 2 is CLOXACILLIN (OPEN FORM) (three-letter code: CXU) (formula: C₁₉H₂₀ClN₃O₅S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 29	C 19	Cl 1	N 3	O 5	S 1	0	0
2	B	1	Total 29	C 19	Cl 1	N 3	O 5	S 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	216	Total O 217 217	0	1
3	B	221	Total O 221 221	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Peptidoglycan glycosyltransferase



• Molecule 1: Peptidoglycan glycosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.52Å 69.89Å 114.30Å 90.00° 97.08° 90.00°	Depositor
Resolution (Å)	30.00 – 1.92 29.47 – 1.92	Depositor EDS
% Data completeness (in resolution range)	99.3 (30.00-1.92) 99.3 (29.47-1.92)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.170 , 0.206 0.178 , 0.214	Depositor DCC
R_{free} test set	4005 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	32.1	Xtriage
Anisotropy	0.168	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6541	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CXU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/3048	0.81	2/4135 (0.0%)
1	B	0.69	0/3074	0.80	4/4172 (0.1%)
All	All	0.70	0/6122	0.81	6/8307 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	759	ARG	NE-CZ-NH1	5.48	123.04	120.30
1	B	759	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	A	731	ARG	NE-CZ-NH2	5.28	122.94	120.30
1	B	559	MSE	CA-CB-CG	-5.22	104.42	113.30
1	B	924	GLY	N-CA-C	-5.19	100.12	113.10
1	B	731	ARG	NE-CZ-NH2	5.04	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3010	0	2942	23	0
1	B	3035	0	2959	29	0
2	A	29	0	19	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	29	0	19	3	0
3	A	217	0	0	0	0
3	B	221	0	0	1	0
All	All	6541	0	5939	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:939[A]:ARG:HH11	1:B:939[A]:ARG:HG3	1.44	0.82
1:B:939[A]:ARG:HH11	1:B:939[A]:ARG:CG	2.03	0.71
1:A:594:ARG:HA	1:A:600:MSE:CE	2.23	0.69
1:B:597:LEU:HB2	1:B:600:MSE:HE2	1.75	0.68
1:B:900:ASN:OD1	1:B:925:ASN:HA	1.95	0.67
1:A:720:TYR:CD1	2:A:1001:CXU:H27A	2.31	0.66
1:A:594:ARG:HA	1:A:600:MSE:HE3	1.79	0.65
1:B:660:ILE:O	1:B:661:ILE:HG23	1.97	0.64
1:A:597:LEU:HB2	1:A:600:MSE:HE2	1.80	0.63
1:A:718:HIS:HB2	2:A:1001:CXU:H31	1.80	0.63
1:B:718:HIS:HB2	2:B:1001:CXU:H31	1.81	0.62
1:B:720:TYR:CD1	2:B:1001:CXU:H27A	2.34	0.61
1:B:850:LYS:NZ	3:B:1208:HOH:O	2.33	0.61
1:B:746:GLY:HA3	1:B:793:SER:OG	2.02	0.60
1:A:594:ARG:CA	1:A:600:MSE:HE3	2.34	0.58
1:B:584:LEU:O	1:B:607:GLY:HA3	2.06	0.56
1:B:882:VAL:HG21	1:B:942:ALA:HB2	1.87	0.55
1:A:594:ARG:HD2	1:A:604:ASN:HD22	1.72	0.55
1:A:720:TYR:HB2	2:A:1001:CXU:S4	2.46	0.55
1:A:681:GLY:O	1:A:795:GLY:HA3	2.07	0.54
1:A:718:HIS:HB2	2:A:1001:CXU:C31	2.36	0.54
1:B:939[A]:ARG:CG	1:B:939[A]:ARG:NH1	2.68	0.54
1:B:544:LYS:HB2	1:B:550:TYR:CE2	2.43	0.53
1:B:594:ARG:HA	1:B:600:MSE:CE	2.40	0.52
1:B:527:GLN:HB3	1:B:606:THR:HG23	1.93	0.50
1:B:924:GLY:O	1:B:925:ASN:OD1	2.31	0.49
1:B:720:TYR:CE1	2:B:1001:CXU:H27A	2.49	0.48
1:B:597:LEU:N	1:B:597:LEU:HD12	2.29	0.48
1:B:868:MSE:HE2	1:B:889:GLY:HA2	1.96	0.48
1:B:594:ARG:HD2	1:B:604:ASN:HD22	1.80	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:895:ASP:OD1	1:B:901:PHE:CE2	2.68	0.47
1:B:594:ARG:HA	1:B:600:MSE:HE3	1.95	0.47
1:B:819:ALA:HA	1:B:917:VAL:HG21	1.96	0.46
1:B:593:TRP:CE2	1:B:594:ARG:HG3	2.51	0.45
1:A:878:MSE:SE	1:A:881:ARG:HH21	2.49	0.45
1:A:901:PHE:CE2	1:A:928:ASN:HA	2.52	0.45
1:B:681:GLY:O	1:B:795:GLY:HA3	2.17	0.45
1:A:791:TRP:HH2	2:A:1001:CXU:H30	1.82	0.44
1:B:900:ASN:CG	1:B:925:ASN:HA	2.36	0.44
2:A:1001:CXU:CL	2:A:1001:CXU:C22	3.03	0.44
1:A:584:LEU:O	1:A:607:GLY:HA3	2.18	0.43
1:A:900:ASN:N	1:A:900:ASN:OD1	2.51	0.43
1:B:946:ASN:O	1:B:950:LEU:HD23	2.19	0.43
1:A:632:SER:HB2	1:A:673:THR:HG22	1.99	0.43
1:A:597:LEU:HD12	1:A:597:LEU:N	2.33	0.43
1:A:791:TRP:CH2	2:A:1001:CXU:H30	2.53	0.43
1:A:594:ARG:CB	1:A:600:MSE:HE3	2.50	0.42
1:A:819:ALA:HA	1:A:917:VAL:HG21	2.01	0.42
1:B:913:HIS:N	1:B:914:PRO:HD3	2.35	0.42
1:B:594:ARG:CA	1:B:600:MSE:HE3	2.50	0.41
1:A:527:GLN:HB2	1:A:550:TYR:CZ	2.56	0.41
1:A:597:LEU:HD13	1:A:600:MSE:HE2	2.02	0.41
1:A:594:ARG:HH21	1:A:600:MSE:SE	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	409/482 (85%)	402 (98%)	6 (2%)	1 (0%)	51	41
1	B	414/482 (86%)	401 (97%)	12 (3%)	1 (0%)	51	41

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	823/964 (85%)	803 (98%)	18 (2%)	2 (0%)	51 41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	547	ASP
1	B	661	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	318/356 (89%)	310 (98%)	8 (2%)	53 44
1	B	321/356 (90%)	317 (99%)	4 (1%)	75 74
All	All	639/712 (90%)	627 (98%)	12 (2%)	66 56

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	545	GLN
1	A	661	ILE
1	A	711	GLU
1	A	722	ASN
1	A	737	SER
1	A	895	ASP
1	A	900	ASN
1	A	939	ARG
1	B	592	ASP
1	B	667	SER
1	B	939[A]	ARG
1	B	939[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	604	ASN
1	B	604	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	CXU	A	1001	1	21,31,31	2.79	6 (28%)	23,46,46	1.65	8 (34%)
2	CXU	B	1001	1	21,31,31	2.92	6 (28%)	23,46,46	1.70	7 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CXU	A	1001	1	-	0/6/37/37	0/2/3/3
2	CXU	B	1001	1	-	0/6/37/37	0/2/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1001	CXU	C5-S4	-4.40	1.76	1.85
2	A	1001	CXU	C5-S4	-3.67	1.77	1.85
2	B	1001	CXU	C2-C1	2.17	1.53	1.50
2	A	1001	CXU	C2-C1	2.37	1.53	1.50
2	B	1001	CXU	C27-C23	3.07	1.52	1.48
2	A	1001	CXU	C29-CL	3.12	1.81	1.73
2	A	1001	CXU	C27-C23	3.14	1.52	1.48
2	B	1001	CXU	C29-CL	3.15	1.81	1.73
2	A	1001	CXU	C28-C29	4.43	1.47	1.39
2	B	1001	CXU	C28-C29	4.92	1.48	1.39
2	A	1001	CXU	C22-C26	9.91	1.51	1.41
2	B	1001	CXU	C22-C26	10.03	1.51	1.41

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1001	CXU	C52-C5-S4	-2.89	104.36	109.20
2	A	1001	CXU	C52-C5-S4	-2.47	105.08	109.20
2	A	1001	CXU	C26-C28-C29	-2.37	117.37	121.03
2	A	1001	CXU	C28-C29-CL	-2.29	117.24	120.46
2	B	1001	CXU	O1-C1-C2	-2.24	119.24	125.22
2	B	1001	CXU	C51-C5-C52	-2.09	107.43	110.81
2	B	1001	CXU	C30-C29-CL	2.07	122.69	118.39
2	A	1001	CXU	O21-C21-C22	2.10	124.12	120.88
2	B	1001	CXU	C2-N2-C21	2.23	126.26	122.30
2	A	1001	CXU	C33-C28-C26	2.35	123.71	119.56
2	A	1001	CXU	C30-C29-CL	2.71	124.02	118.39
2	A	1001	CXU	C51-C5-S4	2.77	113.84	109.20
2	A	1001	CXU	C3-C2-N2	2.91	116.82	109.80
2	B	1001	CXU	C51-C5-S4	3.25	114.64	109.20
2	B	1001	CXU	C3-C2-N2	3.79	118.96	109.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	CXU	7	0
2	B	1001	CXU	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/482 (83%)	0.22	33 (8%) 12 14	23, 34, 72, 117	0
1	B	406/482 (84%)	0.27	43 (10%) 7 8	23, 35, 77, 111	0
All	All	810/964 (84%)	0.25	76 (9%) 9 10	23, 35, 76, 117	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	603	ILE	8.4
1	A	605	THR	6.1
1	B	603	ILE	6.1
1	B	897	GLU	5.9
1	A	896	VAL	5.7
1	B	605	THR	4.7
1	A	660	ILE	4.5
1	B	925	ASN	4.4
1	B	660	ILE	4.4
1	B	898	ASN	4.4
1	B	844	VAL	4.2
1	A	901	PHE	4.1
1	B	842	ALA	4.0
1	A	900	ASN	3.9
1	A	899	GLY	3.9
1	B	841	GLY	3.8
1	B	549	THR	3.7
1	B	932	TYR	3.6
1	B	525	TYR	3.6
1	B	840	GLU	3.5
1	B	601	ALA	3.5
1	B	526	VAL	3.4
1	A	906	ILE	3.3
1	B	661	ILE	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	606	THR	3.2
1	B	551	VAL	3.2
1	B	896	VAL	3.2
1	B	546	ASP	3.2
1	B	926	GLY	3.1
1	A	895	ASP	3.1
1	A	917	VAL	3.0
1	B	548	GLY	2.9
1	A	525	TYR	2.9
1	B	843	VAL	2.9
1	B	607	GLY	2.8
1	A	913	HIS	2.7
1	A	604	ASN	2.7
1	A	950	LEU	2.7
1	A	926	GLY	2.7
1	B	550	TYR	2.6
1	A	526	VAL	2.6
1	B	582	GLU	2.6
1	B	927	GLU	2.6
1	A	840	GLU	2.5
1	B	900	ASN	2.5
1	B	709	THR	2.5
1	B	901	PHE	2.5
1	B	722	ASN	2.5
1	A	550	TYR	2.4
1	A	918	VAL	2.4
1	A	722	ASN	2.4
1	B	839	PRO	2.3
1	A	718	HIS	2.3
1	A	949	ALA	2.3
1	A	548	GLY	2.3
1	B	950	LEU	2.3
1	B	591	SER	2.3
1	A	549	THR	2.3
1	A	686	THR	2.3
1	A	925	ASN	2.3
1	B	718	HIS	2.3
1	B	687	VAL	2.2
1	B	662	GLU	2.2
1	A	546	ASP	2.2
1	A	602	GLY	2.2
1	B	544	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	686	THR	2.2
1	A	811	VAL	2.2
1	A	724	ASP	2.2
1	A	544	LYS	2.2
1	B	949	ALA	2.1
1	B	524	ALA	2.0
1	B	724	ASP	2.0
1	B	543	VAL	2.0
1	A	842	ALA	2.0
1	B	614	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CXU	A	1001	29/29	0.85	0.23	1.29	31,50,83,98	0
2	CXU	B	1001	29/29	0.89	0.17	0.62	36,51,86,88	0

6.5 Other polymers [i](#)

There are no such residues in this entry.