



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 03:39 am GMT

PDB ID : 6R1R
Title : RIBONUCLEOTIDE REDUCTASE E441D MUTANT R1 PROTEIN FROM
ESCHERICHIA COLI
Authors : Eriksson, M.; Eklund, H.
Deposited on : 1997-09-17
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1099 (3.12-3.08)
Ramachandran outliers	110173	1057 (3.12-3.08)
Sidechain outliers	110143	1057 (3.12-3.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	761	
1	B	761	
1	C	761	
2	D	20	
2	E	20	
2	F	20	
2	P	20	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 18163 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBONUCLEOTIDE REDUCTASE R1 PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	738	Total	C	N	O	S	0	0	0
			5874	3728	1010	1111	25			
1	B	738	Total	C	N	O	S	0	0	0
			5874	3728	1010	1111	25			
1	C	738	Total	C	N	O	S	0	0	0
			5874	3728	1010	1111	25			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	441	ASP	GLU	ENGINEERED	UNP P00452
B	441	ASP	GLU	ENGINEERED	UNP P00452
C	441	ASP	GLU	ENGINEERED	UNP P00452

- Molecule 2 is a protein called RIBONUCLEOTIDE REDUCTASE R2 PROTEIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	18	Total	C	N	O	0	0	0
			140	84	21	35			
2	E	18	Total	C	N	O	0	0	0
			140	84	21	35			
2	F	18	Total	C	N	O	0	0	0
			140	84	21	35			
2	P	4	Total	C	N	O	0	0	0
			31	22	4	5			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	28	Total	O	0	0
			28	28		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	30	Total 30	O 30	0	0
3	C	32	Total 32	O 32	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

• Molecule 1: RIBONUCLEOTIDE REDUCTASE R1 PROTEIN




• Molecule 1: RIBONUCLEOTIDE REDUCTASE R1 PROTEIN






• Molecule 1: RIBONUCLEOTIDE REDUCTASE R1 PROTEIN

Chain C: 60% 31% 6%



• Molecule 2: RIBONUCLEOTIDE REDUCTASE R2 PROTEIN

Chain D: 65% 20% 5% 10%



● Molecule 2: RIBONUCLEOTIDE REDUCTASE R2 PROTEIN



● Molecule 2: RIBONUCLEOTIDE REDUCTASE R2 PROTEIN



● Molecule 2: RIBONUCLEOTIDE REDUCTASE R2 PROTEIN



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	224.49Å 224.49Å 336.25Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 3.10	Depositor
% Data completeness (in resolution range)	89.7 (20.00-3.10)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
Refinement program	REFMAC, TNT	Depositor
R, R_{free}	0.194 , 0.240	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	18163	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/6002	1.40	48/8129 (0.6%)
1	B	0.50	0/6002	1.41	53/8129 (0.7%)
1	C	0.51	0/6002	1.44	49/8129 (0.6%)
2	D	0.44	0/140	1.08	0/188
2	E	0.47	0/140	1.04	0/188
2	F	0.45	0/140	1.10	0/188
2	P	0.85	0/31	2.35	1/41 (2.4%)
All	All	0.50	0/18457	1.41	151/24992 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	2
All	All	0	6

There are no bond length outliers.

The worst 5 of 151 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	530	ARG	NE-CZ-NH1	20.18	130.39	120.30
1	C	530	ARG	NE-CZ-NH1	19.45	130.02	120.30
1	B	703	ARG	NE-CZ-NH2	-17.22	111.69	120.30
1	C	703	ARG	NE-CZ-NH1	16.34	128.47	120.30
1	A	703	ARG	NE-CZ-NH1	16.24	128.42	120.30

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	46	HIS	Mainchain
1	A	75	PRO	Mainchain
1	B	667	GLU	Mainchain
1	B	697	THR	Mainchain
1	C	46	HIS	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5874	0	5797	167	0
1	B	5874	0	5797	156	0
1	C	5874	0	5797	161	0
2	D	140	0	123	4	0
2	E	140	0	123	2	0
2	F	140	0	123	4	0
2	P	31	0	34	3	0
3	A	28	0	0	3	0
3	B	30	0	0	4	0
3	C	32	0	0	3	0
All	All	18163	0	17794	490	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 490 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:GLN:HE21	1:A:195:ARG:HH21	1.04	0.98
1:B:191:GLN:HE21	1:B:195:ARG:HH21	1.03	0.94
1:C:191:GLN:HE21	1:C:195:ARG:HH21	0.95	0.91
1:C:1:MET:H2	1:C:3:GLN:HG3	1.40	0.87
1:A:1:MET:H2	1:A:3:GLN:HG3	1.40	0.86

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	736/761 (97%)	680 (92%)	48 (6%)	8 (1%)	17	54
1	B	736/761 (97%)	681 (92%)	47 (6%)	8 (1%)	17	54
1	C	736/761 (97%)	685 (93%)	44 (6%)	7 (1%)	18	57
2	D	16/20 (80%)	14 (88%)	1 (6%)	1 (6%)	1	10
2	E	16/20 (80%)	13 (81%)	2 (12%)	1 (6%)	1	10
2	F	16/20 (80%)	14 (88%)	1 (6%)	1 (6%)	1	10
2	P	2/20 (10%)	0	2 (100%)	0	100	100
All	All	2258/2363 (96%)	2087 (92%)	145 (6%)	26 (1%)	15	51

5 of 26 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	294	GLN
1	A	735	ARG
2	D	371	SER
1	B	294	GLN
1	B	735	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	632/651 (97%)	554 (88%)	78 (12%)	5	23
1	B	632/651 (97%)	560 (89%)	72 (11%)	7	27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	632/651 (97%)	558 (88%)	74 (12%)	6	26
2	D	17/19 (90%)	16 (94%)	1 (6%)	23	58
2	E	17/19 (90%)	15 (88%)	2 (12%)	6	25
2	F	17/19 (90%)	15 (88%)	2 (12%)	6	25
2	P	3/19 (16%)	1 (33%)	2 (67%)	0	0
All	All	1950/2029 (96%)	1719 (88%)	231 (12%)	6	25

5 of 231 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	168	SER
1	B	451	ASP
1	C	542	LYS
1	B	204	LYS
1	B	298	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 62 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	415	GLN
1	B	633	ASN
1	C	654	GLN
1	B	548	GLN
1	B	686	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.