



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 15, 2017 – 06:00 PM EDT

PDB ID : 2R4J
Title : Crystal structure of Escherichia coli SeMet substituted Glycerol-3-phosphate Dehydrogenase in complex with DHAP
Authors : Yeh, J.I.; Du, S.; Chinte, U.
Deposited on : unknown
Resolution : 1.96 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	501	<div> <div>3%</div> <div>67%</div> <div>26%</div> <div>5%</div> <div>•</div> </div>
1	B	501	<div> <div>6%</div> <div>65%</div> <div>27%</div> <div>6%</div> <div>••</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BOG	A	800	-	-	-	X
2	BOG	B	800	-	-	-	X
3	PO4	B	801	-	-	-	X
5	EDO	A	1952	-	-	-	X
5	EDO	A	1955	-	-	-	X
5	EDO	A	1956	-	-	-	X
5	EDO	A	1957	-	-	X	X
5	EDO	A	1958	-	-	X	X
5	EDO	A	1959	-	-	X	-
5	EDO	A	1962	-	-	-	X
5	EDO	B	802	-	-	-	X
5	EDO	B	806	-	-	-	X
5	EDO	B	807	-	-	X	X
5	EDO	B	808	-	-	-	X
5	EDO	B	810	-	-	-	X
5	EDO	B	811	-	-	-	X
5	EDO	B	813	-	-	X	-
6	IMD	A	1960	-	-	X	X
6	IMD	A	1961	-	-	X	-
6	IMD	A	1963	-	-	X	X
6	IMD	A	1966	-	-	X	-
6	IMD	A	1967	-	-	-	X
7	13P	B	816	-	-	X	-
8	BCN	A	1969	-	-	-	X
8	BCN	B	820	-	-	-	X
8	BCN	B	821	-	-	-	X
9	TAM	B	812	-	-	X	X

2 Entry composition [i](#)


There are 10 unique types of molecules in this entry. The entry contains 8645 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aerobic glycerol-3-phosphate dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	494	Total	C	N	O	S	Se	0	0	0
			3953	2510	703	727	6	7			
1	B	497	Total	C	N	O	S	Se	0	0	0
			3981	2527	710	731	6	7			

- Molecule 2 is B-OCTYLGLUCOSIDE (three-letter code: BOG) (formula: $C_{14}H_{28}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			20	14	6		
2	A	1	Total	C	O	0	0
			20	14	6		
2	A	1	Total	C	O	0	0
			20	14	6		
2	A	1	Total	C	O	0	0
			20	14	6		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			20	14	6		
2	B	1	Total	C	O	0	0
			20	14	6		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0
			53	27	9	15	2	
4	B	1	Total	C	N	O	P	0
			53	27	9	15	2	

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total 4	C 2	O 2	0	0
5	A	1	Total 4	C 2	O 2	0	0
5	A	1	Total 4	C 2	O 2	0	0
5	A	1	Total 4	C 2	O 2	0	0
5	A	1	Total 4	C 2	O 2	0	0
5	A	1	Total 4	C 2	O 2	0	0
5	A	1	Total 4	C 2	O 2	0	0
5	A	1	Total 4	C 2	O 2	0	0
5	A	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0
5	B	1	Total 4	C 2	O 2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	N	0	0
			5	3	2		
6	A	1	Total	C	N	0	0
			5	3	2		
6	A	1	Total	C	N	0	0
			5	3	2		
6	A	1	Total	C	N	0	0
			5	3	2		
6	A	1	Total	C	N	0	0
			5	3	2		
6	B	1	Total	C	N	0	0
			5	3	2		

- Molecule 7 is 1,3-DIHYDROXYACETONEPHOSPHATE (three-letter code: 13P) (formula: $C_3H_7O_6P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	O	P	0	0
			10	3	6	1		
7	B	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 8 is BICINE (three-letter code: BCN) (formula: $C_6H_{13}NO_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			11	6	1	4		
8	B	1	Total	C	N	O	0	0
			11	6	1	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	C	N	O	0	0
			11	6	1	4		

- Molecule 9 is TRIS(HYDROXYETHYL)AMINOMETHANE (three-letter code: TAM) (formula: C₇H₁₇NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	B	1	Total	C	N	O	0	0
			11	7	1	3		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	140	Total	O	0	0
			140	140		
10	B	137	Total	O	0	0
			137	137		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aerobic glycerol-3-phosphate dehydrogenase



- Molecule 1: Aerobic glycerol-3-phosphate dehydrogenase





4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	113.92Å 114.14Å 193.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.96 40.32 – 1.96	Depositor EDS
% Data completeness (in resolution range)	100.0 (10.00-1.96) 91.0 (40.32-1.96)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	133.85 (at 1.95Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.211 , 0.263 0.218 , 0.265	Depositor DCC
R_{free} test set	4086 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	0.369	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.450 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8645	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BCN, IMD, PO4, EDO, 13P, TAM, FAD, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.10	4/4041 (0.1%)	1.03	6/5461 (0.1%)
1	B	1.12	2/4069 (0.0%)	1.05	12/5498 (0.2%)
All	All	1.11	6/8110 (0.1%)	1.04	18/10959 (0.2%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	295	VAL	C-O	5.60	1.33	1.23
1	A	447	TYR	CD2-CE2	5.58	1.47	1.39
1	A	165	GLU	CB-CG	5.50	1.62	1.52
1	B	12	GLY	N-CA	-5.27	1.38	1.46
1	A	236	VAL	CB-CG2	5.19	1.63	1.52

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	169	ARG	NE-CZ-NH2	-5.88	117.36	120.30
1	A	247	ILE	CB-CA-C	-5.74	100.11	111.60
1	A	355	LEU	CB-CG-CD1	-5.71	101.28	111.00
1	B	148	ASP	CB-CG-OD1	5.68	123.41	118.30
1	A	355	LEU	CA-CB-CG	5.60	128.17	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3953	0	3903	144	0
1	B	3981	0	3935	177	0
2	A	80	0	112	7	0
2	B	40	0	56	9	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	53	0	31	1	0
4	B	53	0	31	0	0
5	A	44	0	66	28	0
5	B	60	0	90	24	0
6	A	25	0	25	21	0
6	B	5	0	5	1	0
7	A	10	0	5	0	0
7	B	10	0	5	4	0
8	A	11	0	12	1	0
8	B	22	0	24	5	0
9	B	11	0	17	7	0
10	A	140	0	0	9	0
10	B	137	0	0	13	0
All	All	8645	0	8317	347	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 347 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:27:LEU:HD21	1:B:376:ILE:CD1	1.54	1.37
1:B:374:GLN:CB	1:B:375:GLY:HA3	1.59	1.27
1:B:467:LEU:HA	10:B:938:HOH:O	1.37	1.22
5:B:813:EDO:H22	10:B:869:HOH:O	1.37	1.20
1:B:428:LEU:HD12	1:B:428:LEU:O	1.36	1.20

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	492/501 (98%)	456 (93%)	30 (6%)	6 (1%)	15	5
1	B	495/501 (99%)	454 (92%)	26 (5%)	15 (3%)	5	1
All	All	987/1002 (98%)	910 (92%)	56 (6%)	21 (2%)	8	2

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	113	LYS
1	A	118	PRO
1	B	113	LYS
1	B	118	PRO
1	B	324	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	413/412 (100%)	377 (91%)	36 (9%)	12	3
1	B	416/412 (101%)	366 (88%)	50 (12%)	6	1
All	All	829/824 (101%)	743 (90%)	86 (10%)	8	2

5 of 86 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	4	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	116	SER
1	B	420	THR
1	B	53	LEU
1	B	103	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	424	ASN
1	A	478	ASN
1	B	494	GLN
1	A	444	HIS
1	A	473	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

48 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BOG	A	1949	-	20,20,20	0.97	1 (5%)	25,25,25	0.92	2 (8%)
2	BOG	A	1950	-	20,20,20	0.96	1 (5%)	25,25,25	1.71	5 (20%)
3	PO4	A	1951	-	4,4,4	0.64	0	6,6,6	0.61	0
5	EDO	A	1952	-	3,3,3	0.72	0	2,2,2	0.25	0
5	EDO	A	1953	-	3,3,3	0.63	0	2,2,2	0.93	0
5	EDO	A	1954	-	3,3,3	0.57	0	2,2,2	0.27	0
5	EDO	A	1955	-	3,3,3	0.69	0	2,2,2	0.38	0
5	EDO	A	1956	-	3,3,3	0.50	0	2,2,2	0.23	0
5	EDO	A	1957	-	3,3,3	1.10	0	2,2,2	0.87	0
5	EDO	A	1958	-	3,3,3	0.49	0	2,2,2	0.23	0
5	EDO	A	1959	-	3,3,3	0.68	0	2,2,2	0.50	0
6	IMD	A	1960	-	3,5,5	0.52	0	4,5,5	0.93	0
6	IMD	A	1961	-	3,5,5	0.55	0	4,5,5	0.61	0
5	EDO	A	1962	-	3,3,3	0.55	0	2,2,2	0.35	0
6	IMD	A	1963	-	3,5,5	0.73	0	4,5,5	0.50	0
5	EDO	A	1964	-	3,3,3	0.56	0	2,2,2	0.46	0
5	EDO	A	1965	-	3,3,3	0.63	0	2,2,2	0.66	0
6	IMD	A	1966	-	3,5,5	0.59	0	4,5,5	1.13	0
6	IMD	A	1967	-	3,5,5	0.73	0	4,5,5	0.50	0
7	13P	A	1968	-	9,9,9	5.41	6 (66%)	10,12,12	3.94	4 (40%)
8	BCN	A	1969	-	7,10,10	0.77	0	8,11,11	2.60	3 (37%)
4	FAD	A	600	-	51,58,58	1.77	18 (35%)	54,89,89	2.10	14 (25%)
2	BOG	A	700	-	20,20,20	0.76	1 (5%)	25,25,25	1.18	2 (8%)
2	BOG	A	800	-	20,20,20	0.63	1 (5%)	25,25,25	1.59	5 (20%)
4	FAD	B	600	-	51,58,58	2.17	11 (21%)	54,89,89	2.76	16 (29%)
2	BOG	B	700	-	20,20,20	0.80	1 (5%)	25,25,25	1.10	2 (8%)
2	BOG	B	800	-	20,20,20	0.57	0	25,25,25	1.31	3 (12%)
3	PO4	B	801	-	4,4,4	0.40	0	6,6,6	0.86	0
5	EDO	B	802	-	3,3,3	0.96	0	2,2,2	0.88	0
5	EDO	B	803	-	3,3,3	0.82	0	2,2,2	0.80	0
5	EDO	B	804	-	3,3,3	0.52	0	2,2,2	0.88	0
5	EDO	B	805	-	3,3,3	0.53	0	2,2,2	0.67	0
5	EDO	B	806	-	3,3,3	1.09	0	2,2,2	0.37	0
5	EDO	B	807	-	3,3,3	1.00	0	2,2,2	0.62	0
5	EDO	B	808	-	3,3,3	0.84	0	2,2,2	0.34	0
5	EDO	B	809	-	3,3,3	0.59	0	2,2,2	0.41	0
5	EDO	B	810	-	3,3,3	0.58	0	2,2,2	0.42	0
5	EDO	B	811	-	3,3,3	0.32	0	2,2,2	0.78	0
9	TAM	B	812	-	7,10,10	0.71	0	9,12,12	2.74	5 (55%)
5	EDO	B	813	-	3,3,3	0.26	0	2,2,2	0.85	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	B	814	-	3,3,3	0.73	0	2,2,2	0.21	0
5	EDO	B	815	-	3,3,3	0.38	0	2,2,2	0.89	0
7	13P	B	816	-	9,9,9	5.43	6 (66%)	10,12,12	3.64	4 (40%)
5	EDO	B	817	-	3,3,3	0.64	0	2,2,2	0.32	0
5	EDO	B	818	-	3,3,3	0.55	0	2,2,2	0.28	0
6	IMD	B	819	-	3,5,5	0.61	0	4,5,5	0.45	0
8	BCN	B	820	-	7,10,10	0.82	0	8,11,11	3.26	4 (50%)
8	BCN	B	821	-	7,10,10	0.64	0	8,11,11	3.85	4 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BOG	A	1949	-	-	0/11/31/31	0/1/1/1
2	BOG	A	1950	-	-	0/11/31/31	0/1/1/1
3	PO4	A	1951	-	-	0/0/0/0	0/0/0/0
5	EDO	A	1952	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1953	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1954	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1955	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1956	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1957	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1958	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1959	-	-	0/1/1/1	0/0/0/0
6	IMD	A	1960	-	-	0/0/0/0	0/1/1/1
6	IMD	A	1961	-	-	0/0/0/0	0/1/1/1
5	EDO	A	1962	-	-	0/1/1/1	0/0/0/0
6	IMD	A	1963	-	-	0/0/0/0	0/1/1/1
5	EDO	A	1964	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1965	-	-	0/1/1/1	0/0/0/0
6	IMD	A	1966	-	-	0/0/0/0	0/1/1/1
6	IMD	A	1967	-	-	0/0/0/0	0/1/1/1
7	13P	A	1968	-	-	0/7/8/8	0/0/0/0
8	BCN	A	1969	-	-	0/8/10/10	0/0/0/0
4	FAD	A	600	-	-	0/28/50/50	0/6/6/6
2	BOG	A	700	-	-	0/11/31/31	0/1/1/1
2	BOG	A	800	-	-	0/11/31/31	0/1/1/1
4	FAD	B	600	-	-	0/28/50/50	0/6/6/6
2	BOG	B	700	-	-	0/11/31/31	0/1/1/1
2	BOG	B	800	-	-	0/11/31/31	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	B	801	-	-	0/0/0/0	0/0/0/0
5	EDO	B	802	-	-	0/1/1/1	0/0/0/0
5	EDO	B	803	-	-	0/1/1/1	0/0/0/0
5	EDO	B	804	-	-	0/1/1/1	0/0/0/0
5	EDO	B	805	-	-	0/1/1/1	0/0/0/0
5	EDO	B	806	-	-	0/1/1/1	0/0/0/0
5	EDO	B	807	-	-	0/1/1/1	0/0/0/0
5	EDO	B	808	-	-	0/1/1/1	0/0/0/0
5	EDO	B	809	-	-	0/1/1/1	0/0/0/0
5	EDO	B	810	-	-	0/1/1/1	0/0/0/0
5	EDO	B	811	-	-	0/1/1/1	0/0/0/0
9	TAM	B	812	-	-	0/12/12/12	0/0/0/0
5	EDO	B	813	-	-	0/1/1/1	0/0/0/0
5	EDO	B	814	-	-	0/1/1/1	0/0/0/0
5	EDO	B	815	-	-	0/1/1/1	0/0/0/0
7	13P	B	816	-	-	0/7/8/8	0/0/0/0
5	EDO	B	817	-	-	0/1/1/1	0/0/0/0
5	EDO	B	818	-	-	0/1/1/1	0/0/0/0
6	IMD	B	819	-	-	0/0/0/0	0/1/1/1
8	BCN	B	820	-	-	0/8/10/10	0/0/0/0
8	BCN	B	821	-	-	0/8/10/10	0/0/0/0

The worst 5 of 46 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	600	FAD	C4'-C3'	-3.75	1.46	1.53
4	A	600	FAD	C6-C5X	-3.64	1.36	1.41
4	B	600	FAD	C6-C5X	-3.18	1.37	1.41
4	A	600	FAD	O4'-C4'	-2.83	1.37	1.43
4	B	600	FAD	O4B-C4B	-2.76	1.38	1.45

The worst 5 of 73 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	600	FAD	N3A-C2A-N1A	-8.74	121.25	128.86
7	A	1968	13P	O3P-P-O1	-8.74	83.48	106.73
4	A	600	FAD	N3A-C2A-N1A	-7.92	121.96	128.86
7	B	816	13P	O3P-P-O1	-7.88	85.77	106.73
4	A	600	FAD	O2A-PA-O5B	-4.47	87.04	108.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

36 monomers are involved in 97 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1950	BOG	1	0
5	A	1953	EDO	2	0
5	A	1955	EDO	2	0
5	A	1956	EDO	3	0
5	A	1957	EDO	6	0
5	A	1958	EDO	4	0
5	A	1959	EDO	7	0
6	A	1960	IMD	4	0
6	A	1961	IMD	5	0
5	A	1962	EDO	1	0
6	A	1963	IMD	8	0
5	A	1964	EDO	1	0
5	A	1965	EDO	3	0
6	A	1966	IMD	4	0
8	A	1969	BCN	1	0
4	A	600	FAD	1	0
2	A	700	BOG	1	0
2	A	800	BOG	5	0
2	B	700	BOG	2	0
2	B	800	BOG	7	0
5	B	802	EDO	1	0
5	B	803	EDO	2	0
5	B	804	EDO	1	0
5	B	806	EDO	1	0
5	B	807	EDO	7	0
5	B	808	EDO	2	0
5	B	810	EDO	2	0
5	B	811	EDO	3	0
9	B	812	TAM	7	0
5	B	813	EDO	6	0
5	B	815	EDO	1	0
7	B	816	13P	4	0
5	B	817	EDO	1	0
6	B	819	IMD	1	0
8	B	820	BCN	2	0
8	B	821	BCN	3	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	487/501 (97%)	0.45	17 (3%) 44 55	20, 41, 65, 84	0
1	B	490/501 (97%)	0.53	30 (6%) 22 31	19, 41, 66, 98	0
All	All	977/1002 (97%)	0.49	47 (4%) 31 42	19, 41, 66, 98	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	7.6
1	B	494	GLN	6.6
1	A	1	MET	6.6
1	A	325	ASP	6.6
1	B	113	LYS	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	BOG	A	800	20/20	0.34	0.94	18.95	131,136,138,138	0
8	BCN	B	820	11/11	0.74	0.44	13.55	64,69,72,74	0
8	BCN	A	1969	11/11	0.38	0.67	13.40	63,76,79,79	0
3	PO4	B	801	5/5	0.86	0.48	11.51	68,68,70,71	0
5	EDO	A	1956	4/4	0.88	0.40	11.37	41,44,50,54	0
5	EDO	B	808	4/4	0.75	0.28	9.36	52,56,57,60	0
2	BOG	B	800	20/20	0.54	0.62	9.05	133,136,139,140	0
8	BCN	B	821	11/11	0.64	0.42	8.74	58,64,70,70	0
5	EDO	B	802	4/4	0.78	0.27	8.03	38,44,48,50	0
5	EDO	A	1955	4/4	0.89	0.30	7.66	38,50,50,52	0
6	IMD	A	1960	5/5	0.85	0.18	4.44	58,58,60,60	0
6	IMD	A	1967	5/5	0.84	0.20	4.02	63,64,65,67	0
5	EDO	A	1957	4/4	0.74	0.20	3.72	43,48,48,49	0
5	EDO	A	1952	4/4	0.79	0.20	3.65	63,63,64,64	0
6	IMD	A	1963	5/5	0.93	0.17	3.26	42,47,49,50	0
9	TAM	B	812	11/11	0.81	0.20	3.17	56,63,66,70	0
5	EDO	B	807	4/4	0.72	0.21	3.07	46,47,50,52	0
5	EDO	B	810	4/4	0.92	0.18	3.05	53,53,54,58	0
5	EDO	A	1962	4/4	0.75	0.18	2.62	70,71,72,74	0
5	EDO	B	811	4/4	0.97	0.15	2.59	36,43,44,49	0
5	EDO	B	806	4/4	0.85	0.20	2.34	34,45,50,55	0
5	EDO	A	1958	4/4	0.94	0.21	2.01	32,34,40,46	0
5	EDO	B	813	4/4	0.96	0.19	1.99	36,39,41,43	0
2	BOG	B	700	20/20	0.82	0.25	1.75	63,71,77,79	0
5	EDO	A	1953	4/4	0.88	0.19	1.67	32,45,46,47	0
7	13P	B	816	10/10	0.87	0.19	1.18	28,40,45,45	0
6	IMD	A	1961	5/5	0.89	0.17	1.10	73,73,74,74	0
7	13P	A	1968	10/10	0.87	0.19	0.95	28,38,43,44	0
5	EDO	A	1959	4/4	0.92	0.14	0.75	43,48,49,52	0
2	BOG	A	700	20/20	0.78	0.22	0.71	86,91,94,94	0
5	EDO	A	1965	4/4	0.88	0.16	0.34	37,37,42,42	0
4	FAD	B	600	53/53	0.97	0.15	0.33	19,25,29,32	0
4	FAD	A	600	53/53	0.96	0.15	0.13	18,26,29,36	0
5	EDO	B	815	4/4	0.93	0.11	0.03	54,57,59,62	0
5	EDO	B	809	4/4	0.84	0.18	-	65,66,67,68	0
6	IMD	A	1966	5/5	0.83	0.37	-	70,70,72,72	0
2	BOG	A	1949	20/20	0.16	0.53	-	92,95,97,97	0
5	EDO	B	803	4/4	0.92	0.25	-	47,49,52,53	0
5	EDO	B	814	4/4	0.72	0.21	-	54,56,56,56	0
5	EDO	B	805	4/4	0.82	0.18	-	63,64,64,68	0
2	BOG	A	1950	20/20	0.44	0.52	-	90,93,94,95	0
5	EDO	A	1954	4/4	0.90	0.18	-	64,65,65,66	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	EDO	A	1964	4/4	0.84	0.19	-	64,66,66,67	0
6	IMD	B	819	5/5	0.84	0.36	-	78,78,79,81	0
3	PO4	A	1951	5/5	0.90	0.26	-	84,84,84,85	0
5	EDO	B	817	4/4	0.83	0.41	-	55,62,65,66	0
5	EDO	B	818	4/4	0.68	0.55	-	72,75,75,75	0
5	EDO	B	804	4/4	0.91	0.12	-	49,49,51,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.