



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 06:17 pm GMT

PDB ID : 4R4U
Title : Crystal structure of acyl-CoA thioesterase tesB from Yersinia pestis in complex with coenzyme A
Authors : Swarbrick, C.M.D.; Forwood, J.K.
Deposited on : 2014-08-19
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	<div> <div> <div></div> <div>84%</div> <div>11%</div> <div>• •</div> </div> </div>
1	B	289	<div> <div>2%</div> <div> <div></div> <div>80%</div> <div>11%</div> <div>• 8%</div> </div> </div>
1	C	289	<div> <div>2%</div> <div> <div></div> <div>87%</div> <div>8%</div> <div>6%</div> </div> </div>
1	D	289	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>8%</div> <div>• 9%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9126 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-CoA thioesterase II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	S	0	0	0
			2219	1414	387	411	7			
1	D	264	Total	C	N	O	S	0	0	0
			2108	1340	371	390	7			
1	C	272	Total	C	N	O	S	0	0	0
			2168	1378	380	403	7			
1	B	265	Total	C	N	O	S	0	0	0
			2112	1342	370	393	7			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP Q8D151
A	-1	ASN	-	EXPRESSION TAG	UNP Q8D151
A	0	ALA	-	EXPRESSION TAG	UNP Q8D151
D	-2	SER	-	EXPRESSION TAG	UNP Q8D151
D	-1	ASN	-	EXPRESSION TAG	UNP Q8D151
D	0	ALA	-	EXPRESSION TAG	UNP Q8D151
C	-2	SER	-	EXPRESSION TAG	UNP Q8D151
C	-1	ASN	-	EXPRESSION TAG	UNP Q8D151
C	0	ALA	-	EXPRESSION TAG	UNP Q8D151
B	-2	SER	-	EXPRESSION TAG	UNP Q8D151
B	-1	ASN	-	EXPRESSION TAG	UNP Q8D151
B	0	ALA	-	EXPRESSION TAG	UNP Q8D151

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		
2	D	1	Total	Na	0	0
			1	1		

- Molecule 3 is COENZYME A (three-letter code: COA) (formula: C₂₁H₃₆N₇O₁₆P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	D	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
3	A	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		

- Molecule 4 is MALONIC ACID (three-letter code: MLA) (formula: C₃H₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			7	3	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	117	Total 117	O 117	0	0
5	D	115	Total 115	O 115	0	0
5	C	98	Total 98	O 98	0	0
5	B	84	Total 84	O 84	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Acyl-CoA thioesterase II




• Molecule 1: Acyl-CoA thioesterase II




• Molecule 1: Acyl-CoA thioesterase II



• Molecule 1: Acyl-CoA thioesterase II





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.98Å 171.37Å 73.66Å 90.00° 109.51° 90.00°	Depositor
Resolution (Å)	85.68 – 2.20 45.91 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (85.68-2.20) 99.9 (45.91-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.34 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.201 , 0.266 0.204 , 0.267	Depositor DCC
R_{free} test set	2995 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	32.5	Xtriage
Anisotropy	0.112	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.049 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9126	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, NA, MLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/2277	0.76	1/3081 (0.0%)
1	B	0.67	0/2167	0.71	0/2935
1	C	0.69	0/2225	0.73	0/3014
1	D	0.73	0/2163	0.74	1/2929 (0.0%)
All	All	0.70	0/8832	0.73	2/11959 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	66	ARG	NE-CZ-NH1	5.68	123.14	120.30
1	A	268	ARG	NE-CZ-NH1	5.05	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2219	0	2149	27	0
1	B	2112	0	2037	23	0
1	C	2168	0	2094	14	0
1	D	2108	0	2037	16	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1	0	0	0	0
3	A	48	0	32	2	0
3	D	48	0	32	1	0
4	D	7	0	2	0	0
5	A	117	0	0	0	0
5	B	84	0	0	1	0
5	C	98	0	0	2	0
5	D	115	0	0	3	0
All	All	9126	0	8383	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:PRO:N	1:A:146:GLU:HA	2.00	0.74
1:A:191:ASP:OD2	1:A:193:ARG:NH2	2.28	0.66
1:C:164:LYS:HE2	5:C:390:HOH:O	1.97	0.64
1:C:27:GLU:HB3	1:C:29:LEU:HD13	1.81	0.63
1:A:217:ILE:HG21	1:A:224:MET:HE3	1.79	0.62
1:A:128:GLU:OE2	1:A:268:ARG:NH2	2.33	0.59
1:D:241:LEU:HD13	1:D:245:LEU:HD12	1.85	0.59
1:C:154:GLY:N	5:C:358:HOH:O	2.35	0.58
1:B:54:GLU:OE1	1:B:110:SER:HA	2.04	0.57
3:A:302:COA:O8A	3:A:302:COA:H4B	2.04	0.57
1:A:256:GLY:O	1:A:257:ALA:HB3	2.06	0.56
1:A:34:VAL:HG12	1:A:68:GLY:HA3	1.87	0.56
1:C:230:ASP:O	1:C:278:GLN:HG3	2.07	0.55
1:D:66:ARG:NH1	1:D:100:PRO:O	2.40	0.55
1:A:147:LYS:O	1:A:148:VAL:HB	2.07	0.55
1:C:60:PHE:CZ	1:C:233:MET:HE3	2.42	0.55
1:D:154:GLY:N	5:D:492:HOH:O	2.40	0.55
1:A:168:PRO:HB2	1:A:220:LEU:HD12	1.90	0.54
1:A:230:ASP:O	1:A:278:GLN:HG3	2.08	0.54
1:A:64:PHE:HB3	3:A:302:COA:H71	1.91	0.53
1:B:2:SER:HB3	1:B:185:ASN:O	2.09	0.53
1:A:112:GLU:OE2	1:B:283:ARG:NE	2.37	0.52
1:D:60:PHE:CZ	1:D:233:MET:HE3	2.45	0.52
1:A:112:GLU:CD	1:B:283:ARG:HH11	2.13	0.52
1:D:133:GLU:HA	1:D:136:ILE:HG22	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:48:LYS:HE2	1:C:193:ARG:HD2	1.91	0.51
1:D:130:LEU:O	1:D:161:ARG:HD3	2.10	0.51
1:D:115:PHE:HB3	1:C:215:HIS:CE1	2.46	0.51
1:D:119:ASN:HB2	5:D:463:HOH:O	2.10	0.51
1:A:146:GLU:CD	1:A:147:LYS:H	2.14	0.50
1:B:109:GLN:HG2	1:B:110:SER:N	2.27	0.50
1:D:4:ALA:N	5:D:452:HOH:O	2.44	0.50
3:D:301:COA:O9P	3:D:301:COA:H141	2.12	0.49
1:D:226:ILE:HA	1:D:281:VAL:O	2.13	0.48
1:A:63:TYR:CE1	1:B:61:HIS:CE1	3.02	0.47
1:B:203:SER:OG	1:B:233:MET:HE1	2.14	0.47
1:A:61:HIS:HE1	1:B:230:ASP:HB2	1.79	0.47
1:B:138:ARG:HG2	1:B:139:GLN:N	2.30	0.47
1:A:148:VAL:HG13	1:A:148:VAL:O	2.16	0.46
1:D:241:LEU:HD13	1:D:245:LEU:CD1	2.44	0.46
1:B:241:LEU:HD11	1:B:245:LEU:HD12	1.96	0.46
1:A:111:GLN:HG3	1:A:112:GLU:N	2.31	0.45
1:B:2:SER:OG	1:B:3:GLN:N	2.48	0.45
1:A:112:GLU:O	1:B:256:GLY:HA2	2.17	0.45
1:A:146:GLU:OE2	1:A:147:LYS:N	2.42	0.45
1:C:215:HIS:O	1:C:217:ILE:HG13	2.17	0.44
1:B:15:GLU:OE1	1:B:23:ARG:NE	2.47	0.44
1:C:128:GLU:OE2	1:C:268:ARG:NH2	2.50	0.44
1:D:207:PHE:CE2	1:D:278:GLN:HB3	2.53	0.44
1:B:226:ILE:HA	1:B:281:VAL:O	2.17	0.43
1:D:61:HIS:ND1	1:C:61:HIS:HB3	2.34	0.43
1:A:28:ASP:HA	1:A:70:SER:HB3	2.01	0.43
1:A:54:GLU:OE2	1:A:111:GLN:N	2.53	0.42
1:B:167:ASN:OD1	1:B:170:GLN:HB2	2.19	0.42
1:C:164:LYS:NZ	1:C:174:GLU:OE1	2.49	0.42
1:B:15:GLU:CD	1:B:23:ARG:HH21	2.23	0.42
1:B:154:GLY:N	5:B:362:HOH:O	2.52	0.42
1:B:203:SER:OG	1:B:233:MET:CE	2.67	0.42
1:A:226:ILE:HA	1:A:281:VAL:O	2.20	0.42
1:B:244:TRP:O	1:B:245:LEU:HD23	2.20	0.42
1:A:18:GLU:OE2	1:D:97:ASN:N	2.53	0.41
1:C:60:PHE:CE2	1:C:233:MET:HB3	2.56	0.41
1:A:145:PRO:N	1:A:146:GLU:CA	2.75	0.41
1:B:233:MET:HB2	1:B:233:MET:HE3	1.44	0.41
1:D:48:LYS:HD2	1:D:48:LYS:O	2.20	0.41
1:B:241:LEU:HD12	1:B:241:LEU:HA	1.71	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:247:TYR:CD1	1:B:265:ILE:HG12	2.54	0.41
1:C:233:MET:HB2	1:C:233:MET:HE3	1.68	0.41
1:C:190:ASP:HA	1:C:240:ARG:HH12	1.86	0.41
1:D:230:ASP:O	1:D:278:GLN:HG3	2.21	0.40
1:A:256:GLY:HA2	1:B:112:GLU:O	2.21	0.40
1:A:256:GLY:O	1:A:257:ALA:CB	2.70	0.40
1:A:34:VAL:HG12	1:A:68:GLY:CA	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/289 (94%)	261 (96%)	9 (3%)	1 (0%)	38	41
1	B	259/289 (90%)	247 (95%)	12 (5%)	0	100	100
1	C	268/289 (93%)	262 (98%)	6 (2%)	0	100	100
1	D	258/289 (89%)	252 (98%)	6 (2%)	0	100	100
All	All	1056/1156 (91%)	1022 (97%)	33 (3%)	1 (0%)	55	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	148	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/248 (96%)	229 (96%)	9 (4%)	38	47
1	B	226/248 (91%)	218 (96%)	8 (4%)	41	51
1	C	232/248 (94%)	228 (98%)	4 (2%)	66	79
1	D	225/248 (91%)	220 (98%)	5 (2%)	57	70
All	All	921/992 (93%)	895 (97%)	26 (3%)	49	61

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	A	85	ASN
1	A	111	GLN
1	A	112	GLU
1	A	148	VAL
1	A	149	ARG
1	A	241	LEU
1	A	247	TYR
1	A	255	SER
1	D	102	PHE
1	D	128	GLU
1	D	241	LEU
1	D	247	TYR
1	D	273	VAL
1	C	31	LEU
1	C	93	SER
1	C	141	SER
1	C	247	TYR
1	B	51	VAL
1	B	93	SER
1	B	102	PHE
1	B	138	ARG
1	B	207	PHE
1	B	247	TYR
1	B	268	ARG
1	B	273	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	COA	A	302	-	43,50,50	0.99	3 (6%)	48,75,75	1.80	8 (16%)
3	COA	D	301	-	43,50,50	1.01	3 (6%)	48,75,75	1.81	8 (16%)
4	MLA	D	303	-	0,6,6	0.00	-	0,7,7	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	COA	A	302	-	-	0/44/64/64	0/3/3/3
3	COA	D	301	-	-	0/44/64/64	0/3/3/3
4	MLA	D	303	-	-	0/0/4/4	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302	COA	C2A-N3A	2.06	1.35	1.32
3	D	301	COA	O4B-C1B	2.07	1.44	1.41
3	A	302	COA	P3B-O3B	2.80	1.64	1.59
3	D	301	COA	C5A-C4A	3.13	1.47	1.40
3	D	301	COA	P3B-O3B	3.25	1.65	1.59
3	A	302	COA	C5A-C4A	3.29	1.47	1.40

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	301	COA	N3A-C2A-N1A	-8.25	121.67	128.86
3	A	302	COA	N3A-C2A-N1A	-5.96	123.67	128.86
3	A	302	COA	CEP-CBP-CAP	-4.49	101.04	108.82
3	D	301	COA	C4A-C5A-N7A	-4.30	105.25	109.41
3	A	302	COA	C4A-C5A-N7A	-3.24	106.28	109.41
3	D	301	COA	O5P-C5P-C6P	-3.05	116.28	122.01
3	D	301	COA	C1B-N9A-C4A	-2.99	121.46	126.64
3	D	301	COA	C4B-O4B-C1B	-2.72	106.87	109.77
3	A	302	COA	C1B-N9A-C4A	-2.15	122.92	126.64
3	A	302	COA	O9P-C9P-N8P	2.03	126.97	123.07
3	A	302	COA	C3B-C2B-C1B	2.03	104.52	99.95
3	D	301	COA	C5A-C6A-N6A	2.15	124.85	120.47
3	D	301	COA	C2A-N1A-C6A	2.28	122.75	118.77
3	D	301	COA	CEP-CBP-CAP	2.44	113.05	108.82
3	A	302	COA	CDP-CBP-CAP	3.65	115.14	108.82
3	A	302	COA	CEP-CBP-CCP	4.35	114.76	108.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	COA	2	0
3	D	301	COA	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	277/289 (95%)	-0.34	3 (1%) 80 79	17, 30, 62, 89	0
1	B	265/289 (91%)	-0.04	6 (2%) 61 58	21, 38, 63, 99	0
1	C	272/289 (94%)	-0.09	6 (2%) 62 60	21, 35, 58, 91	0
1	D	264/289 (91%)	-0.13	8 (3%) 51 48	19, 32, 61, 91	0
All	All	1078/1156 (93%)	-0.15	23 (2%) 64 61	17, 34, 62, 99	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	28	ASP	3.6
1	D	138	ARG	3.1
1	D	139	GLN	3.1
1	C	155	PRO	3.1
1	B	285	HIS	3.0
1	D	134	THR	3.0
1	C	154	GLY	2.8
1	D	170	GLN	2.8
1	B	155	PRO	2.6
1	C	2	SER	2.6
1	C	140	PHE	2.4
1	A	29	LEU	2.3
1	C	28	ASP	2.3
1	B	187	LYS	2.3
1	B	169	LEU	2.2
1	B	165	PHE	2.2
1	D	154	GLY	2.2
1	B	222	PRO	2.2
1	A	140	PHE	2.2
1	D	136	ILE	2.2
1	D	187	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	169	LEU	2.1
1	C	32	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	COA	A	302	48/48	0.92	0.18	2.00	35,49,113,119	0
3	COA	D	301	48/48	0.90	0.18	1.42	40,64,130,134	0
4	MLA	D	303	7/7	0.94	0.14	-0.58	29,35,39,40	0
2	NA	D	302	1/1	0.91	0.06	-2.84	25,25,25,25	0
2	NA	A	301	1/1	0.98	0.04	-4.11	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.