



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 05:14 am GMT

PDB ID : 3R6Y  
Title : Crystal structure of chymotrypsin-treated aspartase from Bacillus sp. YM55-1  
Authors : Fibriansah, G.; Puthan Veetil, V.; Poelarends, G.J.; Thunnissen, A.-M.W.H.  
Deposited on : 2011-03-22  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1692 (3.00-3.00)
Clashscore	112137	2037 (3.00-3.00)
Ramachandran outliers	110173	1973 (3.00-3.00)
Sidechain outliers	110143	1976 (3.00-3.00)
RSRZ outliers	101464	1716 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	<div> <div>%</div> <div> <div></div> <div>73%</div> <div>24%</div> <div>• •</div> </div> </div>
1	B	401	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>22%</div> <div>• •</div> </div> </div>
1	C	401	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>20%</div> <div>•</div> </div> </div>
1	D	401	<div> <div>%</div> <div> <div></div> <div>76%</div> <div>21%</div> <div>•</div> </div> </div>
1	E	401	<div> <div>%</div> <div> <div></div> <div>74%</div> <div>23%</div> <div>•</div> </div> </div>
1	F	401	<div> <div>2%</div> <div> <div></div> <div>72%</div> <div>25%</div> <div>•</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	401	<div><div></div><div>70%25%••</div></div>
1	H	401	<div><div></div><div>72%24%••</div></div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 23984 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	390	Total	C	N	O	S	0	0	0
			3000	1892	511	578	19			
1	B	390	Total	C	N	O	S	0	0	0
			3000	1892	511	578	19			
1	C	383	Total	C	N	O	S	0	0	0
			2952	1862	503	569	18			
1	D	390	Total	C	N	O	S	0	0	0
			3000	1892	511	578	19			
1	E	391	Total	C	N	O	S	0	0	0
			3009	1897	512	581	19			
1	F	390	Total	C	N	O	S	0	0	0
			3000	1892	511	578	19			
1	G	384	Total	C	N	O	S	0	0	0
			2961	1867	504	572	18			
1	H	390	Total	C	N	O	S	0	0	0
			3000	1892	511	578	19			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	F	1	Total	Ca	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	7	Total	O	0	0
			7	7		
3	B	8	Total	O	0	0
			8	8		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	8	Total 8	O 8	0	0
3	D	12	Total 12	O 12	0	0
3	E	10	Total 10	O 10	0	0
3	F	4	Total 4	O 4	0	0
3	G	5	Total 5	O 5	0	0
3	H	6	Total 6	O 6	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Aspartase




#### • Molecule 1: Aspartase





#### • Molecule 1: Aspartase







• Molecule 1: Aspartase




• Molecule 1: Aspartase




• Molecule 1: Aspartase






• Molecule 1: Aspartase

Chain G: 70% 25%



• Molecule 1: Aspartase

Chain H: 72% 24%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.31Å 168.94Å 149.08Å 90.00° 92.23° 90.00°	Depositor
Resolution (Å)	40.00 – 3.00 38.26 – 3.00	Depositor EDS
% Data completeness (in resolution range)	98.6 (40.00-3.00) 88.0 (38.26-3.00)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.240 , 0.297 0.238 , 0.286	Depositor DCC
$R_{free}$ test set	3466 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	69.5	Xtriage
Anisotropy	0.119	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 15.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.126 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	23984	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.38	0/3049	0.50	0/4126
1	B	0.38	0/3049	0.49	0/4126
1	C	0.37	0/2999	0.50	0/4058
1	D	0.38	0/3049	0.50	0/4126
1	E	0.39	0/3058	0.51	0/4138
1	F	0.39	0/3049	0.50	0/4126
1	G	0.38	0/3008	0.50	0/4070
1	H	0.39	0/3049	0.51	0/4126
All	All	0.38	0/24310	0.50	0/32896

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3000	0	3015	88	0
1	B	3000	0	3015	95	0
1	C	2952	0	2961	68	0
1	D	3000	0	3015	74	0
1	E	3009	0	3021	98	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	3000	0	3015	86	0
1	G	2961	0	2967	100	0
1	H	3000	0	3015	105	0
2	B	1	0	0	0	0
2	F	1	0	0	0	0
3	A	7	0	0	0	0
3	B	8	0	0	1	0
3	C	8	0	0	1	0
3	D	12	0	0	0	0
3	E	10	0	0	0	0
3	F	4	0	0	0	0
3	G	5	0	0	0	0
3	H	6	0	0	0	0
All	All	23984	0	24024	582	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 582 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:ILE:HD12	1:D:379:VAL:HG12	1.27	1.12
1:E:320:ILE:HD13	1:H:236:LEU:HD23	1.28	1.12
1:F:336:VAL:HG21	1:F:379:VAL:HG11	1.29	1.12
1:E:36:ILE:HD12	1:H:379:VAL:HG12	1.31	1.10
1:D:336:VAL:HG21	1:D:379:VAL:HG11	1.23	1.10

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/401 (97%)	365 (94%)	22 (6%)	1 (0%)	44	81
1	B	388/401 (97%)	369 (95%)	19 (5%)	0	100	100
1	C	379/401 (94%)	357 (94%)	22 (6%)	0	100	100
1	D	388/401 (97%)	364 (94%)	24 (6%)	0	100	100
1	E	389/401 (97%)	360 (92%)	28 (7%)	1 (0%)	44	81
1	F	388/401 (97%)	359 (92%)	29 (8%)	0	100	100
1	G	380/401 (95%)	360 (95%)	20 (5%)	0	100	100
1	H	388/401 (97%)	359 (92%)	29 (8%)	0	100	100
All	All	3088/3208 (96%)	2893 (94%)	193 (6%)	2 (0%)	55	89

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	318	SER
1	A	354	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/339 (97%)	319 (97%)	9 (3%)	50	82
1	B	328/339 (97%)	319 (97%)	9 (3%)	50	82
1	C	322/339 (95%)	313 (97%)	9 (3%)	49	82
1	D	328/339 (97%)	319 (97%)	9 (3%)	50	82
1	E	329/339 (97%)	321 (98%)	8 (2%)	54	85
1	F	328/339 (97%)	319 (97%)	9 (3%)	50	82
1	G	323/339 (95%)	314 (97%)	9 (3%)	49	82
1	H	328/339 (97%)	314 (96%)	14 (4%)	33	72
All	All	2614/2712 (96%)	2538 (97%)	76 (3%)	48	82

5 of 76 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	352	GLU
1	E	357	GLU
1	H	190	GLN
1	D	357	GLU
1	E	170	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	249	HIS
1	H	256	HIS
1	G	190	GLN
1	C	26	GLN
1	F	226	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	390/401 (97%)	-0.05	5 (1%) 77 51	35, 60, 107, 124	0
1	B	390/401 (97%)	-0.02	8 (2%) 64 34	37, 65, 106, 120	0
1	C	383/401 (95%)	-0.04	7 (1%) 69 40	40, 63, 105, 119	0
1	D	390/401 (97%)	-0.06	6 (1%) 74 47	36, 61, 101, 116	0
1	E	391/401 (97%)	0.01	5 (1%) 77 51	36, 60, 107, 124	0
1	F	390/401 (97%)	0.01	8 (2%) 64 34	37, 64, 106, 120	0
1	G	384/401 (95%)	-0.04	2 (0%) 90 74	40, 64, 105, 119	0
1	H	390/401 (97%)	-0.06	2 (0%) 90 74	36, 61, 101, 116	0
All	All	3108/3208 (96%)	-0.03	43 (1%) 75 49	35, 62, 105, 124	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	237	ASN	9.5
1	B	237	ASN	6.0
1	E	134	HIS	5.5
1	C	237	ASN	5.4
1	C	21	ALA	4.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	CA	F	402	1/1	0.94	0.24	1.72	87,87,87,87	0
2	CA	B	402	1/1	0.93	0.23	1.60	80,80,80,80	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.