



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:24 pm GMT

PDB ID : 4R9T  
Title : L-ficolin complexed to sulphates  
Authors : Laffly, E.; Lacroix, M.; Martin, L.; Vassal-Stermann, E.; Thielens, N.; Gaboriaud, C.  
Deposited on : 2014-09-08  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1062 (2.26-2.26)
Clashscore	112137	1178 (2.26-2.26)
Ramachandran outliers	110173	1145 (2.26-2.26)
Sidechain outliers	110143	1146 (2.26-2.26)
RSRZ outliers	101464	1066 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	218	<div> <div>22%</div> <div> <div></div> <div>80%</div> <div>15%</div> <div>••</div> </div> </div>
1	B	218	<div> <div>91%</div> <div>8%</div> <div>•</div> </div>
1	C	218	<div> <div>2%</div> <div>87%</div> <div>10%</div> <div>••</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	B	303	-	-	-	X
5	SO4	B	305	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5330 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ficolin-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	218	Total	C	N	O	S	3	0	0
			1744	1098	307	330	9			
1	A	217	Total	C	N	O	S	1	1	0
			1745	1099	307	330	9			
1	C	214	Total	C	N	O	S	5	1	0
			1721	1084	302	327	8			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	71	ASN	-	EXPRESSION TAG	UNP Q15485
A	71	ASN	-	EXPRESSION TAG	UNP Q15485
C	71	ASN	-	EXPRESSION TAG	UNP Q15485

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	33	Total	O	0	0
			33	33		
6	A	7	Total	O	0	0
			7	7		
6	C	21	Total	O	0	0
			21	21		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Ficolin-2

Chain B:  91% 8%




#### • Molecule 1: Ficolin-2

Chain A:  22% 80% 15%



#### • Molecule 1: Ficolin-2

Chain C:  2% 87% 10%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.11Å 96.11Å 141.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.05 – 2.25 48.06 – 2.25	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.05-2.25) 100.0 (48.06-2.25)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.97 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.188 , 0.218 0.194 , 0.223	Depositor DCC
$R_{free}$ test set	1821 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.4	Xtriage
Anisotropy	0.301	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 42.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5330	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CA, NAG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.69	1/1793 (0.1%)	0.88	2/2425 (0.1%)
1	B	0.92	1/1792 (0.1%)	0.94	5/2425 (0.2%)
1	C	0.89	0/1771	0.96	9/2395 (0.4%)
All	All	0.84	2/5356 (0.0%)	0.93	16/7245 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	144	ARG	CD-NE	-5.49	1.37	1.46
1	A	100	ASP	CB-CG	5.10	1.62	1.51

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	144	ARG	NE-CZ-NH2	-10.46	115.07	120.30
1	C	253	ARG	NE-CZ-NH2	-9.19	115.70	120.30
1	C	253	ARG	NE-CZ-NH1	8.74	124.67	120.30
1	C	144	ARG	NE-CZ-NH2	-8.64	115.98	120.30
1	B	132	ARG	NE-CZ-NH1	7.14	123.87	120.30
1	B	132	ARG	NE-CZ-NH2	-7.12	116.74	120.30
1	A	144	ARG	NE-CZ-NH2	-6.95	116.82	120.30
1	C	78	ARG	CG-CD-NE	6.93	126.35	111.80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	144	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	C	132	ARG	NE-CZ-NH1	6.79	123.70	120.30
1	B	224	ASP	CB-CG-OD1	6.42	124.08	118.30
1	C	85	ASP	CB-CG-OD1	5.67	123.40	118.30
1	C	111	ASP	CB-CG-OD2	5.61	123.35	118.30
1	C	125	ASP	CB-CG-OD1	5.36	123.12	118.30
1	A	224	ASP	CB-CG-OD1	5.09	122.89	118.30
1	C	123	ARG	NE-CZ-NH2	-5.07	117.76	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	160	ALA	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1745	0	1620	38	0
1	B	1744	0	1615	7	0
1	C	1721	0	1595	6	0
2	B	28	0	25	0	0
3	B	4	0	3	0	0
3	C	4	0	3	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	5	0	0	0	0
5	B	10	0	0	0	0
5	C	5	0	0	0	0
6	A	7	0	0	0	0
6	B	33	0	0	0	0
6	C	21	0	0	0	0
All	All	5330	0	4861	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:247:VAL:CG2	1:A:269:LYS:HG2	1.60	1.29
1:A:161:GLN:HB3	1:A:162:GLY:HA3	1.36	1.05
1:A:247:VAL:HG23	1:A:269:LYS:HG2	1.44	0.96
1:A:247:VAL:CG2	1:A:269:LYS:CG	2.51	0.88
1:A:247:VAL:HG23	1:A:269:LYS:CG	2.06	0.85
1:A:247:VAL:HG22	1:A:269:LYS:HG2	1.64	0.79
1:A:189:ASP:OD1	1:A:191:ALA:O	2.02	0.78
1:A:160:ALA:O	1:A:161:GLN:HB2	1.84	0.77
1:A:247:VAL:HG21	1:A:269:LYS:HG2	1.65	0.77
1:A:216:GLN:HB3	1:A:243:LYS:HD3	1.67	0.77
1:A:269:LYS:HG3	1:A:270:SER:N	2.03	0.72
1:A:161:GLN:HB3	1:A:162:GLY:CA	2.18	0.72
1:A:161:GLN:CB	1:A:162:GLY:HA3	2.16	0.71
1:A:121[A]:GLN:HG2	1:A:283:MET:HG3	1.78	0.66
1:A:191:ALA:O	1:A:192:GLU:HB2	1.98	0.64
1:A:247:VAL:HG23	1:A:269:LYS:CB	2.33	0.58
1:A:163:THR:HG22	1:A:288:ALA:HB3	1.87	0.56
1:B:156:HIS:HD2	1:B:187:VAL:O	1.89	0.55
1:A:192:GLU:O	1:A:193:LYS:HG2	2.07	0.54
1:A:237:GLN:HE21	1:A:264:ASN:HD22	1.56	0.54
1:A:247:VAL:HG23	1:A:269:LYS:HB3	1.90	0.53
1:B:223:GLN:HE21	1:B:225:ASN:HD21	1.56	0.52
1:A:216:GLN:CB	1:A:243:LYS:HD3	2.39	0.52
1:A:237:GLN:NE2	1:A:264:ASN:HD22	2.08	0.52
1:B:228:ASN:ND2	1:B:244:ASN:OD1	2.42	0.52
1:C:223:GLN:HE21	1:C:225:ASN:HD21	1.59	0.49
1:C:265:GLY:H	1:C:267:ASN:HD21	1.60	0.49
1:C:156:HIS:HD2	1:C:187:VAL:O	1.95	0.49
1:A:156:HIS:HD2	1:A:187:VAL:O	1.96	0.49
1:C:93:TRP:CZ2	1:C:144:ARG:HA	2.48	0.49
1:A:121[A]:GLN:OE1	1:A:251:ASN:OD1	2.30	0.49
1:A:213:HIS:HA	1:A:216:GLN:CD	2.33	0.48
1:A:216:GLN:CD	1:A:243:LYS:HG2	2.35	0.47
1:A:193:LYS:HG3	1:A:217:SER:OG	2.14	0.47
1:A:73:CYS:SG	1:A:101:CYS:SG	3.04	0.47
1:B:265:GLY:H	1:B:267:ASN:HD21	1.62	0.47
1:A:216:GLN:NE2	1:A:243:LYS:HG2	2.31	0.45
1:A:121[A]:GLN:HG2	1:A:283:MET:CG	2.47	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:227:LEU:HG	1:A:243:LYS:HE2	2.00	0.42
1:A:265:GLY:H	1:A:267:ASN:HD21	1.67	0.42
1:C:122:ARG:HD2	1:C:282:GLU:OE2	2.19	0.42
1:B:240:TRP:CH2	1:B:250:LEU:HB2	2.55	0.42
1:A:243:LYS:HD2	1:A:243:LYS:HA	1.61	0.42
1:C:228:ASN:ND2	1:C:244:ASN:OD1	2.41	0.42
1:A:212:PHE:O	1:A:216:GLN:NE2	2.53	0.41
1:B:71:ASN:N	1:B:72:PRO:HD2	2.35	0.41
1:A:213:HIS:HA	1:A:216:GLN:NE2	2.35	0.41
1:A:246:HIS:HE1	1:A:263:ALA:O	2.04	0.41
1:A:93:TRP:CZ2	1:A:144:ARG:HA	2.55	0.40
1:B:93:TRP:CZ2	1:B:144:ARG:HA	2.56	0.40
1:A:240:TRP:CH2	1:A:250:LEU:HB2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	216/218 (99%)	198 (92%)	16 (7%)	2 (1%)	20	17
1	B	216/218 (99%)	205 (95%)	11 (5%)	0	100	100
1	C	213/218 (98%)	201 (94%)	12 (6%)	0	100	100
All	All	645/654 (99%)	604 (94%)	39 (6%)	2 (0%)	44	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	161	GLN
1	A	100	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/183 (100%)	170 (93%)	13 (7%)	17	16
1	B	183/183 (100%)	177 (97%)	6 (3%)	43	53
1	C	180/183 (98%)	172 (96%)	8 (4%)	33	37
All	All	546/549 (100%)	519 (95%)	27 (5%)	31	32

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	71	ASN
1	B	90	LEU
1	B	145	LEU
1	B	150	LEU
1	B	183	ARG
1	B	250	LEU
1	A	90	LEU
1	A	100	ASP
1	A	111	ASP
1	A	121[A]	GLN
1	A	121[B]	GLN
1	A	145	LEU
1	A	161	GLN
1	A	183	ARG
1	A	193	LYS
1	A	210	LEU
1	A	243	LYS
1	A	250	LEU
1	A	269	LYS
1	C	78	ARG
1	C	90	LEU
1	C	122	ARG
1	C	145	LEU
1	C	150	LEU
1	C	192[A]	GLU
1	C	192[B]	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	250	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	156	HIS
1	B	223	GLN
1	B	246	HIS
1	B	267	ASN
1	A	156	HIS
1	A	161	GLN
1	A	195	ASN
1	A	214	ASN
1	A	216	GLN
1	A	223	GLN
1	A	237	GLN
1	A	246	HIS
1	A	267	ASN
1	C	156	HIS
1	C	195	ASN
1	C	223	GLN
1	C	267	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	B	301	1,2	14,14,15	1.28	1 (7%)	15,19,21	1.74	5 (33%)
2	NAG	B	302	2	14,14,15	0.58	0	15,19,21	1.41	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	301	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	302	2	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	NAG	O5-C1	-3.38	1.38	1.43

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	NAG	C1-C2-N2	-2.90	105.54	110.49
2	B	301	NAG	C3-C4-C5	-2.23	106.30	110.22
2	B	301	NAG	O3-C3-C2	2.25	114.22	109.39
2	B	301	NAG	O3-C3-C4	2.43	115.65	110.36
2	B	302	NAG	O4-C4-C5	2.96	116.74	109.28
2	B	301	NAG	C6-C5-C4	2.99	119.99	113.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	SO4	A	302	-	4,4,4	0.39	0	6,6,6	0.38	0
3	ACT	B	303	-	1,3,3	0.29	0	0,3,3	0.00	-
5	SO4	B	305	-	4,4,4	0.53	0	6,6,6	1.10	1 (16%)
5	SO4	B	306	-	4,4,4	0.33	0	6,6,6	0.51	0
3	ACT	C	301	-	1,3,3	0.06	0	0,3,3	0.00	-
5	SO4	C	303	-	4,4,4	0.39	0	6,6,6	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SO4	A	302	-	-	0/0/0/0	0/0/0/0
3	ACT	B	303	-	-	0/0/0/0	0/0/0/0
5	SO4	B	305	-	-	0/0/0/0	0/0/0/0
5	SO4	B	306	-	-	0/0/0/0	0/0/0/0
3	ACT	C	301	-	-	0/0/0/0	0/0/0/0
5	SO4	C	303	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	305	SO4	O3-S-O1	2.14	121.07	109.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	217/218 (99%)	1.24	47 (21%) 1 1	33, 79, 141, 220	0
1	B	218/218 (100%)	-0.55	1 (0%) 90 91	18, 29, 51, 78	2 (0%)
1	C	214/218 (98%)	-0.33	5 (2%) 61 64	20, 32, 57, 101	1 (0%)
All	All	649/654 (99%)	0.12	53 (8%) 12 13	18, 38, 116, 220	3 (0%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	229	THR	12.4
1	A	227	LEU	11.5
1	A	216	GLN	7.3
1	A	245	CYS	6.2
1	A	230	GLY	6.0
1	A	100	ASP	6.0
1	A	212	PHE	5.9
1	C	288	ALA	5.3
1	A	193	LYS	5.0
1	A	264	ASN	4.4
1	A	262	PHE	4.4
1	A	263	ALA	4.3
1	A	202	VAL	4.3
1	A	235	MET	3.9
1	C	161	GLN	3.9
1	A	204	GLY	3.7
1	A	236	PHE	3.7
1	A	269	LYS	3.7
1	A	260	GLY	3.7
1	A	271	GLY	3.4
1	A	203	GLU	3.4
1	C	99	PRO	3.4
1	A	243	LYS	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	219	SER	3.3
1	A	234	VAL	3.3
1	A	177	TYR	3.3
1	A	102	ARG	3.2
1	A	205	SER	3.1
1	A	218	PHE	3.1
1	A	180	ALA	3.0
1	A	201	PHE	3.0
1	A	131	TYR	2.9
1	A	217	SER	2.9
1	A	191	ALA	2.8
1	A	103	PRO	2.8
1	A	244	ASN	2.7
1	A	225	ASN	2.7
1	A	223	GLN	2.7
1	A	179	PHE	2.6
1	A	163	THR	2.6
1	C	76	GLY	2.5
1	A	274	TYR	2.5
1	A	242	TYR	2.4
1	A	233	ALA	2.3
1	A	75	THR	2.3
1	A	211	THR	2.3
1	A	237	GLN	2.2
1	A	101	CYS	2.1
1	B	288	ALA	2.1
1	A	195	ASN	2.1
1	A	90	LEU	2.0
1	C	75	THR	2.0
1	A	270	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	B	301	14/15	0.94	0.13	-	35,42,47,53	0
2	NAG	B	302	14/15	0.96	0.15	-	43,52,58,59	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	SO4	B	305	5/5	0.84	0.26	13.75	27,36,44,47	5
3	ACT	B	303	4/4	0.93	0.24	5.81	33,35,38,38	0
3	ACT	C	301	4/4	0.94	0.20	1.75	26,26,27,29	4
4	CA	A	301	1/1	0.52	0.12	-1.16	71,71,71,71	1
4	CA	B	304	1/1	0.99	0.08	-1.29	26,26,26,26	0
4	CA	C	302	1/1	0.92	0.04	-1.62	42,42,42,42	0
5	SO4	A	302	5/5	0.84	0.29	-	59,60,63,64	5
5	SO4	B	306	5/5	0.92	0.27	-	42,43,47,47	5
5	SO4	C	303	5/5	0.98	0.13	-	38,39,44,45	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.