



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:37 am GMT

PDB ID : 3RCB  
Title : Crystal structure of the K102E mutant of KijD10, a 3-ketoreductase from *Actinomadura kijaniata* in complex with TDP-benzene and NADP  
Authors : Holden, H.M.; Kubiak, R.L.  
Deposited on : 2011-03-30  
Resolution : 2.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	<div> <div style="display: flex; align-items: center;"> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; left: 0; top: -5px; width: 5px; height: 10px; background-color: red;"></div> <div style="position: absolute; left: 5px; top: -5px; width: 72%; height: 10px; background-color: green;"></div> <div style="position: absolute; left: 77%; top: -5px; width: 17%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; left: 94%; top: -5px; width: 10%; height: 10px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; width: 100%; margin-top: 2px;"> <span>%</span> <span>72%</span> <span>17%</span> <span>• 10%</span> </div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	TLO	A	338	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2758 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sugar 3-ketoreductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	0	0
			2478	1571	454	447	6			

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	EXPRESSION TAG	UNP B3TMR8
A	-16	GLY	-	EXPRESSION TAG	UNP B3TMR8
A	-15	SER	-	EXPRESSION TAG	UNP B3TMR8
A	-14	SER	-	EXPRESSION TAG	UNP B3TMR8
A	-13	HIS	-	EXPRESSION TAG	UNP B3TMR8
A	-12	HIS	-	EXPRESSION TAG	UNP B3TMR8
A	-11	HIS	-	EXPRESSION TAG	UNP B3TMR8
A	-10	HIS	-	EXPRESSION TAG	UNP B3TMR8
A	-9	HIS	-	EXPRESSION TAG	UNP B3TMR8
A	-8	HIS	-	EXPRESSION TAG	UNP B3TMR8
A	-7	GLU	-	EXPRESSION TAG	UNP B3TMR8
A	-6	ASN	-	EXPRESSION TAG	UNP B3TMR8
A	-5	LEU	-	EXPRESSION TAG	UNP B3TMR8
A	-4	TYR	-	EXPRESSION TAG	UNP B3TMR8
A	-3	PHE	-	EXPRESSION TAG	UNP B3TMR8
A	-2	GLN	-	EXPRESSION TAG	UNP B3TMR8
A	-1	GLY	-	EXPRESSION TAG	UNP B3TMR8
A	0	HIS	-	EXPRESSION TAG	UNP B3TMR8
A	102	GLU	LYS	ENGINEERED MUTATION	UNP B3TMR8

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	
			48	21	7	17	3	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl		
			1	1	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na		
			1	1	0	0

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 6 is 5'-O-[(S)-HYDROXY{[(S)-HYDROXY(PHENOXY)PHOSPHORYL]OXY}P HOSPHORYL]THYMIDINE (three-letter code: TLO) (formula: C<sub>16</sub>H<sub>20</sub>N<sub>2</sub>O<sub>11</sub>P<sub>2</sub>).




- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	163	Total 163	O 163	0	0



- Molecule 1: Sugar 3-ketoreductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.37Å 103.50Å 145.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.00 – 2.49 64.77 – 2.49	Depositor EDS
% Data completeness (in resolution range)	97.0 (65.00-2.49) 96.9 (64.77-2.49)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.36 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, $R_{free}$	0.189 , 0.253 0.188 , 0.249	Depositor DCC
$R_{free}$ test set	959 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.7	Xtriage
Anisotropy	0.174	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 36.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2758	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, PO4, TLO, NAP, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.51	0/2539	1.15	11/3451 (0.3%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	50	ARG	NE-CZ-NH2	-10.83	114.89	120.30
1	A	121	ARG	NE-CZ-NH1	7.08	123.84	120.30
1	A	327	ARG	NE-CZ-NH2	-6.47	117.07	120.30
1	A	23	ARG	NE-CZ-NH1	-6.23	117.19	120.30
1	A	181	LEU	CB-CG-CD1	-6.10	100.64	111.00
1	A	323	ARG	NE-CZ-NH2	-5.71	117.44	120.30
1	A	112	GLU	OE1-CD-OE2	-5.38	116.85	123.30
1	A	323	ARG	NE-CZ-NH1	5.15	122.88	120.30
1	A	46	ASP	CB-CA-C	-5.15	100.11	110.40
1	A	54	ARG	NE-CZ-NH1	5.06	122.83	120.30
1	A	192	GLY	N-CA-C	-5.00	100.59	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2478	0	2430	42	0
2	A	48	0	25	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	5	0	0	0	0
6	A	62	0	36	6	0
7	A	163	0	0	5	0
All	All	2758	0	2491	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:VAL:HG21	1:A:26:LEU:HD21	1.61	0.82
1:A:311:ASP:OD2	6:A:338:TLO:H1'	1.91	0.71
1:A:186:TYR:HB2	1:A:187:PRO:CD	2.21	0.70
1:A:263:ALA:O	7:A:412:HOH:O	2.08	0.70
1:A:47:ARG:NH1	1:A:50:ARG:HH22	1.91	0.68
1:A:90:ARG:HD2	7:A:415:HOH:O	1.98	0.62
1:A:11:ARG:HB3	1:A:38:THR:HG21	1.82	0.62
1:A:103:PRO:HG3	1:A:181:LEU:HD22	1.83	0.58
1:A:186:TYR:HB2	1:A:187:PRO:HD3	1.88	0.56
1:A:11:ARG:HB3	1:A:38:THR:CG2	2.37	0.55
1:A:302:GLY:HA2	7:A:380:HOH:O	2.08	0.54
1:A:133:HIS:CD2	1:A:291:SER:HB2	2.43	0.53
1:A:262:PRO:HD2	1:A:265:TYR:CB	2.40	0.51
1:A:37:VAL:HG23	7:A:419:HOH:O	2.11	0.50
1:A:15:ILE:O	1:A:78:PRO:HD2	2.12	0.49
1:A:47:ARG:O	1:A:48:ALA:C	2.52	0.48
1:A:307:GLU:HB3	6:A:338:TLO:H2'A	1.96	0.48
1:A:243:ASN:OD1	1:A:243:ASN:C	2.51	0.48
1:A:159:PHE:CE1	6:A:337:TLO:HE2	2.49	0.48
1:A:138:ARG:HD3	6:A:338:TLO:O2B	2.14	0.47
1:A:262:PRO:HD2	1:A:265:TYR:HB2	1.97	0.47
1:A:186:TYR:CB	1:A:187:PRO:CD	2.87	0.47
1:A:108:ARG:NE	1:A:112:GLU:OE1	2.48	0.47
6:A:338:TLO:H6	6:A:338:TLO:O5'	2.15	0.47
1:A:181:LEU:HD23	1:A:181:LEU:HA	1.56	0.47
1:A:151:GLU:O	1:A:248:GLY:HA2	2.15	0.46
1:A:311:ASP:OD1	1:A:314:ARG:NH1	2.49	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:LEU:HD23	1:A:199:LEU:HA	1.82	0.45
1:A:305:PRO:HG2	1:A:308:TRP:CE2	2.51	0.45
1:A:26:LEU:O	1:A:27:PRO:C	2.52	0.45
1:A:331:PHE:HA	1:A:332:PRO:C	2.37	0.45
1:A:163:PRO:HB3	7:A:385:HOH:O	2.16	0.44
1:A:59:PRO:HG2	1:A:59:PRO:O	2.18	0.44
1:A:207:ARG:NH2	1:A:331:PHE:CE1	2.86	0.44
1:A:13:GLY:HA2	1:A:39:ALA:O	2.18	0.44
1:A:331:PHE:C	1:A:331:PHE:CD2	2.92	0.43
1:A:311:ASP:OD2	6:A:338:TLO:C1'	2.63	0.42
1:A:67:LEU:HD23	1:A:67:LEU:HA	1.86	0.42
1:A:26:LEU:O	1:A:30:GLU:HB2	2.21	0.41
1:A:129:PHE:O	1:A:130:MET:C	2.58	0.41
1:A:252:ARG:HD2	1:A:252:ARG:HH21	1.72	0.41
1:A:79:LEU:O	1:A:84:HIS:NE2	2.50	0.40
1:A:34:LEU:HA	1:A:34:LEU:HD23	1.79	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/350 (89%)	299 (96%)	11 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	251/281 (89%)	244 (97%)	7 (3%)	49 76

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	ILE
1	A	102	GLU
1	A	159	PHE
1	A	210	ARG
1	A	219	ASN
1	A	231	GLN
1	A	244	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	297	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAP	A	333	-	44,52,52	0.82	2 (4%)	51,80,80	2.01	13 (25%)
5	PO4	A	336	-	4,4,4	0.77	0	6,6,6	1.43	1 (16%)
6	TLO	A	337	-	26,33,33	0.79	1 (3%)	32,49,49	1.60	8 (25%)
6	TLO	A	338	-	26,33,33	0.79	0	32,49,49	2.57	6 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	333	-	-	0/27/67/67	0/5/5/5
5	PO4	A	336	-	-	0/0/0/0	0/0/0/0
6	TLO	A	337	-	-	0/17/33/33	0/3/3/3
6	TLO	A	338	-	-	0/17/33/33	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	337	TLO	O3B-CG	-2.15	1.35	1.40
2	A	333	NAP	O4D-C1D	2.16	1.44	1.41
2	A	333	NAP	C2A-N3A	2.28	1.36	1.32

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	333	NAP	N3A-C2A-N1A	-8.82	121.17	128.86
6	A	338	TLO	C5-C4-N3	-6.84	117.70	125.24
2	A	333	NAP	O7N-C7N-C3N	-4.29	114.61	119.62
2	A	333	NAP	O5B-C5B-C4B	-3.76	95.66	109.00
6	A	337	TLO	C5-C4-N3	-3.22	121.69	125.24
6	A	337	TLO	C5-C6-N1	-2.98	118.93	122.15
2	A	333	NAP	O3D-C3D-C2D	-2.89	102.58	111.83
6	A	337	TLO	CE2-CD2-CG	-2.55	114.75	118.95
6	A	337	TLO	O3'-C3'-C4'	-2.53	100.12	110.13
2	A	333	NAP	O2B-C2B-C1B	-2.45	100.92	110.06

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	333	NAP	C4A-C5A-N7A	-2.29	107.20	109.41
6	A	338	TLO	O3'-C3'-C2'	-2.26	102.60	110.83
6	A	338	TLO	C2'-C1'-N1	-2.21	109.02	114.23
2	A	333	NAP	O2A-PA-O5B	-2.14	98.05	108.14
2	A	333	NAP	O4B-C4B-C3B	2.05	109.24	105.17
6	A	337	TLO	CZ-CE2-CD2	2.10	123.09	120.21
2	A	333	NAP	O7N-C7N-N7N	2.10	125.57	122.58
5	A	336	PO4	O4-P-O2	2.23	116.09	107.90
6	A	338	TLO	C2'-C3'-C4'	2.25	107.52	102.73
6	A	337	TLO	C2'-C3'-C4'	2.37	107.78	102.73
2	A	333	NAP	C3B-C2B-C1B	2.38	107.41	102.75
2	A	333	NAP	O5D-PN-O1N	2.43	119.05	109.25
6	A	337	TLO	CD1-CG-CD2	2.47	124.08	120.19
6	A	337	TLO	C4-N3-C2	2.55	117.39	115.16
2	A	333	NAP	O3X-P2B-O1X	2.61	120.71	110.50
2	A	333	NAP	O2A-PA-O1A	4.36	134.86	112.28
6	A	338	TLO	C5M-C5-C4	4.80	125.70	120.17
6	A	338	TLO	C4-N3-C2	10.35	124.21	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	337	TLO	1	0
6	A	338	TLO	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	314/350 (89%)	-0.38	2 (0%) 89 89	4, 15, 32, 44	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	176	GLY	2.3
1	A	175	VAL	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	TLO	A	338	31/31	0.94	0.20	2.26	21,34,40,40	0
4	NA	A	335	1/1	0.99	0.15	0.35	6,6,6,6	0
6	TLO	A	337	31/31	0.98	0.14	0.17	11,18,25,26	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAP	A	333	48/48	-	-	-	15,31,54,56	48
3	CL	A	334	1/1	0.99	0.15	-	33,33,33,33	0
5	PO4	A	336	5/5	0.97	0.15	-	36,37,42,43	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.