



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 3, 2017 – 11:46 PM EDT

PDB ID : 2RDJ
Title : Snapshots of a Y-family DNA polymerase in replication: Dpo4 in apo and binary/ternary complex forms
Authors : Wong, J.H.Y.; Ling, H.
Deposited on : unknown
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	14	<div> <div>57%</div> <div>36%</div> <div>7%</div> </div>
1	E	14	<div> <div>36%</div> <div>50%</div> <div>14%</div> </div>
2	D	16	<div> <div>44%</div> <div>38%</div> <div>19%</div> </div>
3	F	15	<div> <div>60%</div> <div>33%</div> <div>7%</div> </div>
4	A	352	<div> <div>13%</div> <div>86%</div> <div>10%</div> <div>...</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	B	352	<div> <div>12%</div> <div>82%</div> <div>14%</div> <div>••</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	TMP	A	355	X	-	-	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7024 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*DGP*DGP*DGP*DAP*DCP*DCP*DCP*DTP*DTP*DCP*DGP*DAP*DAP*DT)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	14	Total	C	N	O	P	0	0	0
			284	136	53	82	13			
1	E	14	Total	C	N	O	P	0	0	0
			284	136	53	82	13			

- Molecule 2 is a DNA chain called DNA (5'-D(P*DTP*DTP*DAP*DTP*DTP*DCP*DGP*DAP*DAP*DGP*DGP*DGP*DTP*DCP*DCP*DC)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	16	Total	C	N	O	P	0	0	0
			327	156	57	98	16			

- Molecule 3 is a DNA chain called DNA (5'-D(P*DTP*DAP*DTP*DTP*DCP*DGP*DAP*DAP*DGP*DGP*DGP*DTP*DCP*DCP*DC)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	15	Total	C	N	O	P	0	0	0
			307	146	55	91	15			


- Molecule 4 is a protein called DNA polymerase IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	A	341	Total	C	N	O	S	0	0	0
			2743	1760	472	504	7			
4	B	341	Total	C	N	O	S	0	0	0
			2743	1760	472	504	7			

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	2	Total Ca 2 2	0	0
5	A	2	Total Ca 2 2	0	0

- Molecule 6 is THYMIDINE-5'-PHOSPHATE (three-letter code: TMP) (formula: $C_{10}H_{15}N_2O_8P$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N O P 21 10 2 8 1	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	C	19	Total	O	0	0
			19	19		
8	D	16	Total	O	0	0
			16	16		
8	E	20	Total	O	0	0
			20	20		
8	F	18	Total	O	0	0
			18	18		
8	A	143	Total	O	0	0
			143	143		
8	B	89	Total	O	0	0
			89	89		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(*DGP*DGP*DGP*DAP*DCP*DCP*DCP*DTP*DTP*DCP*DGP*DAP*DAP*DT)-3')

Chain C: 



- Molecule 1: DNA (5'-D(*DGP*DGP*DGP*DAP*DCP*DCP*DCP*DTP*DTP*DCP*DGP*DAP*DAP*DT)-3')

Chain E: 




- Molecule 2: DNA (5'-D(P*DTP*DTP*DAP*DTP*DTP*DCP*DGP*DAP*DAP*DGP*DGP*DGP*DTP*DCP*DCP*DC)-3')

Chain D: 





- Molecule 3: DNA (5'-D(P*DTP*DAP*DTP*DTP*DCP*DGP*DAP*DAP*DGP*DGP*DGP*DTP*DCP*DCP*DC)-3')


Chain F: 





- Molecule 4: DNA polymerase IV

Chain A: 





• Molecule 4: DNA polymerase IV



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	97.99Å 102.51Å 106.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.11 – 2.20 23.61 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (24.11-2.20) 95.3 (23.61-2.20)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.22Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.210 , 0.256 0.235 , 0.274	Depositor DCC
R_{free} test set	2657 reflections (5.12%)	DCC
Wilson B-factor (Å ²)	42.1	Xtriage
Anisotropy	0.142	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.022 for -h,l,k	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7024	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.29 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.9697e-04.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TMP, GOL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.98	0/318	1.59	4/489 (0.8%)
1	E	1.03	0/318	2.03	17/489 (3.5%)
2	D	1.02	1/365 (0.3%)	1.86	8/561 (1.4%)
3	F	1.03	0/343	1.62	4/527 (0.8%)
4	A	0.61	0/2782	0.65	0/3736
4	B	0.58	0/2782	0.66	1/3736 (0.0%)
All	All	0.70	1/6908 (0.0%)	1.01	34/9538 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	DT	C1'-N1	5.31	1.56	1.49

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	12	DG	O4'-C1'-N9	14.55	118.19	108.00
1	E	11	DG	O4'-C1'-N9	9.94	114.96	108.00
1	E	10	DC	O4'-C1'-N1	8.88	114.22	108.00
1	E	5	DC	O4'-C1'-N1	8.14	113.70	108.00
2	D	13	DG	O4'-C1'-N9	7.93	113.55	108.00
2	D	14	DG	O4'-C4'-C3'	-7.92	101.25	106.00
1	E	14	DT	O4'-C1'-N1	7.47	113.23	108.00
3	F	8	DC	O4'-C1'-N1	7.05	112.94	108.00
1	E	7	DC	C2-N3-C4	6.94	123.37	119.90
2	D	10	DA	P-O3'-C3'	6.86	127.93	119.70
1	C	4	DA	P-O3'-C3'	6.71	127.76	119.70
1	E	12	DA	O4'-C4'-C3'	-6.67	101.83	104.50
2	D	16	DC	P-O3'-C3'	6.47	127.46	119.70
1	E	3	DG	C5-C6-O6	-6.33	124.80	128.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	5	DC	N1-C2-O2	6.30	122.68	118.90
2	D	3	DT	N3-C2-O2	-6.14	118.62	122.30
3	F	6	DT	C1'-O4'-C4'	-6.06	104.04	110.10
3	F	11	DA	P-O3'-C3'	6.01	126.92	119.70
1	E	14	DT	C6-C5-C7	-5.84	119.39	122.90
1	C	8	DT	C6-C5-C7	-5.83	119.40	122.90
1	E	8	DT	O4'-C1'-N1	-5.65	104.05	108.00
1	E	12	DA	C4'-C3'-C2'	-5.63	98.03	103.10
1	E	7	DC	OP1-P-OP2	5.62	128.03	119.60
4	B	7	ASP	CB-CG-OD1	5.54	123.28	118.30
3	F	16	DC	P-O3'-C3'	5.53	126.33	119.70
2	D	3	DT	C6-N1-C2	-5.48	118.56	121.30
1	C	10	DC	N1-C1'-C2'	5.47	123.00	112.60
1	E	7	DC	C5-C6-N1	5.43	123.71	121.00
1	C	11	DG	O4'-C1'-N9	-5.41	104.22	108.00
1	E	4	DA	N1-C2-N3	-5.40	126.60	129.30
1	E	10	DC	P-O3'-C3'	5.24	125.99	119.70
1	E	7	DC	O5'-P-OP2	-5.22	101.01	105.70
2	D	14	DG	C4'-C3'-C2'	-5.17	98.45	103.10
1	E	7	DC	C4-C5-C6	-5.05	114.88	117.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	284	0	159	5	0
1	E	284	0	159	4	0
2	D	327	0	182	8	0
3	F	307	0	170	4	0
4	A	2743	0	2889	39	0
4	B	2743	0	2889	34	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	21	0	13	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	6	0	8	0	0
8	A	143	0	0	6	0
8	B	89	0	0	5	0
8	C	19	0	0	1	0
8	D	16	0	0	0	0
8	E	20	0	0	1	0
8	F	18	0	0	1	0
All	All	7024	0	6469	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:14:GLN:HE22	4:A:139:THR:H	1.03	0.94
1:C:14:DT:H3'	6:A:355:TMP:H2'2	1.49	0.92
4:B:285:HIS:HD2	8:B:355:HOH:O	1.57	0.87
4:B:14:GLN:HE22	4:B:139:THR:H	1.15	0.87
4:A:285:HIS:HD2	8:A:493:HOH:O	1.62	0.81
4:A:289:VAL:HB	4:A:332:ARG:HB2	1.66	0.78
4:B:279:ARG:HD3	8:B:430:HOH:O	1.85	0.76
6:A:355:TMP:H5'1	8:A:445:HOH:O	1.85	0.75
4:A:212:LYS:HE2	8:A:484:HOH:O	1.86	0.75
4:A:14:GLN:NE2	4:A:139:THR:H	1.84	0.70
4:B:14:GLN:NE2	4:B:139:THR:H	1.89	0.70
4:A:157:MET:CE	4:A:166:ILE:HD11	2.22	0.69
4:A:250:THR:HA	4:A:332:ARG:HG3	1.75	0.69
4:A:157:MET:HE2	4:A:166:ILE:HD11	1.76	0.67
1:E:11:DG:N7	8:E:29:HOH:O	2.28	0.65
4:B:304:HIS:HD2	4:B:305:GLY:O	1.80	0.64
4:A:248:ILE:HG12	4:A:334:GLY:HA3	1.81	0.63
4:B:248:ILE:CD1	4:B:332:ARG:HB3	2.29	0.62
4:A:248:ILE:CD1	4:A:332:ARG:HB3	2.30	0.60
1:E:14:DT:H2''	4:B:12:TYR:CD2	2.35	0.60
4:B:185:GLY:O	4:B:221:LYS:HE2	2.02	0.60
1:C:4:DA:H2''	1:C:5:DC:O5'	2.04	0.58
4:B:157:MET:CE	4:B:164:LYS:HE3	2.35	0.57
4:B:289:VAL:HB	4:B:332:ARG:HB2	1.88	0.56
4:A:242:ARG:HB2	4:A:242:ARG:HH11	1.71	0.55
4:B:285:HIS:CD2	8:B:355:HOH:O	2.44	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:12:DG:H2''	2:D:13:DG:C8	2.42	0.55
4:A:285:HIS:CD2	8:A:493:HOH:O	2.47	0.54
4:A:157:MET:CE	4:A:164:LYS:HE3	2.38	0.54
4:B:98:LYS:HB2	4:B:110:ASP:HB3	1.89	0.54
4:B:62:VAL:HG23	8:B:380:HOH:O	2.09	0.53
4:B:36:ARG:NH2	4:B:254:ASN:OD1	2.32	0.52
4:B:157:MET:HE2	4:B:164:LYS:HE3	1.91	0.52
2:D:12:DG:H2''	2:D:13:DG:H8	1.73	0.52
4:A:248:ILE:HD11	4:A:332:ARG:HB3	1.91	0.52
4:B:248:ILE:HD11	4:B:332:ARG:HB3	1.92	0.52
4:B:46:ALA:HB1	4:B:50:ALA:HB3	1.92	0.51
4:A:157:MET:HE1	4:A:166:ILE:HD11	1.93	0.51
4:B:166:ILE:HG23	4:B:170:GLU:HB3	1.91	0.51
3:F:5:DA:H5'	4:B:41:GLY:HA2	1.93	0.50
1:C:14:DT:C6	6:A:355:TMP:H2'1	2.46	0.50
4:B:186:ILE:HD11	4:B:225:LEU:HD21	1.93	0.49
4:A:14:GLN:HE22	4:A:139:THR:N	1.88	0.49
4:A:79:GLU:H	4:A:79:GLU:CD	2.15	0.49
3:F:13:DG:H5''	8:F:29:HOH:O	2.11	0.49
2:D:4:DT:H5'	4:A:41:GLY:HA2	1.95	0.49
4:A:133:LEU:O	4:A:137:LYS:HE3	2.13	0.49
4:B:159:LYS:NZ	8:B:368:HOH:O	2.46	0.49
1:C:14:DT:C5	6:A:355:TMP:H2'1	2.49	0.48
1:C:14:DT:H3'	6:A:355:TMP:C2'	2.34	0.47
8:C:31:HOH:O	4:A:285:HIS:CE1	2.67	0.47
4:A:166:ILE:HG23	4:A:170:GLU:HB3	1.98	0.46
4:A:76:MET:CE	4:A:78:LYS:HD3	2.45	0.46
2:D:4:DT:OP2	4:A:332:ARG:NH1	2.48	0.46
4:B:88:ILE:HG23	4:B:132:ILE:HG12	1.97	0.46
4:A:157:MET:HE1	4:A:164:LYS:HE3	1.98	0.46
4:A:46:ALA:HB1	4:A:50:ALA:HB3	1.97	0.46
1:E:14:DT:H2''	4:B:12:TYR:CE2	2.50	0.45
4:A:94:GLU:O	4:A:94:GLU:HG3	2.16	0.45
4:B:44:ALA:HB1	4:B:76:MET:HE2	1.99	0.45
4:A:157:MET:HE3	4:A:164:LYS:HE3	1.99	0.44
1:E:14:DT:H5''	4:B:105:ASP:CG	2.39	0.43
4:B:9:ASP:O	4:B:10:TYR:C	2.55	0.43
4:B:242:ARG:HD2	4:B:244:SER:O	2.18	0.43
4:A:304:HIS:HD2	4:A:305:GLY:O	2.02	0.43
4:A:323:LEU:HD13	4:A:330:ILE:HD11	2.01	0.43
4:B:258:LEU:O	4:B:262:LYS:HG3	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:11:DA:H2''	2:D:12:DG:H5''	2.00	0.43
2:D:7:DT:H5''	4:A:242:ARG:HH21	1.84	0.42
4:A:157:MET:HE3	4:A:164:LYS:CE	2.49	0.42
4:A:146:LYS:HB3	8:A:402:HOH:O	2.19	0.42
2:D:13:DG:H2'	2:D:14:DG:O4'	2.18	0.42
2:D:7:DT:H5''	4:A:242:ARG:HE	1.85	0.42
3:F:8:DC:H5''	4:B:242:ARG:NH1	2.35	0.42
4:A:98:LYS:HB2	4:A:110:ASP:HB3	2.01	0.41
4:A:277:ASP:HB3	8:A:361:HOH:O	2.21	0.41
4:B:106:GLU:OE2	4:B:152:LYS:NZ	2.52	0.41
4:B:32:VAL:HG22	4:B:76:MET:HE3	2.02	0.41
4:A:284:ILE:HD12	4:A:337:PHE:CE1	2.56	0.41
4:B:262:LYS:HB2	4:B:263:PRO:HD3	2.02	0.41
4:A:10:TYR:HA	6:A:355:TMP:P	2.60	0.40
4:A:100:GLU:HG3	4:A:240:ARG:HG3	2.03	0.40
4:B:62:VAL:HG12	4:B:66:LYS:NZ	2.36	0.40
3:F:8:DC:H5''	4:B:242:ARG:HH11	1.87	0.40
4:A:199:ILE:HG23	4:A:204:ASP:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	A	339/352 (96%)	329 (97%)	10 (3%)	0	100	100
4	B	339/352 (96%)	331 (98%)	8 (2%)	0	100	100
All	All	678/704 (96%)	660 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	A	300/309 (97%)	294 (98%)	6 (2%)	60	74
4	B	300/309 (97%)	286 (95%)	14 (5%)	30	37
All	All	600/618 (97%)	580 (97%)	20 (3%)	43	54

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	A	76	MET
4	A	97	GLU
4	A	137	LYS
4	A	242	ARG
4	A	248	ILE
4	A	323	LEU
4	B	38	GLU
4	B	76	MET
4	B	97	GLU
4	B	99	ILE
4	B	105	ASP
4	B	172	LYS
4	B	216	MET
4	B	221	LYS
4	B	242	ARG
4	B	248	ILE
4	B	278	LYS
4	B	294	ASP
4	B	323	LEU
4	B	336	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
4	A	14	GLN
4	A	188	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	A	285	HIS
4	A	304	HIS
4	B	14	GLN
4	B	188	ASN
4	B	285	HIS
4	B	304	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	TMP	A	355	5	17,22,22	0.73	0	24,33,33	2.11	5 (20%)
7	GOL	A	356	-	5,5,5	0.38	0	5,5,5	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	TMP	A	355	5	1/1/4/4	0/6/22/22	0/2/2/2
7	GOL	A	356	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	355	TMP	C5-C4-N3	-5.08	119.64	125.24
6	A	355	TMP	O5'-P-O1P	-3.10	97.79	106.47
6	A	355	TMP	O4'-C1'-N1	2.21	111.50	107.78
6	A	355	TMP	C3'-C2'-C1'	2.68	109.35	102.48
6	A	355	TMP	C4-N3-C2	5.92	120.34	115.16

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	A	355	TMP	C1'

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	355	TMP	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	C	14/14 (100%)	-0.09	0	100 100	36, 50, 75, 78	0
1	E	14/14 (100%)	-0.41	0	100 100	37, 43, 48, 49	0
2	D	16/16 (100%)	0.34	0	100 100	35, 53, 83, 83	0
3	F	15/15 (100%)	-0.17	0	100 100	39, 44, 55, 80	0
4	A	341/352 (96%)	0.62	44 (12%)	4 3	26, 44, 55, 61	0
4	B	341/352 (96%)	0.67	41 (12%)	5 4	28, 44, 56, 64	0
All	All	741/763 (97%)	0.59	85 (11%)	5 5	26, 44, 57, 83	0

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	B	66	LYS	4.7
4	B	4	LEU	4.4
4	B	37	PHE	4.2
4	B	3	VAL	4.0
4	A	6	VAL	4.0
4	B	48	TYR	4.0
4	A	274	TYR	3.8
4	B	6	VAL	3.8
4	B	116	ARG	3.7
4	A	252	LYS	3.6
4	A	335	VAL	3.6
4	A	116	ARG	3.5
4	B	211	ASP	3.4
4	A	333	ILE	3.3
4	B	169	GLU	3.3
4	A	286	VAL	3.3
4	B	209	GLU	3.2
4	B	210	PHE	3.2
4	A	334	GLY	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	A	169	GLU	3.2
4	B	134	GLU	3.2
4	A	304	HIS	3.1
4	B	38	GLU	3.0
4	A	5	PHE	3.0
4	A	287	VAL	3.0
4	B	63	GLU	2.9
4	B	109	LEU	2.9
4	B	335	VAL	2.8
4	A	225	LEU	2.8
4	A	15	VAL	2.8
4	A	341	ILE	2.8
4	B	248	ILE	2.8
4	A	200	ASN	2.7
4	A	4	LEU	2.7
4	B	45	THR	2.7
4	A	88	ILE	2.7
4	B	284	ILE	2.7
4	A	38	GLU	2.7
4	A	278	LYS	2.7
4	A	337	PHE	2.7
4	B	144	ILE	2.7
4	B	107	ALA	2.7
4	B	24	LYS	2.7
4	A	92	LEU	2.6
4	A	119	ARG	2.6
4	B	70	ASN	2.6
4	A	188	ASN	2.6
4	B	69	PRO	2.5
4	B	5	PHE	2.5
4	B	333	ILE	2.5
4	B	98	LYS	2.5
4	B	225	LEU	2.5
4	A	183	VAL	2.5
4	B	238	ARG	2.4
4	B	128	ILE	2.4
4	B	138	ILE	2.4
4	B	88	ILE	2.4
4	A	107	ALA	2.4
4	A	284	ILE	2.3
4	A	303	PRO	2.3
4	A	13	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	A	211	ASP	2.3
4	A	327	GLU	2.3
4	A	8	PHE	2.3
4	B	167	ASP	2.3
4	A	113	ASP	2.2
4	A	11	PHE	2.2
4	A	3	VAL	2.2
4	B	142	VAL	2.2
4	B	274	TYR	2.2
4	B	286	VAL	2.2
4	A	186	ILE	2.2
4	A	228	LEU	2.2
4	A	192	GLU	2.2
4	A	283	ALA	2.1
4	A	149	VAL	2.1
4	A	138	ILE	2.1
4	B	92	LEU	2.1
4	B	239	THR	2.1
4	B	252	LYS	2.1
4	A	336	ARG	2.1
4	A	196	LYS	2.0
4	A	29	VAL	2.0
4	B	8	PHE	2.0
4	B	337	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	TMP	A	355	21/21	0.93	0.15	-0.35	38,40,45,46	0
5	CA	A	353	1/1	0.99	0.07	-1.90	35,35,35,35	0
5	CA	A	354	1/1	0.98	0.04	-2.06	40,40,40,40	0
5	CA	B	353	1/1	0.95	0.07	-2.73	45,45,45,45	0
5	CA	B	354	1/1	0.96	0.03	-3.41	45,45,45,45	0
7	GOL	A	356	6/6	0.53	0.30	-	60,63,65,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.