



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 05:12 pm GMT

PDB ID : 3RPX
Title : Crystal structure of complement component 1, q subcomponent binding protein, C1QBP
Authors : Tempel, W.; Tong, Y.; Crombet, L.; Shen, Y.; Guan, X.; Nedyalkova, L.; Walker, J.R.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Weigelt, J.; Park, H.; Structural Genomics Consortium (SGC)
Deposited on : 2011-04-27
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1026 (2.68-2.64)
Ramachandran outliers	110173	1010 (2.68-2.64)
Sidechain outliers	110143	1010 (2.68-2.64)
RSRZ outliers	101464	3511 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	<div> <div>6%</div> <div>71%</div> <div>5%</div> <div>24%</div> </div>
1	B	194	<div> <div>5%</div> <div>66%</div> <div>7%</div> <div>27%</div> </div>
1	C	194	<div> <div>4%</div> <div>64%</div> <div>•</div> <div>31%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	UNX	A	46	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	UNX	A	55	-	-	-	X
2	UNX	A	60	-	-	-	X
2	UNX	B	45	-	-	-	X
2	UNX	B	47	-	-	-	X
2	UNX	B	50	-	-	-	X
2	UNX	B	51	-	-	-	X
2	UNX	B	52	-	-	-	X

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3257 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Complement component 1 Q subcomponent-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	148	Total	C	N	O	S	0	1	0
			1145	729	192	221	3			
1	B	141	Total	C	N	O	S	0	0	0
			1086	693	171	219	3			
1	C	133	Total	C	N	O	S	0	1	0
			1007	643	162	199	3			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	283	ALA	-	EXPRESSION TAG	UNP Q07021
A	284	HIS	-	EXPRESSION TAG	UNP Q07021
A	285	HIS	-	EXPRESSION TAG	UNP Q07021
A	286	HIS	-	EXPRESSION TAG	UNP Q07021
A	287	HIS	-	EXPRESSION TAG	UNP Q07021
A	288	HIS	-	EXPRESSION TAG	UNP Q07021
A	289	HIS	-	EXPRESSION TAG	UNP Q07021
B	283	ALA	-	EXPRESSION TAG	UNP Q07021
B	284	HIS	-	EXPRESSION TAG	UNP Q07021
B	285	HIS	-	EXPRESSION TAG	UNP Q07021
B	286	HIS	-	EXPRESSION TAG	UNP Q07021
B	287	HIS	-	EXPRESSION TAG	UNP Q07021
B	288	HIS	-	EXPRESSION TAG	UNP Q07021
B	289	HIS	-	EXPRESSION TAG	UNP Q07021
C	283	ALA	-	EXPRESSION TAG	UNP Q07021
C	284	HIS	-	EXPRESSION TAG	UNP Q07021
C	285	HIS	-	EXPRESSION TAG	UNP Q07021
C	286	HIS	-	EXPRESSION TAG	UNP Q07021
C	287	HIS	-	EXPRESSION TAG	UNP Q07021
C	288	HIS	-	EXPRESSION TAG	UNP Q07021
C	289	HIS	-	EXPRESSION TAG	UNP Q07021


- Molecule 2 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	10	Total 10	X 10	0	0
2	A	9	Total 9	X 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Complement component 1 Q subcomponent-binding protein



- Molecule 1: Complement component 1 Q subcomponent-binding protein



- Molecule 1: Complement component 1 Q subcomponent-binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	115.68Å 123.77Å 137.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.70 – 2.65 19.70 – 2.65	Depositor EDS
% Data completeness (in resolution range)	(Not available) (19.70-2.65) 99.9 (19.70-2.65)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.86 (at 2.67Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.195 , 0.224 (Not available) , (Not available)	Depositor DCC
R_{free} test set	NotAvailable	DCC
Wilson B-factor (Å ²)	59.0	Xtriage
Anisotropy	0.754	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 65.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3257	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/1172	0.66	0/1592
1	B	0.54	0/1104	0.65	0/1500
1	C	0.43	0/1026	0.62	0/1394
All	All	0.51	0/3302	0.64	0/4486

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1145	0	1039	6	0
1	B	1086	0	1002	7	0
1	C	1007	0	896	3	0
2	A	9	0	0	0	0
2	B	10	0	0	0	0
All	All	3257	0	2937	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 14 close contacts within the same asymmetric unit are listed below, sorted by their

clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:LEU:HD22	1:A:235:LEU:HD23	1.57	0.85
1:B:231:LEU:HD22	1:B:235:LEU:HD23	1.59	0.82
1:A:209:VAL:HG13	1:B:271:PHE:HZ	1.74	0.51
1:A:135:ILE:O	1:A:136:ASN:HB2	2.13	0.48
1:C:127:GLU:HG2	1:C:174:LYS:HA	1.95	0.47

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/194 (73%)	138 (98%)	3 (2%)	0	100	100
1	B	133/194 (69%)	130 (98%)	3 (2%)	0	100	100
1	C	124/194 (64%)	120 (97%)	3 (2%)	1 (1%)	22	34
All	All	398/582 (68%)	388 (98%)	9 (2%)	1 (0%)	44	62

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	124	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/175 (67%)	114 (97%)	3 (3%)	51	72
1	B	114/175 (65%)	112 (98%)	2 (2%)	64	82
1	C	99/175 (57%)	98 (99%)	1 (1%)	80	90
All	All	330/525 (63%)	324 (98%)	6 (2%)	64	82

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	287	HIS
1	C	136	ASN
1	B	213	SER
1	A	209	VAL
1	B	260	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	167	ASN
1	A	284	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 19 ligands modelled in this entry, 19 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	148/194 (76%)	0.16	12 (8%) 13 10	47, 76, 108, 127	0
1	B	141/194 (72%)	0.10	10 (7%) 17 14	42, 67, 116, 135	0
1	C	133/194 (68%)	0.40	8 (6%) 23 20	72, 101, 130, 139	0
All	All	422/582 (72%)	0.22	30 (7%) 17 14	42, 81, 124, 139	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	214	THR	4.9
1	A	201	SER	4.1
1	C	125	ALA	3.6
1	B	178	GLY	3.6
1	C	203	ILE	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	UNX	B	45	1/1	0.88	0.33	15.92	27,27,27,27	0
2	UNX	B	51	1/1	0.91	0.36	9.28	34,34,34,34	0
2	UNX	A	46	1/1	0.76	0.41	8.35	43,43,43,43	0
2	UNX	B	52	1/1	0.56	0.34	8.09	70,70,70,70	0
2	UNX	A	60	1/1	0.77	0.39	8.03	48,48,48,48	0
2	UNX	A	55	1/1	0.82	0.43	5.79	66,66,66,66	0
2	UNX	B	50	1/1	0.88	0.39	4.21	59,59,59,59	0
2	UNX	B	47	1/1	0.75	0.28	2.86	39,39,39,39	0
2	UNX	B	44	1/1	0.91	0.18	1.07	41,41,41,41	0
2	UNX	B	58	1/1	0.73	0.47	-	59,59,59,59	0
2	UNX	A	49	1/1	0.91	0.26	-	50,50,50,50	0
2	UNX	A	42	1/1	0.69	0.29	-	59,59,59,59	0
2	UNX	B	54	1/1	0.72	0.24	-	67,67,67,67	0
2	UNX	A	59	1/1	0.94	0.45	-	58,58,58,58	0
2	UNX	A	48	1/1	0.89	0.37	-	49,49,49,49	0
2	UNX	A	53	1/1	0.76	0.26	-	66,66,66,66	0
2	UNX	B	57	1/1	0.60	0.40	-	80,80,80,80	0
2	UNX	A	43	1/1	0.75	0.49	-	57,57,57,57	0
2	UNX	B	56	1/1	0.90	0.13	-	56,56,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.