



Full wwPDB NMR Structure Validation Report ⓘ

May 29, 2017 – 05:01 PM EDT

PDB ID : 2RVP
Title : Solution structure of DNA Containing Metallo-Base-Pair
Authors : Dairaku, T.; Furuita, K.; Sato, H.; Sebera, J.; Nakashima, K.; Kondo, J.;
Yamanaka, D.; Kondo, Y.; Okamoto, I.; Ono, A.; Sychrovsky, V.; Kojima, C.;
Tanaka, Y.
Deposited on : 2016-03-22

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20029077
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029077


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR



The overall completeness of chemical shifts assignment is 35%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | NMR archive (#Entries) |
|------------|-----------------------------|---------------------------|
| Clashscore | 125131 | 11601 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 1 | 1 | 15 |  27% 73% |
| 2 | 2 | 15 |  40% 60% |

2 Ensemble composition and analysis ⓘ

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 955 atoms, of which 348 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3').

| Mol | Chain | Residues | Atoms | | | | | | Trace |
|-----|-------|----------|-------|-----|-----|----|----|----|-------|
| 1 | 1 | 15 | Total | C | H | N | O | P | 0 |
| | | | 477 | 149 | 174 | 52 | 88 | 14 | |

- Molecule 2 is a DNA chain called DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3').

| Mol | Chain | Residues | Atoms | | | | | | Trace |
|-----|-------|----------|-------|-----|-----|----|----|----|-------|
| 2 | 2 | 15 | Total | C | H | N | O | P | 0 |
| | | | 477 | 149 | 174 | 52 | 88 | 14 | |

- Molecule 3 is SILVER ION (three-letter code: AG) (formula: Ag).

| Mol | Chain | Residues | Atoms | |
|-----|-------|----------|-------|----|
| 3 | 1 | 1 | Total | Ag |
| | | | 1 | 1 |


4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3')

Chain 1: 



- Molecule 2: DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3')

Chain 2: 




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3')

Chain 1: 



- Molecule 2: DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3')




4.2.2 Score per residue for model 2

- Molecule 1: DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3')



- Molecule 2: DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3')




4.2.3 Score per residue for model 3

- Molecule 1: DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3')



- Molecule 2: DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3')



4.2.4 Score per residue for model 4


- Molecule 1: DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3')

Chain 1:  27% 73%



- Molecule 2: DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3')

Chain 2:  33% 67%



4.2.5 Score per residue for model 5


- Molecule 1: DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3')

Chain 1:  27% 73%



- Molecule 2: DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3')

Chain 2:  33% 67%



4.2.6 Score per residue for model 6

- Molecule 1: DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3')

Chain 1:  47% 53%



- Molecule 2: DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3')


Chain 2:  53% 47%



4.2.7 Score per residue for model 7

- Molecule 1: DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3')

Chain 1:  33% 67%



- Molecule 2: DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3')


Chain 2:  53% 47%




4.2.8 Score per residue for model 8

- Molecule 1: DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3')

Chain 1:  33% 67%



- Molecule 2: DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3')

Chain 2:  33% 67%



4.2.9 Score per residue for model 9

- Molecule 1: DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3')

Chain 1:  40% 60%



- Molecule 2: DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3')


Chain 2:  33% 67%



4.2.10 Score per residue for model 10

- Molecule 1: DNA (5'-D(*TP*AP*AP*TP*AP*TP*AP*CP*TP*TP*AP*AP*TP*TP*A)-3')

Chain 1:  33% 67%



- Molecule 2: DNA (5'-D(*TP*AP*AP*TP*TP*AP*AP*CP*TP*AP*TP*AP*TP*TP*A)-3')

Chain 2:  33% 67%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

| Software name | Classification | Version |
|---------------|--------------------|---------|
| MARDIGRAS | structure solution | |
| CNS | structure solution | |
| CNS | refinement | |

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

| | |
|--|-------------|
| Chemical shift file(s) | 2rvp_cs.cif |
| Number of chemical shift lists | 1 |
| Total number of shifts | 244 |
| Number of shifts mapped to atoms | 244 |
| Number of unparsed shifts | 0 |
| Number of shifts with mapping errors | 0 |
| Number of shifts with mapping warnings | 0 |
| Assignment completeness (well-defined parts) | 35% |

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section:
AG

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes |
|-----|-------|-------|----------|----------|---------|
| 1 | 1 | 303 | 174 | 174 | 8±1 |
| 2 | 2 | 303 | 174 | 174 | 7±1 |
| All | All | 6070 | 3480 | 3480 | 140 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All unique clashes are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Clash(Å) | Distance(Å) | Models | |
|----------------|---------------|----------|-------------|--------|-------|
| | | | | Worst | Total |
| 1:1:8:DC:H2' | 1:1:9:DT:C6 | 0.62 | 2.30 | 3 | 10 |
| 1:1:9:DT:OP2 | 1:1:9:DT:H73 | 0.60 | 1.97 | 8 | 3 |
| 2:2:24:DT:OP2 | 2:2:24:DT:H73 | 0.58 | 1.98 | 10 | 8 |
| 1:1:9:DT:H73 | 1:1:9:DT:OP2 | 0.57 | 1.99 | 10 | 1 |
| 2:2:23:DC:H2' | 2:2:24:DT:C6 | 0.57 | 2.34 | 10 | 10 |
| 2:2:24:DT:H73 | 2:2:24:DT:OP2 | 0.55 | 2.00 | 7 | 2 |
| 2:2:20:DT:H2'' | 2:2:21:DA:OP2 | 0.53 | 2.03 | 4 | 8 |
| 1:1:9:DT:H71 | 1:1:9:DT:OP2 | 0.52 | 2.04 | 4 | 3 |
| 1:1:12:DA:H4' | 2:2:22:DA:H4' | 0.52 | 1.80 | 4 | 1 |
| 1:1:9:DT:OP2 | 1:1:9:DT:H71 | 0.52 | 2.05 | 1 | 3 |
| 1:1:2:DA:H2'' | 1:1:3:DA:OP2 | 0.49 | 2.06 | 1 | 10 |

Continued on next page...

Continued from previous page...

| Atom-1 | Atom-2 | Clash(Å) | Distance(Å) | Models | |
|----------------|---------------|----------|-------------|--------|-------|
| | | | | Worst | Total |
| 1:1:4:DT:H2'' | 1:1:5:DA:OP2 | 0.49 | 2.06 | 1 | 10 |
| 2:2:17:DA:H2'' | 2:2:18:DA:OP2 | 0.49 | 2.08 | 3 | 10 |
| 1:1:14:DT:H2'' | 1:1:15:DA:OP2 | 0.47 | 2.10 | 7 | 10 |
| 1:1:6:DT:H2'' | 1:1:7:DA:OP2 | 0.47 | 2.09 | 10 | 5 |
| 2:2:29:DT:H2'' | 2:2:30:DA:OP2 | 0.46 | 2.11 | 7 | 10 |
| 1:1:7:DA:H2'' | 1:1:8:DC:C6 | 0.46 | 2.45 | 2 | 5 |
| 2:2:27:DA:C8 | 2:2:28:DT:H72 | 0.45 | 2.47 | 10 | 4 |
| 1:1:11:DA:H2'' | 1:1:12:DA:OP2 | 0.44 | 2.13 | 3 | 2 |
| 2:2:28:DT:H2'' | 2:2:29:DT:OP2 | 0.43 | 2.14 | 7 | 8 |
| 2:2:23:DC:C6 | 2:2:24:DT:H72 | 0.43 | 2.48 | 9 | 1 |
| 1:1:8:DC:H2' | 1:1:9:DT:C5 | 0.42 | 2.49 | 7 | 2 |
| 1:1:9:DT:H6 | 1:1:9:DT:O5' | 0.42 | 1.98 | 4 | 4 |
| 1:1:10:DT:H2'' | 1:1:11:DA:C8 | 0.41 | 2.50 | 2 | 3 |
| 2:2:23:DC:H3' | 2:2:24:DT:H72 | 0.41 | 1.92 | 4 | 1 |
| 2:2:23:DC:H2' | 2:2:24:DT:C5 | 0.41 | 2.51 | 6 | 2 |
| 1:1:11:DA:C2 | 2:2:21:DA:C2 | 0.40 | 3.09 | 4 | 1 |
| 2:2:24:DT:O5' | 2:2:24:DT:H6 | 0.40 | 1.99 | 10 | 1 |
| 1:1:9:DT:O5' | 1:1:9:DT:H6 | 0.40 | 1.98 | 5 | 2 |

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 35% for the well-defined parts and 35% for the entire structure.

7.1 Chemical shift list 1

File name: 2rvp_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

| | |
|---|-----|
| Total number of shifts | 244 |
| Number of shifts mapped to atoms | 244 |
| Number of unparsed shifts | 0 |
| Number of shifts with mapping errors | 0 |
| Number of shifts with mapping warnings | 0 |
| Number of shift outliers (ShiftChecker) | 0 |

7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 35%, i.e. 202 atoms were assigned a chemical shift out of a possible 572. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | ¹ H | ¹³ C | ¹⁵ N |
|-----------|---------------|----------------|-----------------|-----------------|
| Backbone | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) |
| Sidechain | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) |
| Aromatic | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) |
| Overall | 202/572 (35%) | 202/332 (61%) | 0/210 (0%) | 0/30 (0%) |

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 35%, i.e. 202 atoms were assigned a chemical shift out of a possible 572. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

| | Total | ¹H | ¹³C | ¹⁵N |
|-----------|---------------|----------------------|-----------------------|-----------------------|
| Backbone | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) |
| Sidechain | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) |
| Aromatic | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) | 0/0 (—%) |
| Overall | 202/572 (35%) | 202/332 (61%) | 0/210 (0%) | 0/30 (0%) |

7.1.4 Statistically unusual chemical shifts ⓘ

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots ⓘ

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned_chem_shift_list_1). RCI is only applicable to proteins.