



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:03 pm GMT

PDB ID : 4RVT
Title : MAP4K4 in complex with a pyridin-2(1H)-one derivative
Authors : Richters, A.; Becker, C.; Kleine, S.; Rauh, D.
Deposited on : 2014-11-27
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	<div> <div>2%</div> <div> <div></div> <div>70%</div> <div>14%</div> <div>•</div> <div>14%</div> </div> </div>
1	B	328	<div> <div>%</div> <div> <div></div> <div>69%</div> <div>20%</div> <div>•</div> <div>9%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MES	A	402	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4714 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mitogen-activated protein kinase kinase kinase kinase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	0	0	0
			2223	1424	382	404	13			
1	B	297	Total	C	N	O	S	0	2	0
			2381	1513	417	438	13			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP O95819
A	-1	SER	-	EXPRESSION TAG	UNP O95819
A	0	HIS	-	EXPRESSION TAG	UNP O95819
B	-2	GLY	-	EXPRESSION TAG	UNP O95819
B	-1	SER	-	EXPRESSION TAG	UNP O95819
B	0	HIS	-	EXPRESSION TAG	UNP O95819

- Molecule 2 is 3-HEXANOYL-4-HYDROXY-5-(4-HYDROXYPHENYL)PYRIDIN-2(1H)-ONE (three-letter code: 3XM) (formula: C₁₇H₁₉NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			22	17	1	4		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	44	Total	O	0	0
			44	44		
4	B	32	Total	O	0	0
			32	32		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.99Å 90.25Å 90.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.49 – 2.40 45.48 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.49-2.40) 100.0 (45.48-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.209 , 0.278 0.213 , 0.281	Depositor DCC
R_{free} test set	1320 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	50.8	Xtriage
Anisotropy	0.098	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 33.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for -h,l,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4714	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 3XM, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2278	0.59	0/3090
1	B	0.38	0/2435	0.58	0/3299
All	All	0.38	0/4713	0.59	0/6389

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2223	0	2183	29	0
1	B	2381	0	2343	45	0
2	A	22	0	17	0	0
3	A	12	0	13	2	0
4	A	44	0	0	1	0
4	B	32	0	0	1	0
All	All	4714	0	4556	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:105:MET:HE2	1:B:171:ASP:H	1.28	0.94
1:B:236:MET:CE	1:B:244:LEU:HD12	1.99	0.93
1:B:265:PHE:CE1	1:B:289:ILE:HD11	2.08	0.89
1:B:105:MET:HE2	1:B:171:ASP:N	1.93	0.83
1:B:105:MET:HE1	1:B:170:VAL:HG13	1.61	0.81
1:A:66:ILE:N	1:A:67:LYS:HB3	1.95	0.80
1:B:236:MET:HE3	1:B:244:LEU:HD12	1.61	0.80
1:B:132:TYR:HA	1:B:292:GLN:HE22	1.58	0.68
1:B:79:HIS:HD2	1:B:81:ASN:H	1.39	0.67
1:B:236:MET:HE1	1:B:244:LEU:HD12	1.75	0.67
1:B:76:TYR:OH	1:B:171:ASP:OD1	2.08	0.67
1:B:279:SER:H	1:B:282:GLN:NE2	1.92	0.66
1:B:65:GLU:O	1:B:69:GLU:HG2	1.96	0.64
1:B:13:ILE:HA	1:B:19:ARG:NH2	2.14	0.62
1:A:151:HIS:CD2	1:A:172:PHE:HB3	2.34	0.62
1:B:265:PHE:CE1	1:B:289:ILE:CD1	2.80	0.62
1:A:65:GLU:HA	1:A:67:LYS:HD3	1.81	0.62
1:B:79:HIS:CD2	1:B:81:ASN:H	2.19	0.60
1:B:36:TYR:H	1:B:38:GLN:NE2	2.00	0.59
1:A:71:ASN:H	1:A:71:ASN:HD22	1.53	0.57
1:B:58:VAL:HG23	1:B:62:GLU:HB2	1.87	0.57
1:B:211:ASP:OD1	1:B:213:ARG:HD3	2.06	0.55
1:B:62:GLU:O	1:B:66:ILE:HG12	2.06	0.55
1:A:41:LYS:HE3	1:A:50:LEU:HD13	1.89	0.54
1:B:36:TYR:H	1:B:38:GLN:HE22	1.54	0.54
1:A:67:LYS:O	1:A:71:ASN:ND2	2.40	0.54
1:B:105:MET:HE1	1:B:170:VAL:CG1	2.37	0.53
1:B:113:ILE:O	1:B:117:VAL:HG23	2.08	0.52
1:A:276:GLN:HA	3:A:402:MES:H62	1.92	0.52
1:B:148:HIS:HA	1:B:212:TYR:OH	2.10	0.52
1:B:31:VAL:HG11	1:B:35:THR:HG22	1.92	0.51
1:A:70:ILE:HG12	1:A:103:LEU:HD21	1.93	0.51
1:B:151:HIS:O	1:B:152:ARG:HB2	2.11	0.50
1:A:173:GLY:O	1:A:174:VAL:HG23	2.12	0.50
1:B:73:LEU:CD2	1:B:175:SER:HB3	2.44	0.48
1:A:67:LYS:HG3	1:A:67:LYS:O	2.13	0.48
1:A:95:PRO:HB2	1:A:96:GLY:CA	2.44	0.48
1:B:80:ARG:HG2	4:B:423:HOH:O	2.13	0.47
1:A:95:PRO:HB2	1:A:96:GLY:HA3	1.96	0.47
1:B:246:PRO:O	1:B:272:LYS:NZ	2.35	0.46
1:A:292:GLN:O	1:A:295:GLU:HG3	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:73:LEU:HD23	1:B:175:SER:HB3	1.97	0.46
1:B:233:LEU:HD22	1:B:236:MET:HE1	1.97	0.46
1:A:68:LEU:HA	1:A:71:ASN:ND2	2.30	0.46
1:B:304:ASP:HB3	1:B:308:ARG:HH22	1.80	0.46
1:B:58:VAL:HG23	1:B:62:GLU:CB	2.46	0.46
1:B:162:THR:OG1	1:B:166:GLU:HB2	2.17	0.45
1:B:279:SER:H	1:B:282:GLN:HE21	1.64	0.45
1:A:79:HIS:HE1	1:A:81:ASN:HD22	1.63	0.45
1:A:189:ILE:HD13	1:B:239:MET:HE3	2.00	0.44
1:A:194:TRP:CH2	1:A:231:PRO:HG3	2.53	0.44
1:B:310:ARG:O	1:B:311:LYS:HB2	2.18	0.43
1:B:173:GLY:HA3	1:B:178:LEU:HD12	2.00	0.43
1:B:177:GLN:O	1:B:181:THR:HG23	2.19	0.43
1:A:188:PHE:CD1	1:A:188:PHE:C	2.91	0.43
1:A:227:ALA:HB1	1:A:258:TRP:CZ2	2.53	0.43
1:B:31:VAL:HG11	1:B:35:THR:CG2	2.49	0.43
1:B:135:ARG:HH22	1:B:295:GLU:CD	2.22	0.43
1:A:53:ILE:HG12	1:A:104:VAL:HG22	1.99	0.43
1:A:132:TYR:HA	1:A:292:GLN:HE22	1.84	0.42
1:A:15:LEU:O	1:A:90:ILE:HD12	2.19	0.42
1:B:105:MET:CE	1:B:171:ASP:N	2.75	0.42
1:A:136:GLU:HB3	1:A:167:VAL:HB	2.02	0.42
1:A:281:GLU:HG2	4:A:540:HOH:O	2.18	0.42
1:B:105:MET:HE2	1:B:171:ASP:C	2.40	0.42
1:A:95:PRO:CB	1:A:96:GLY:HA3	2.49	0.41
1:A:189:ILE:HG13	1:B:238:PRO:HB2	2.02	0.41
1:A:71:ASN:ND2	1:A:71:ASN:H	2.16	0.41
1:A:276:GLN:HB3	3:A:402:MES:H21	2.02	0.41
1:A:163:GLU:O	1:A:303:LYS:HE3	2.19	0.41
1:B:157:GLN:HG2	1:B:157:GLN:O	2.20	0.40
1:B:37:GLY:N	1:B:56:MET:HG3	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/328 (84%)	257 (94%)	13 (5%)	5 (2%)	10	12
1	B	295/328 (90%)	282 (96%)	12 (4%)	1 (0%)	44	60
All	All	570/656 (87%)	539 (95%)	25 (4%)	6 (1%)	17	23

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	94	PRO
1	A	208	ALA
1	A	66	ILE
1	B	94	PRO
1	A	95	PRO
1	A	308	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/287 (83%)	220 (92%)	19 (8%)	14	22
1	B	256/287 (89%)	236 (92%)	20 (8%)	15	23
All	All	495/574 (86%)	456 (92%)	39 (8%)	14	22

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ASP
1	A	15	LEU
1	A	19	ARG
1	A	33	ASN
1	A	56	MET
1	A	66	ILE
1	A	68	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	71	ASN
1	A	93	SER
1	A	121	LYS
1	A	213	ARG
1	A	240	ARG
1	A	252	ARG
1	A	254	LYS
1	A	256	LYS
1	A	276	GLN
1	A	280	THR
1	A	303	LYS
1	A	309	THR
1	B	13	ILE
1	B	19	ARG
1	B	43	ARG
1	B	61	ASP
1	B	69	GLU
1	B	90	ILE
1	B	118	LYS
1	B	149	VAL
1	B	163	GLU
1	B	178	LEU
1	B	180	ARG
1	B	184	ARG
1	B	185	ARG
1	B	210	TYR
1	B	240	ARG
1	B	276	GLN
1	B	280	THR
1	B	283	LEU
1	B	299	ARG
1	B	304	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	ASN
1	A	81	ASN
1	A	276	GLN
1	A	292	GLN
1	A	294	ASN
1	B	38	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	79	HIS
1	B	81	ASN
1	B	186	ASN
1	B	282	GLN
1	B	292	GLN
1	B	301	GLN
1	B	305	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	3XM	A	401	-	22,23,23	1.73	4 (18%)	25,31,31	1.37	6 (24%)
3	MES	A	402	-	12,12,12	2.08	1 (8%)	14,16,16	1.71	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3XM	A	401	-	-	0/13/13/13	0/2/2/2
3	MES	A	402	-	-	0/6/14/14	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	MES	C8-S	-6.82	1.67	1.77
2	A	401	3XM	CAG-CAD	-5.20	1.40	1.49
2	A	401	3XM	CAK-CAP	-2.71	1.41	1.49
2	A	401	3XM	CAQ-CAP	2.41	1.54	1.51
2	A	401	3XM	CAJ-NAI	4.31	1.40	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	3XM	CAG-CAH-NAI	-2.47	120.71	124.49
2	A	401	3XM	CAH-CAG-CAD	-2.45	116.36	119.77
2	A	401	3XM	OAM-CAL-CAK	-2.20	117.79	121.94
2	A	401	3XM	CAC-CAD-CAG	-2.07	117.59	120.91
2	A	401	3XM	CAH-NAI-CAJ	2.52	121.65	116.75
3	A	402	MES	O2S-S-C8	2.58	109.01	106.79
2	A	401	3XM	CAD-CAG-CAL	2.98	125.17	121.97
3	A	402	MES	O1S-S-C8	3.25	109.59	106.79
3	A	402	MES	O3S-S-C8	3.78	110.70	106.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	MES	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	281/328 (85%)	-0.28	8 (2%) 53 51	31, 48, 87, 135	1 (0%)
1	B	297/328 (90%)	-0.20	4 (1%) 77 75	35, 54, 81, 105	0
All	All	578/656 (88%)	-0.24	12 (2%) 64 61	31, 51, 84, 135	1 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	95	PRO	6.2
1	A	188	PHE	6.1
1	A	94	PRO	5.0
1	B	16	SER	4.4
1	A	209	THR	3.3
1	A	97	HIS	3.1
1	A	59	THR	2.4
1	A	93	SER	2.4
1	B	30	VAL	2.2
1	B	14	ASP	2.1
1	B	13	ILE	2.1
1	A	210	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MES	A	402	12/12	0.86	0.24	5.73	83,89,99,100	0
2	3XM	A	401	22/22	0.97	0.14	0.57	45,51,63,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.