



# Full wwPDB X-ray Structure Validation Report i

Feb 14, 2017 – 07:24 am GMT

PDB ID : 3S05  
Title : mEos2 Fluorescent Protein-Green Form  
Authors : Zhang, M.S.; Wu, L.J.; Xu, P.Y.; Liu, Y.F.; Xu, T.  
Deposited on : 2011-05-13  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 7674 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green to red photoconvertible GPF-like protein EosFP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	221	Total	C	N	O	S			
			1793	1148	305	329	11	7	0	0
1	B	222	Total	C	N	O	S			
			1798	1151	306	330	11	10	0	0
1	C	219	Total	C	N	O	S			
			1779	1140	303	326	10	10	0	0
1	D	219	Total	C	N	O	S			
			1779	1140	303	326	10	10	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	11	LYS	ASN	ENGINEERED MUTATION	UNP Q5S6Z9
A	64	CR8	HIS	CHROMOPHORE	UNP Q5S6Z9
A	64	CR8	TYR	CHROMOPHORE	UNP Q5S6Z9
A	64	CR8	GLY	CHROMOPHORE	UNP Q5S6Z9
A	70	LYS	GLU	ENGINEERED MUTATION	UNP Q5S6Z9
A	74	ASN	HIS	ENGINEERED MUTATION	UNP Q5S6Z9
A	121	TYR	HIS	ENGINEERED MUTATION	UNP Q5S6Z9
A	123	THR	VAL	ENGINEERED MUTATION	UNP Q5S6Z9
A	158	HIS	THR	ENGINEERED MUTATION	UNP Q5S6Z9
B	11	LYS	ASN	ENGINEERED MUTATION	UNP Q5S6Z9
B	64	CR8	HIS	CHROMOPHORE	UNP Q5S6Z9
B	64	CR8	TYR	CHROMOPHORE	UNP Q5S6Z9
B	64	CR8	GLY	CHROMOPHORE	UNP Q5S6Z9
B	70	LYS	GLU	ENGINEERED MUTATION	UNP Q5S6Z9
B	74	ASN	HIS	ENGINEERED MUTATION	UNP Q5S6Z9
B	121	TYR	HIS	ENGINEERED MUTATION	UNP Q5S6Z9
B	123	THR	VAL	ENGINEERED MUTATION	UNP Q5S6Z9
B	158	HIS	THR	ENGINEERED MUTATION	UNP Q5S6Z9
C	11	LYS	ASN	ENGINEERED MUTATION	UNP Q5S6Z9
C	64	CR8	HIS	CHROMOPHORE	UNP Q5S6Z9
C	64	CR8	TYR	CHROMOPHORE	UNP Q5S6Z9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	64	CR8	GLY	CHROMOPHORE	UNP Q5S6Z9
C	70	LYS	GLU	ENGINEERED MUTATION	UNP Q5S6Z9
C	74	ASN	HIS	ENGINEERED MUTATION	UNP Q5S6Z9
C	121	TYR	HIS	ENGINEERED MUTATION	UNP Q5S6Z9
C	123	THR	VAL	ENGINEERED MUTATION	UNP Q5S6Z9
C	158	HIS	THR	ENGINEERED MUTATION	UNP Q5S6Z9
D	11	LYS	ASN	ENGINEERED MUTATION	UNP Q5S6Z9
D	64	CR8	HIS	CHROMOPHORE	UNP Q5S6Z9
D	64	CR8	TYR	CHROMOPHORE	UNP Q5S6Z9
D	64	CR8	GLY	CHROMOPHORE	UNP Q5S6Z9
D	70	LYS	GLU	ENGINEERED MUTATION	UNP Q5S6Z9
D	74	ASN	HIS	ENGINEERED MUTATION	UNP Q5S6Z9
D	121	TYR	HIS	ENGINEERED MUTATION	UNP Q5S6Z9
D	123	THR	VAL	ENGINEERED MUTATION	UNP Q5S6Z9
D	158	HIS	THR	ENGINEERED MUTATION	UNP Q5S6Z9


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	132	Total O 132 132	0	0
2	B	135	Total O 135 135	0	0
2	C	148	Total O 148 148	0	0
2	D	110	Total O 110 110	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Green to red photoconvertible GPF-like protein EosFP




- Molecule 1: Green to red photoconvertible GPF-like protein EosFP



- Molecule 1: Green to red photoconvertible GPF-like protein EosFP



- Molecule 1: Green to red photoconvertible GPF-like protein EosFP



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.37 Å    105.80 Å    118.34 Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	45.81 – 2.20 48.29 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.5 (45.81-2.20) 99.5 (48.29-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	8.05 (at 2.20 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
$R$ , $R_{free}$	0.151 , 0.216 0.146 , 0.210	Depositor DCC
$R_{free}$ test set	2345 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.3	Xtriage
Anisotropy	0.205	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 40.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7674	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CR8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.29	4/1815 (0.2%)	0.94	3/2444 (0.1%)
1	B	1.11	3/1820 (0.2%)	0.89	0/2451
1	C	1.14	5/1801 (0.3%)	0.91	2/2426 (0.1%)
1	D	1.14	5/1801 (0.3%)	0.91	3/2426 (0.1%)
All	All	1.17	17/7237 (0.2%)	0.91	8/9747 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	15	GLU	CG-CD	7.32	1.62	1.51
1	D	209	LYS	CD-CE	-6.64	1.34	1.51
1	C	110	GLU	CG-CD	6.30	1.61	1.51
1	D	166	ASN	CB-CG	-6.28	1.36	1.51
1	D	144	GLU	CD-OE1	-5.54	1.19	1.25
1	B	140	GLU	CB-CG	5.46	1.62	1.52
1	D	85	LYS	CG-CD	-5.42	1.34	1.52
1	A	81	GLN	C-O	-5.39	1.13	1.23
1	C	121	TYR	CD2-CE2	5.36	1.47	1.39
1	C	85	LYS	CG-CD	5.33	1.70	1.52
1	A	82	SER	CB-OG	-5.31	1.35	1.42
1	A	89	TRP	CB-CG	5.31	1.59	1.50
1	B	164	GLU	CB-CG	-5.12	1.42	1.52
1	D	120	PHE	CE2-CZ	5.12	1.47	1.37

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	211	TYR	CD1-CE1	-5.09	1.31	1.39
1	B	207	LYS	CB-CG	-5.06	1.38	1.52
1	C	164	GLU	CD-OE1	-5.00	1.20	1.25

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	65	ASN	N-CA-CB	7.39	123.90	110.60
1	D	66	ARG	NE-CZ-NH1	-7.38	116.61	120.30
1	D	119	ARG	NE-CZ-NH1	-6.44	117.08	120.30
1	A	104	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	D	119	ARG	NE-CZ-NH2	5.25	122.92	120.30
1	A	66	ARG	NE-CZ-NH1	-5.14	117.73	120.30
1	C	66	ARG	NE-CZ-NH1	-5.04	117.78	120.30
1	C	66	ARG	NE-CZ-NH2	5.00	122.80	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	61	PHE	Mainchain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1793	0	1730	33	0
1	B	1798	0	1735	33	0
1	C	1779	0	1713	25	0
1	D	1779	0	1713	25	0
2	A	132	0	0	4	0
2	B	135	0	0	3	0
2	C	148	0	0	1	0
2	D	110	0	0	0	0
All	All	7674	0	6891	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:170:ARG:HH11	1:D:149:ARG:NH2	1.18	1.41
1:A:1:MET:SD	1:A:83:PHE:O	1.93	1.27
1:C:170:ARG:NH1	1:D:149:ARG:NH2	1.94	1.13
1:A:149:ARG:NH1	1:B:170:ARG:HH11	1.69	0.91
1:C:217:HIS:HD2	1:C:219:GLY:H	1.17	0.90
1:D:165:GLY:O	1:D:166:ASN:HB2	1.77	0.83
1:A:1:MET:HE2	1:A:84:PRO:HD3	1.62	0.82
1:D:217:HIS:HD2	1:D:219:GLY:H	1.28	0.81
1:A:149:ARG:HH12	1:B:170:ARG:NH1	1.78	0.81
1:C:170:ARG:NH1	1:D:149:ARG:HH22	1.78	0.79
1:A:149:ARG:HH12	1:B:170:ARG:HH11	1.26	0.78
1:C:65:ASN:OD1	1:C:67:VAL:HG12	1.84	0.77
1:C:65:ASN:CG	1:C:67:VAL:HG12	2.05	0.76
1:A:1:MET:CE	1:A:83:PHE:O	2.36	0.73
1:A:174:ARG:HH12	1:B:170:ARG:NH2	1.87	0.73
1:C:170:ARG:HH11	1:D:149:ARG:HH21	1.32	0.72
1:C:217:HIS:CD2	1:C:219:GLY:H	2.04	0.71
1:B:147:TYR:OH	1:B:174:ARG:NH2	2.23	0.71
1:B:217:HIS:HD2	1:B:219:GLY:H	1.39	0.69
1:A:217:HIS:HD2	1:A:219:GLY:H	1.41	0.68
1:A:93:LEU:HG	1:A:173:PHE:CE1	2.30	0.67
1:D:217:HIS:CD2	1:D:219:GLY:H	2.12	0.67
1:A:19:ASN:ND2	2:A:329:HOH:O	2.28	0.67
1:A:91:ARG:HG2	1:A:93:LEU:HD12	1.77	0.66
1:A:222:ASP:O	1:A:223:ASN:HB3	1.95	0.65
1:A:164:GLU:OE2	2:A:267:HOH:O	2.15	0.65
1:C:38:GLN:NE2	2:C:236:HOH:O	2.29	0.65
1:B:147:TYR:HH	1:B:174:ARG:HH22	1.46	0.63
1:B:217:HIS:CD2	1:B:219:GLY:H	2.19	0.61
1:C:170:ARG:HH11	1:D:149:ARG:CZ	2.03	0.61
1:D:193:ASP:OD1	1:D:217:HIS:HE1	1.85	0.59
1:A:222:ASP:O	1:A:223:ASN:CB	2.51	0.58
1:C:143:THR:H	1:D:145:LYS:NZ	2.00	0.58
1:B:67:VAL:HG21	1:B:83:PHE:CE2	2.39	0.58
1:B:7:ASP:OD1	1:B:32:LYS:HE3	2.04	0.57
1:A:1:MET:HE2	1:A:84:PRO:CD	2.33	0.57
1:D:67:VAL:HG21	1:D:83:PHE:CE2	2.39	0.57
1:A:1:MET:HE2	1:A:3:ALA:HB2	1.87	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MET:CE	1:A:84:PRO:HD3	2.35	0.56
1:A:166:ASN:O	1:A:166:ASN:ND2	2.40	0.55
1:B:94:THR:HA	1:B:100:ILE:HD12	1.89	0.55
1:B:145:LYS:HD2	1:B:191:PHE:CE1	2.42	0.54
1:B:164:GLU:O	2:B:289:HOH:O	2.18	0.54
1:C:217:HIS:HD2	1:C:219:GLY:N	1.98	0.54
2:A:452:HOH:O	1:C:128:ASN:HB2	2.09	0.53
1:A:217:HIS:CD2	1:A:219:GLY:H	2.25	0.52
1:C:9:LYS:HE3	1:C:112:ASP:OD2	2.11	0.51
1:B:123:THR:HG22	1:D:90:GLU:OE2	2.11	0.51
1:D:201:HIS:HD2	1:D:202:ASP:O	1.92	0.51
1:C:201:HIS:HD2	1:C:202:ASP:O	1.94	0.50
1:B:83:PHE:HB3	1:B:84:PRO:HA	1.94	0.50
1:C:21:HIS:HE1	1:C:47:GLY:O	1.94	0.49
2:B:500:HOH:O	1:D:150:ASP:HB3	2.12	0.49
1:C:64:CR8:N	1:C:64:CR8:H171	2.27	0.49
1:B:147:TYR:HH	1:B:174:ARG:NH2	2.07	0.48
1:A:143:THR:H	1:B:145:LYS:NZ	2.11	0.48
1:A:166:ASN:O	1:A:166:ASN:CG	2.50	0.48
1:C:36:GLY:O	1:C:213:HIS:HA	2.14	0.47
1:D:83:PHE:HB3	1:D:84:PRO:HA	1.96	0.47
1:B:145:LYS:HD2	1:B:191:PHE:CD1	2.49	0.47
1:C:170:ARG:HD3	1:D:147:TYR:CZ	2.49	0.47
1:A:64:CR8:H23	1:A:212:GLU:HB2	1.97	0.47
1:B:93:LEU:O	1:B:100:ILE:HD12	2.16	0.46
1:A:15:GLU:OE2	1:A:22:HIS:HE1	1.98	0.46
1:B:128:ASN:ND2	1:D:150:ASP:OD1	2.49	0.46
1:C:65:ASN:ND2	1:C:67:VAL:HG12	2.29	0.46
1:D:8:MET:HE3	1:D:8:MET:HB3	1.89	0.46
1:C:137:LEU:HA	1:C:137:LEU:HD23	1.83	0.45
1:B:201:HIS:HD2	1:B:202:ASP:O	1.99	0.45
1:C:65:ASN:ND2	1:C:67:VAL:CG1	2.80	0.45
1:A:83:PHE:HB3	1:A:84:PRO:HA	1.99	0.45
1:B:67:VAL:HG21	1:B:83:PHE:HE2	1.80	0.45
1:A:83:PHE:CE1	1:A:86:GLY:HA2	2.52	0.45
1:B:93:LEU:HD12	1:B:93:LEU:N	2.32	0.45
1:A:193:ASP:OD1	1:A:217:HIS:HE1	2.00	0.44
1:D:64:CR8:H23	1:D:212:GLU:HB2	1.98	0.44
1:B:100:ILE:HG23	1:B:100:ILE:O	2.16	0.44
1:B:212:GLU:HG2	1:B:213:HIS:N	2.32	0.44
1:D:43:GLU:HG3	1:D:207:LYS:HD3	2.00	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:147:TYR:HB3	1:C:189:TYR:CD1	2.53	0.43
1:C:143:THR:H	1:D:145:LYS:HZ1	1.66	0.43
1:D:171:CYS:HG	1:D:173:PHE:HE1	1.66	0.43
1:A:102:ILE:HG22	2:A:268:HOH:O	2.19	0.42
1:B:67:VAL:CG2	1:B:83:PHE:HE2	2.32	0.42
1:B:85:LYS:HG3	1:B:180:LYS:HG2	2.01	0.42
1:D:217:HIS:HD2	1:D:219:GLY:N	2.07	0.42
1:A:135:LYS:HE2	1:A:135:LYS:HB3	1.84	0.42
1:A:147:TYR:CZ	1:B:170:ARG:HD3	2.54	0.41
1:A:212:GLU:HG2	1:A:213:HIS:N	2.35	0.41
1:A:139:TRP:CZ3	1:A:161:LEU:HG	2.56	0.41
1:A:40:MET:HB2	1:A:64:CR8:H10	2.02	0.41
1:A:1:MET:HE2	1:A:3:ALA:CB	2.49	0.41
1:C:21:HIS:CE1	1:C:47:GLY:O	2.74	0.41
1:B:156:ASP:OD1	1:B:174:ARG:HG3	2.21	0.40
1:D:26:ASP:OD2	1:D:45:LYS:HG3	2.21	0.40
1:B:147:TYR:HB3	1:B:189:TYR:CD1	2.57	0.40
1:D:144:GLU:HB2	1:D:157:ILE:CD1	2.51	0.40
1:B:170:ARG:NH1	2:B:480:HOH:O	2.55	0.40
1:B:91:ARG:HD3	1:B:173:PHE:CD2	2.57	0.40
1:B:193:ASP:OD1	1:B:217:HIS:HE1	2.04	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	218/224 (97%)	213 (98%)	5 (2%)	0	100 100
1	B	219/224 (98%)	218 (100%)	1 (0%)	0	100 100
1	C	216/224 (96%)	213 (99%)	3 (1%)	0	100 100
1	D	216/224 (96%)	214 (99%)	2 (1%)	0	100 100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	869/896 (97%)	858 (99%)	11 (1%)	0	100   100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	189/191 (99%)	187 (99%)	2 (1%)	78   88
1	B	189/191 (99%)	188 (100%)	1 (0%)	91   96
1	C	187/191 (98%)	184 (98%)	3 (2%)	68   81
1	D	187/191 (98%)	184 (98%)	3 (2%)	68   81
All	All	752/764 (98%)	743 (99%)	9 (1%)	75   86

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	ARG
1	A	116	ASN
1	B	66	ARG
1	C	66	ARG
1	C	145	LYS
1	C	172	ASP
1	D	66	ARG
1	D	145	LYS
1	D	150	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	HIS
1	A	81	GLN
1	A	124	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	133	GLN
1	A	166	ASN
1	A	201	HIS
1	A	206	ASN
1	A	217	HIS
1	B	81	GLN
1	B	128	ASN
1	B	133	GLN
1	B	201	HIS
1	B	206	ASN
1	B	217	HIS
1	C	21	HIS
1	C	22	HIS
1	C	133	GLN
1	C	201	HIS
1	C	206	ASN
1	C	217	HIS
1	D	201	HIS
1	D	206	ASN
1	D	217	HIS

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CR8	A	64	1	18,27,28	2.28	8 (44%)	16,37,39	1.61	2 (12%)
1	CR8	B	64	1	18,27,28	2.33	8 (44%)	16,37,39	1.70	4 (25%)
1	CR8	C	64	1	18,27,28	2.21	6 (33%)	16,37,39	1.23	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CR8	D	64	1	18,27,28	2.05	6 (33%)	16,37,39	1.46	2 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CR8	A	64	1	-	0/8/25/26	0/3/3/3
1	CR8	B	64	1	-	0/8/25/26	0/3/3/3
1	CR8	C	64	1	-	0/8/25/26	0/3/3/3
1	CR8	D	64	1	-	0/8/25/26	0/3/3/3

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	64	CR8	C2-C1	-4.30	1.35	1.46
1	D	64	CR8	C2-C1	-3.82	1.36	1.46
1	A	64	CR8	C4-C1	-3.78	1.37	1.46
1	B	64	CR8	C4-C1	-3.75	1.37	1.46
1	A	64	CR8	C2-C1	-3.70	1.37	1.46
1	C	64	CR8	C2-C1	-3.57	1.37	1.46
1	C	64	CR8	C17-N13	-3.55	1.42	1.49
1	D	64	CR8	O25-C12	-3.53	1.22	1.32
1	A	64	CR8	O25-C12	-3.36	1.22	1.32
1	B	64	CR8	O3-C1	-3.33	1.14	1.24
1	C	64	CR8	O25-C12	-3.26	1.22	1.32
1	A	64	CR8	O3-C1	-3.23	1.14	1.24
1	C	64	CR8	C4-C1	-3.17	1.38	1.46
1	A	64	CR8	C17-N13	-3.14	1.43	1.49
1	B	64	CR8	O25-C12	-3.10	1.23	1.32
1	B	64	CR8	C17-N13	-2.80	1.44	1.49
1	D	64	CR8	C4-C1	-2.71	1.39	1.46
1	C	64	CR8	O3-C1	-2.65	1.16	1.24
1	B	64	CR8	C17-C	-2.50	1.43	1.48
1	D	64	CR8	O3-C1	-2.10	1.17	1.24
1	D	64	CR8	C23-N11	-2.01	1.29	1.35
1	A	64	CR8	C23-N11	-2.00	1.29	1.35
1	B	64	CR8	C9-C8	2.08	1.48	1.40
1	A	64	CR8	C9-C8	2.09	1.48	1.40
1	D	64	CR8	C8-C7	3.93	1.46	1.36
1	A	64	CR8	C8-C7	4.44	1.47	1.36

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	64	CR8	C8-C7	4.49	1.47	1.36
1	C	64	CR8	C8-C7	4.97	1.49	1.36

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	CR8	O19-C-C17	-5.00	110.61	126.26
1	D	64	CR8	O3-C1-C2	-3.67	115.63	121.54
1	C	64	CR8	C4-C5-C7	-2.54	119.79	121.95
1	B	64	CR8	O3-C1-C4	-2.36	117.74	121.54
1	B	64	CR8	C6-C2-C1	-2.21	118.05	121.21
1	B	64	CR8	C2-C6-C7	-2.17	120.11	121.95
1	A	64	CR8	C4-C1-C2	2.13	120.59	116.70
1	D	64	CR8	O3-C1-C4	2.45	125.48	121.54
1	B	64	CR8	C4-C1-C2	3.48	123.05	116.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	64	CR8	2	0
1	C	64	CR8	1	0
1	D	64	CR8	1	0

## 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	220/224 (98%)	-0.70	1 (0%) 90 90	5, 13, 37, 82	3 (1%)
1	B	221/224 (98%)	-0.72	0 100 100	5, 13, 34, 58	4 (1%)
1	C	218/224 (97%)	-0.80	0 100 100	6, 14, 33, 66	4 (1%)
1	D	218/224 (97%)	-0.74	0 100 100	7, 17, 41, 56	4 (1%)
All	All	877/896 (97%)	-0.74	1 (0%) 95 95	5, 14, 36, 82	15 (1%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	5.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CR8	D	64	25/26	0.95	0.12	-	3,11,16,22	0
1	CR8	C	64	25/26	0.95	0.10	-	3,7,14,17	0
1	CR8	B	64	25/26	0.96	0.11	-	2,8,13,15	0
1	CR8	A	64	25/26	0.96	0.09	-	2,6,12,13	0

### 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.